

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
PCNL	730	1,180.1	1,174	1,305.8	1,478	1,153.2	1,570	1,081.1	2,135	1,115.6
Age	-	-	-	-	-	-	-	-	-	-
< 18	16	989.5	37	1,388.4	40	1,005.0	35	690.5	54	789.9
18-24	16	584.6	39	990.9	51	950.6	42	691.2	68	812.7
25-34	61	761.3	96	786.4	136	793.2	125	612.2	216	774.9
35-39	32	526.4	98	1,054.6	107	819.0	116	761.4	172	839.8
40-44	93	1,266.2	147	1,261.4	152	923.6	141	744.1	245	972.1
45-54	235	1,204.8	345	1,205.9	458	1,115.1	469	1,004.8	598	983.1
55-64	277	1,672.6	412	1,912.3	534	1,718.3	642	1,957.1	782	1,871.2
Region	-	-	-	-	-	-	-	-	-	-
Northeast	45	862.2	93	1,076.8	131	1,164.1	160	1,148.2	238	1,192.4
South	223	720.3	264	662.0	345	533.3	313	425.1	627	621.2
Midwest	377	2,049.7	532	2,393.2	694	2,365.5	731	2,220.7	1,039	2,105.5
West	85	1,166.6	285	1,487.2	308	1,346.0	366	1,479.6	231	1,092.9
Gender	-	-	-	-	-	-	-	-	-	-
Female	374	1,529.6	584	1,581.5	751	1,405.3	832	1,346.6	1,140	1,384.9
Male	356	951.7	590	1,113.6	727	972.9	738	884.5	995	912.3

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
ESWL	10,452	16,897.0	14,938	16,614.4	21,492	16,769.3	23,901	16,457.9	30,128	15,742.2
Age	-	-	-	-	-	-	-	-	-	-
< 18	114	7,050.1	177	6,641.7	234	5,879.4	247	4,872.8	321	4,695.7
18-24	311	11,362.8	435	11,051.8	590	10,997.2	665	10,944.7	884	10,565.3
25-34	1,197	14,938.2	1,664	13,631.5	2,251	13,128.4	2,598	12,724.7	3,477	12,473.1
35-39	1,005	16,532.3	1,380	14,849.9	1,903	14,565.6	2,090	13,717.5	2,855	13,939.1
40-44	1,214	16,528.3	1,855	15,917.3	2,573	15,633.7	2,847	15,025.3	3,559	14,121.9
45-54	3,374	17,298.1	4,904	17,140.9	6,920	16,848.5	7,687	16,468.8	9,339	15,352.9
55-64	3,237	19,545.9	4,523	20,993.3	7,021	22,592.3	7,767	23,677.7	9,693	23,193.4
Region	-	-	-	-	-	-	-	-	-	-
Northeast	781	14,964.6	1,386	16,047.2	1,770	15,729.1	2,084	14,955.1	2,912	14,589.9
South	2,838	9,167.0	3,321	8,327.7	4,399	6,800.1	5,026	6,825.5	7,476	7,406.3
Midwest	5,831	31,702.3	7,587	34,129.6	11,937	40,687.8	13,250	40,252.8	16,755	33,953.4
West	1,002	13,752.4	2,644	13,796.7	3,386	14,797.7	3,541	14,314.6	2,985	14,122.2
Gender	-	-	-	-	-	-	-	-	-	-
Female	3,890	15,909.4	5,923	16,039.3	8,448	15,808.1	9,896	16,016.3	12,414	15,080.7
Male	6,562	17,542.6	9,015	17,015.2	13,044	17,456.7	14,005	16,784.9	17,714	16,241.4

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Open Stone Surgery	262	423.6	319	354.8	382	298.1	367	252.7	516	269.6
Age	-	-	-	-	-	-	-	-	-	-
< 18	25	1,546.1	38	1,425.9	48	1,206.0	34	670.7	51	746.1
18-24	16	584.6	19	482.7	15	279.6	31	510.2	33	394.4
25-34	18	224.6	33	270.3	45	262.5	29	142.0	77	276.2
35-39	26	427.7	36	387.4	25	191.4	23	151.0	41	200.2
40-44	29	394.8	27	231.7	38	230.9	52	274.4	59	234.1
45-54	79	405.0	85	297.1	108	263.0	94	201.4	148	243.3
55-64	69	416.6	81	376.0	103	331.4	104	317.0	107	256.0
Region	-	-	-	-	-	-	-	-	-	-
Northeast	29	555.7	66	764.2	84	746.5	66	473.6	67	335.7
South	81	261.6	68	170.5	88	136.0	75	101.9	141	139.7
Midwest	125	679.6	132	593.8	146	497.6	165	501.3	248	502.6
West	27	370.6	53	276.6	64	279.7	61	246.6	60	283.9
Gender	-	-	-	-	-	-	-	-	-	-
Female	129	527.6	167	452.2	195	364.9	189	305.9	264	320.7
Male	133	355.6	152	286.9	187	250.3	178	213.3	252	231.1

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Uteroscopy	8,421	13,613.7	11,692	13,004.1	15,748	12,287.5	18,007	12,399.4	23,412	12,233.0
Age	-	-	-	-	-	-	-	-	-	-
< 18	134	8,287.0	210	7,879.9	255	6,407.0	301	5,938.1	365	5,339.4
18-24	441	16,112.5	579	14,710.4	651	12,134.2	776	12,771.6	1,081	12,919.8
25-34	1,140	14,226.9	1,569	12,853.3	2,011	11,728.7	2,300	11,265.1	3,066	10,998.7
35-39	902	14,838.0	1,155	12,428.7	1,514	11,588.2	1,773	11,636.9	2,353	11,488.1
40-44	977	13,301.6	1,511	12,965.5	1,926	11,702.5	2,076	10,956.3	2,858	11,340.4
45-54	2,548	13,063.3	3,446	12,044.7	4,725	11,504.2	5,369	11,502.7	7,010	11,524.1
55-64	2,279	13,761.2	3,222	14,954.7	4,666	15,014.3	5,412	16,498.5	6,679	15,981.5
Region	-	-	-	-	-	-	-	-	-	-
Northeast	535	10,251.0	881	10,200.3	1,159	10,299.5	1,375	9,867.2	1,887	9,454.4
South	2,460	7,946.0	2,935	7,359.8	3,699	5,718.0	4,239	5,756.7	6,363	6,303.7
Midwest	4,749	25,819.6	6,039	27,166.0	8,623	29,391.9	9,835	29,878.2	12,886	26,113.0
West	677	9,291.8	1,837	9,585.7	2,267	9,907.4	2,558	10,340.8	2,276	10,767.8
Gender	-	-	-	-	-	-	-	-	-	-
Female	3,323	13,590.4	4,682	12,678.7	6,292	11,773.7	7,487	12,117.4	9,785	11,887.0
Male	5,098	13,628.8	7,010	13,230.9	9,456	12,654.9	10,520	12,608.2	13,627	12,494.2

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Laparoscopic Removal	1	1.6	5	5.6	6	4.7	8	5.5	3	1.6
Age	-	-	-	-	-	-	-	-	-	-
< 18	0	0.0	0	0.0	3	75.4	0	0.0	1	14.6
18-24	0	0.0	1	25.4	0	0.0	1	16.5	1	12.0
25-34	0	0.0	0	0.0	0	0.0	2	9.8	0	0.0
35-39	0	0.0	0	0.0	1	7.7	0	0.0	0	0.0
40-44	0	0.0	0	0.0	1	6.1	2	10.6	1	4.0
45-54	0	0.0	2	7.0	0	0.0	2	4.3	0	0.0
55-64	1	6.0	2	9.3	1	3.2	1	3.0	0	0.0
Region	-	-	-	-	-	-	-	-	-	-
Northeast	0	0.0	1	11.6	0	0.0	0	0.0	0	0.0
South	0	0.0	3	7.5	2	3.1	5	6.8	2	2.0
Midwest	0	0.0	1	4.5	4	13.6	3	9.1	1	2.0
West	1	13.7	0	0.0	0	0.0	0	0.0	0	0.0
Gender	-	-	-	-	-	-	-	-	-	-
Female	0	0.0	4	10.8	3	5.6	5	8.1	2	2.4
Male	1	2.7	1	1.9	3	4.0	3	3.6	1	0.9

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002		2003		2004		2005		2006	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
All Procedures	19,866	32,116.0	28,128	31,284.6	39,106	30,512.7	43,853	30,196.6	56,194	29,361.9
Age	-	-	-	-	-	-	-	-	-	-
< 18	289	17,872.6	462	17,335.8	580	14,572.9	617	12,172.0	792	11,585.7
18-24	784	28,644.5	1,073	27,261.2	1,307	24,361.6	1,515	24,934.2	2,067	24,704.2
25-34	2,416	30,151.0	3,362	27,541.6	4,443	25,912.7	5,054	24,753.9	6,836	24,522.9
35-39	1,965	32,324.4	2,669	28,720.5	3,550	27,171.8	4,002	26,266.7	5,421	26,467.1
40-44	2,313	31,490.8	3,540	30,375.8	4,690	28,496.8	5,118	27,010.8	6,722	26,672.5
45-54	6,236	31,971.3	8,782	30,695.6	12,211	29,730.7	13,621	29,182.0	17,095	28,103.4
55-64	5,863	35,402.5	8,240	38,245.5	12,325	39,659.6	13,926	42,453.4	17,261	41,302.2
Region	-	-	-	-	-	-	-	-	-	-
Northeast	1,390	26,633.5	2,427	28,100.0	3,144	27,939.2	3,685	26,444.2	5,104	25,572.4
South	5,602	18,094.9	6,591	16,527.5	8,533	13,190.6	9,658	13,115.9	14,609	14,472.8
Midwest	11,082	60,251.2	14,291	64,287.0	21,404	72,956.6	23,984	72,862.0	30,929	62,676.6
West	1,792	24,595.1	4,819	25,146.1	6,025	26,330.7	6,526	26,381.5	5,552	26,266.7
Gender	-	-	-	-	-	-	-	-	-	-
Female	7,716	31,557.0	11,360	30,762.6	15,689	29,357.6	18,409	29,794.3	23,605	28,675.7
Male	12,150	32,481.4	16,768	31,648.5	23,417	31,338.8	25,444	30,494.5	32,589	29,879.8

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
PCNL	2,289	1,064.0	3,248	1,090.8	3,814	1,170.0	3,924	1,251.5	4,647	1,226.4
Age	-	-	-	-	-	-	-	-	-	-
< 18	68	852.7	84	787.9	88	743.1	94	816.6	115	819.4
18-24	65	682.1	93	706.3	96	647.2	126	877.4	140	680.8
25-34	192	595.8	315	718.2	332	667.8	319	659.2	437	781.8
35-39	176	740.5	243	737.5	274	751.1	296	823.2	287	688.8
40-44	241	853.6	310	795.9	319	746.6	375	897.9	415	848.2
45-54	662	957.8	954	1,007.4	1,107	1,058.8	1,197	1,179.6	1,321	1,096.0
55-64	885	1,999.1	1,249	1,967.7	1,598	2,426.9	1,517	2,524.1	1,932	2,499.7
Region	-	-	-	-	-	-	-	-	-	-
Northeast	222	1,036.9	568	1,076.9	589	1,159.8	632	1,209.2	941	1,194.1
South	707	636.5	839	633.6	1,046	713.2	1,087	772.4	1,245	769.5
Midwest	1,059	1,841.4	1,362	1,828.2	1,704	2,042.5	1,697	2,287.6	1,801	2,098.3
West	301	1,198.2	479	1,256.9	475	1,053.2	508	1,095.5	660	1,257.6
Gender	-	-	-	-	-	-	-	-	-	-
Female	1,177	1,257.2	1,745	1,324.5	2,028	1,385.7	2,101	1,483.0	2,551	1,482.1
Male	1,112	915.2	1,503	905.3	1,786	994.2	1,823	1,060.6	2,096	1,013.6

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
ESWL	31,657	14,715.8	43,202	14,509.0	43,604	13,376.0	42,263	13,478.6	48,093	12,692.5
Age	-	-	-	-	-	-	-	-	-	-
< 18	301	3,774.3	410	3,845.8	349	2,947.1	337	2,927.6	357	2,543.6
18-24	958	10,053.5	1,194	9,067.4	1,339	9,026.6	1,178	8,203.3	1,527	7,425.6
25-34	3,495	10,844.6	4,751	10,832.2	4,784	9,623.2	4,521	9,342.6	5,009	8,960.8
35-39	3,178	13,371.5	4,234	12,850.2	4,295	11,773.6	4,064	11,302.4	4,256	10,215.1
40-44	3,708	13,133.6	5,024	12,899.3	5,077	11,883.0	4,848	11,608.6	5,534	11,311.2
45-54	9,805	14,185.7	13,422	14,173.0	13,392	12,809.2	12,959	12,770.8	14,405	11,951.3
55-64	10,212	23,067.0	14,167	22,319.4	14,368	21,821.3	14,356	23,886.5	17,005	22,001.6
Region	-	-	-	-	-	-	-	-	-	-
Northeast	2,766	12,919.8	6,609	12,530.8	6,139	12,088.0	6,102	11,674.7	9,220	11,699.9
South	8,228	7,407.2	10,763	8,128.7	11,267	7,681.7	10,519	7,474.3	11,668	7,211.6
Midwest	17,601	30,604.6	20,975	28,154.0	20,934	25,092.0	20,550	27,701.8	21,404	24,937.7
West	3,062	12,189.0	4,855	12,739.4	5,264	11,671.8	5,092	10,981.0	5,801	11,053.7
Gender	-	-	-	-	-	-	-	-	-	-
Female	13,347	14,256.6	18,279	13,874.7	18,892	12,908.3	18,585	13,118.3	21,105	12,261.9
Male	18,310	15,069.7	24,923	15,012.2	24,712	13,756.9	23,678	13,775.6	26,988	13,051.0

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Open Stone Surgery	521	242.2	686	230.4	439	134.7	381	121.5	413	109.0
Age	-	-	-	-	-	-	-	-	-	-
< 18	69	865.2	85	797.3	55	464.4	55	477.8	63	448.9
18-24	25	262.4	34	258.2	25	168.5	19	132.3	24	116.7
25-34	58	180.0	95	216.6	53	106.6	38	78.5	52	93.0
35-39	40	168.3	47	142.6	32	87.7	18	50.1	26	62.4
40-44	60	212.5	74	190.0	35	81.9	23	55.1	33	67.5
45-54	141	204.0	150	158.4	125	119.6	114	112.3	98	81.3
55-64	128	289.1	201	316.7	114	173.1	114	189.7	117	151.4
Region	-	-	-	-	-	-	-	-	-	-
Northeast	71	331.6	117	221.8	87	171.3	76	145.4	110	139.6
South	155	139.5	171	129.1	111	75.7	76	54.0	91	56.2
Midwest	232	403.4	309	414.8	173	207.4	176	237.3	143	166.6
West	63	250.8	89	233.5	68	150.8	53	114.3	69	131.5
Gender	-	-	-	-	-	-	-	-	-	-
Female	280	299.1	342	259.6	219	149.6	200	141.2	223	129.6
Male	241	198.4	344	207.2	220	122.5	181	105.3	190	91.9

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Uteroscopy	25,621	11,910.0	35,548	11,938.4	38,345	11,762.7	38,493	12,276.3	46,178	12,187.1
Age	-	-	-	-	-	-	-	-	-	-
< 18	372	4,664.6	542	5,084.0	559	4,720.5	530	4,604.3	581	4,139.7
18-24	1,167	12,246.8	1,462	11,102.7	1,577	10,631.0	1,553	10,814.8	2,204	10,717.8
25-34	3,269	10,143.4	4,560	10,396.7	4,783	9,621.2	4,627	9,561.7	5,318	9,513.6
35-39	2,634	11,082.6	3,629	11,014.0	3,840	10,526.3	3,912	10,879.7	4,131	9,915.0
40-44	2,988	10,583.4	4,046	10,388.2	4,255	9,959.0	4,325	10,356.3	5,335	10,904.4
45-54	7,537	10,904.4	10,617	11,211.1	11,346	10,852.2	11,495	11,328.0	13,491	11,193.0
55-64	7,654	17,289.0	10,692	16,844.7	11,985	18,202.1	12,051	20,051.2	15,118	19,560.1
Region	-	-	-	-	-	-	-	-	-	-
Northeast	2,008	9,379.2	4,539	8,606.0	4,630	9,116.7	4,967	9,503.1	7,903	10,028.7
South	7,439	6,696.9	9,893	7,471.6	10,973	7,481.3	10,513	7,470.1	12,119	7,490.4
Midwest	13,630	23,699.8	17,047	22,881.6	18,172	21,781.4	18,148	24,463.8	20,748	24,173.4
West	2,544	10,127.0	4,069	10,677.0	4,570	10,133.0	4,865	10,491.5	5,408	10,304.9
Gender	-	-	-	-	-	-	-	-	-	-
Female	11,034	11,785.9	15,635	11,867.8	16,996	11,612.9	17,064	12,044.7	20,814	12,092.8
Male	14,587	12,005.6	19,913	11,994.5	21,349	11,884.8	21,429	12,467.1	25,364	12,265.6

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Laparoscopic Removal	7	3.3	9	3.0	11	3.4	12	3.8	15	4.0
Age	-	-	-	-	-	-	-	-	-	-
< 18	0	0.0	0	0.0	0	0.0	2	17.4	1	7.1
18-24	1	10.5	2	15.2	0	0.0	0	0.0	2	9.7
25-34	0	0.0	0	0.0	2	4.0	2	4.1	1	1.8
35-39	0	0.0	3	9.1	2	5.5	1	2.8	0	0.0
40-44	0	0.0	0	0.0	1	2.3	1	2.4	0	0.0
45-54	2	2.9	4	4.2	1	1.0	3	3.0	6	5.0
55-64	4	9.0	0	0.0	5	7.6	3	5.0	5	6.5
Region	-	-	-	-	-	-	-	-	-	-
Northeast	3	14.0	0	0.0	0	0.0	1	1.9	3	3.8
South	0	0.0	2	1.5	6	4.1	1	0.7	3	1.9
Midwest	3	5.2	6	8.1	5	6.0	7	9.4	6	7.0
West	1	4.0	1	2.6	0	0.0	3	6.5	3	5.7
Gender	-	-	-	-	-	-	-	-	-	-
Female	2	2.1	6	4.6	4	2.7	6	4.2	6	3.5
Male	5	4.1	3	1.8	7	3.9	6	3.5	9	4.4

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.

Table 12 - Surgical Procedures among Privately Insured Adults with A Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007		2008		2009		2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
All Procedures	60,095	27,935.3	82,693	27,771.6	86,213	26,446.7	85,073	27,131.7	99,346	26,219.0
Age	-	-	-	-	-	-	-	-	-	-
< 18	810	10,156.7	1,121	10,515.0	1,051	8,875.2	1,018	8,843.7	1,117	7,958.7
18-24	2,216	23,255.3	2,785	21,149.8	3,037	20,473.2	2,876	20,027.9	3,897	18,950.6
25-34	7,014	21,763.7	9,721	22,163.7	9,954	20,022.9	9,507	19,646.2	10,817	19,351.0
35-39	6,028	25,362.9	8,156	24,753.4	8,443	23,144.2	8,291	23,058.1	8,700	20,881.3
40-44	6,997	24,783.1	9,454	24,273.4	9,687	22,672.9	9,572	22,920.4	11,317	23,131.3
45-54	18,147	26,254.7	25,147	26,554.1	25,971	24,840.7	25,768	25,393.7	29,321	24,326.5
55-64	18,883	42,653.2	26,309	41,448.5	28,070	42,631.1	28,041	46,656.5	34,177	44,219.2
Region	-	-	-	-	-	-	-	-	-	-
Northeast	5,070	23,681.6	11,833	22,435.6	11,445	22,535.7	11,778	22,534.3	18,177	23,066.1
South	16,529	14,880.1	21,668	16,364.6	23,403	15,955.9	22,196	15,771.5	25,126	15,529.6
Midwest	32,525	56,554.4	39,699	53,286.5	40,988	49,129.2	40,578	54,699.9	44,102	51,383.0
West	5,971	23,769.0	9,493	24,909.5	10,377	23,008.9	10,521	22,688.7	11,941	22,753.4
Gender	-	-	-	-	-	-	-	-	-	-
Female	25,840	27,600.9	36,007	27,331.2	38,139	26,059.2	37,956	26,791.5	44,699	25,969.8
Male	34,255	28,193.0	46,686	28,121.0	48,074	26,762.3	47,117	27,412.1	54,647	26,426.5

SOURCE: Truven Health MarketScan®, 2002-2011.

NOTE: X represents the number of privately insured adults with a primary diagnosis of urolithiasis.

NOTE: Rate is per 100,000 privately insured adults with a primary diagnosis of urolithiasis.