

Table 7b - Other Ambulatory Visit Rates for Medicare Beneficiaries with a Primary Diagnosis of Urolithiasis, 2002 - 2006

	2002			2003			2004			2005			2006		
	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator
<b>Total</b>	43,720	9,647	-	46,320	9,539	-	51,060	9,623	-	53,560	9,453	-	53,100	9,078	-
<b>Age</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69	14,100	11,529	R	15,840	11,704	R	16,920	11,491	R	17,800	11,320	R	17,200	10,531	R
70-74	12,100	10,168	*	12,200	9,741	*	13,980	10,333	*	15,380	10,839	*	15,020	10,142	*
75-79	10,260	9,801	*	10,160	9,340	*	11,100	9,348	*	10,620	8,541	*	10,740	8,554	*
80-84	4,700	7,100	*	5,420	7,602	*	5,760	7,368	*	6,260	7,202	*	6,620	7,521	*
85+	2,560	6,244	*	2,700	6,008	*	3,300	6,453	*	3,500	6,228	*	3,520	5,875	*
<b>Race</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
White	38,840	9,629	R	41,020	9,534	R	44,740	9,544	R	47,600	9,500	R	48,700	9,389	R
Black	2,740	10,604	*	2,780	9,816	*	3,580	11,526	*	3,300	9,627	*	2,240	6,470	*
Hispanic	460	7,012	*	740	9,788	*	820	9,214	*	580	6,532	*	680	6,827	*
Asian	380	7,480	*	320	6,038	*	340	5,060	*	520	6,633	*	540	6,553	*
Other	1,300	10,552	*	1,460	10,296	*	1,580	10,422	*	1,560	10,744	*	940	7,015	*
<b>Region</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northeast	9,520	8,583	*	9,940	8,609	*	11,200	9,149	*	10,120	7,826	*	11,000	8,135	*
South	15,420	8,714	*	16,500	8,616	*	17,060	8,114	*	17,740	7,890	*	19,020	8,103	*
Midwest	12,740	12,471	R	13,900	12,797	R	15,820	13,135	R	18,000	13,870	R	16,080	12,204	R
West	6,040	9,566	*	5,980	8,540	*	6,980	9,009	*	7,700	9,315	*	7,000	8,411	*
<b>Gender</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female	16,900	9,225	R	18,480	9,445	R	20,280	9,394	R	20,120	8,743	R	21,560	9,021	R
Male	26,820	9,934	*	27,840	9,602	*	30,780	9,780	*	33,440	9,939	*	31,540	9,118	*

SOURCE: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

NOTE: X represents the number of Medicare beneficiaries diagnosed with any diagnosis of urolithiasis

NOTE: Unweighted counts have been multiplied by 20 to arrive at counts in table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 adult Medicare beneficiaries with a primary diagnosis of urolithiasis.

NOTE: \* indicates that the rate is significantly different from reference group (R)

NOTE: R indicates the reference group

Table 7b - Other Ambulatory Visit Rates for Medicare Beneficiaries with a Primary Diagnosis of Urolithiasis, 2007 - 2011

	2007			2008			2009			2010			2011		
	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator
<b>Total</b>	53,080	9,162	-	62,200	10,147	-	65,220	10,093	-	72,920	10,706	-	79,840	11,063	-
<b>Age</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69	17,640	10,894	R	22,320	12,875	R	21,760	11,842	R	25,120	12,878	R	28,800	13,682	R
70-74	13,260	9,264	*	16,920	11,030	*	17,560	10,733	*	19,400	11,303	*	21,260	11,724	*
75-79	11,120	9,023	*	11,240	9,049	*	12,280	9,422	*	14,820	10,840	*	15,120	10,606	*
80-84	7,140	7,904	*	7,180	7,534	*	8,740	8,944	*	8,740	8,738	*	9,100	8,695	*
85+	3,920	6,454	*	4,540	6,809	*	4,880	6,893	*	4,840	6,231	*	5,560	6,728	*
<b>Race</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
White	48,400	9,376	R	55,940	10,274	R	59,260	10,317	R	66,820	11,045	R	72,860	11,363	R
Black	2,300	7,302	*	3,140	8,982	*	2,900	7,425	*	3,360	8,231	*	3,700	8,669	*
Hispanic	680	7,188	*	720	8,392	*	560	6,250	*	820	7,854	*	740	6,715	*
Asian	500	5,669	*	540	5,477	*	720	8,090	*	780	7,500	*	1,040	9,665	*
Other	1,200	8,982	*	1,860	12,318	*	1,780	11,946	*	1,140	7,873	*	1,500	9,340	*
<b>Region</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northeast	11,000	8,158	*	12,820	9,299	*	13,520	9,472	*	14,740	9,951	*	16,640	10,749	*
South	17,320	7,488	*	23,040	9,258	*	22,620	8,529	*	24,860	8,854	*	29,300	9,723	*
Midwest	17,140	13,060	R	17,680	12,873	R	19,060	13,375	R	22,200	14,737	R	22,260	14,618	R
West	7,620	9,293	*	8,660	9,744	*	10,020	10,464	*	11,120	10,947	*	11,640	10,275	*
<b>Gender</b>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female	20,440	8,692	R	23,760	9,549	R	25,880	9,972	R	27,080	9,899	R	30,600	10,592	R
Male	32,640	9,482	*	38,440	10,556	*	39,340	10,174	*	45,840	11,247	*	49,240	11,377	*

SOURCE: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

NOTE: X represents the number of Medicare beneficiaries diagnosed with any diagnosis of urolithiasis

NOTE: Unweighted counts have been multiplied by 20 to arrive at counts in table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 adult Medicare beneficiaries with a primary diagnosis of urolithiasis.

NOTE: \* indicates that the rate is significantly different from reference group (R)

NOTE: R indicates the reference group