

Table 8 - Hospitalization Rate for Medicare Beneficiaries with Urolithiasis, 2002 - 2006

	2002			2003			2004			2005			2006		
	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator
Total	48,440	10,688	-	50,740	10,449	-	51,900	9,781	-	51,460	9,082	-	51,440	8,794	-
Age	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69	11,820	9,663	R	12,240	9,044	R	12,140	8,245	R	12,220	7,771	R	12,320	7,544	R
70-74	11,420	9,597	*	12,360	9,869	*	12,440	9,194	*	12,120	8,541	*	11,240	7,590	*
75-79	11,620	11,100	*	11,000	10,112	*	11,420	9,618	*	10,880	8,750	*	11,220	8,936	*
80-84	7,420	11,208	*	8,520	11,950	*	8,760	11,205	*	9,240	10,630	*	9,240	10,498	*
85+	6,160	15,024	*	6,620	14,731	*	7,140	13,962	*	7,000	12,456	*	7,420	12,383	*
Race	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
White	43,100	10,685	R	45,820	10,650	R	46,220	9,860	R	46,020	9,184	R	45,800	8,830	R
Black	3,360	13,003	*	2,700	9,534	*	3,400	10,947	*	3,440	10,035	*	3,300	9,532	*
Hispanic	740	11,246	*	540	7,143	*	740	8,315	*	680	7,658	*	780	7,831	*
Asian	320	6,299	*	480	9,057	*	200	2,976	*	420	5,357	*	600	7,282	*
Other	920	7,468	*	1,200	8,463	*	1,340	8,839	*	900	6,190	*	960	7,164	*
Region	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northeast	9,840	8,870	*	10,900	9,441	*	10,280	8,397	*	10,660	8,242	*	11,620	8,593	*
South	19,460	10,997	*	20,580	10,747	*	20,460	9,731	*	20,040	8,913	*	19,020	8,103	*
Midwest	13,340	13,058	R	13,160	12,116	R	14,460	12,006	R	14,380	11,080	R	14,760	11,202	R
West	5,800	9,186	*	6,100	8,712	*	6,700	8,647	*	6,380	7,718	*	6,040	7,258	*
Gender	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female	20,660	11,276	R	21,780	11,132	R	23,300	10,793	R	23,660	10,281	R	23,900	10,000	R
Male	27,780	10,290	*	28,960	9,988	*	28,600	9,087	*	27,800	8,262	*	27,540	7,961	*

SOURCE: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

NOTE: X represents the number of Medicare beneficiaries diagnosed with any diagnosis of urolithiasis

NOTE: Unweighted counts have been multiplied by 20 to arrive at counts in table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 adult Medicare beneficiaries with a primary diagnosis of urolithiasis.

NOTE: * indicates that the rate is significantly different from reference group (R)

NOTE: R indicates the reference group

Table 8 - Hospitalization Rate for Medicare Beneficiaries with Urolithiasis, 2007 - 2011

	2007			2008			2009			2010			2011		
	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator	Count of X	Rate per 100,000 Patients	95% Significance Indicator
Total	47,460	8,192	-	50,400	8,222	-	48,660	7,530	-	53,380	7,837	-	62,020	8,594	-
Age	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65-69	11,960	7,386	R	12,860	7,418	R	11,520	6,269	R	12,840	6,583	R	15,120	7,183	R
70-74	10,160	7,098	*	11,500	7,496	*	11,300	6,907	*	11,780	6,863	*	13,820	7,621	*
75-79	9,480	7,692	*	9,520	7,664	*	9,420	7,227	*	10,880	7,958	*	12,300	8,628	*
80-84	8,640	9,564	*	9,520	9,990	*	8,560	8,760	*	9,480	9,478	*	9,480	9,058	*
85+	7,220	11,887	*	7,000	10,498	*	7,860	11,102	*	8,400	10,814	*	11,300	13,674	*
Race	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
White	42,300	8,194	R	45,100	8,283	R	42,660	7,427	R	47,100	7,785	R	55,100	8,594	R
Black	3,120	9,905	*	3,400	9,725	*	4,120	10,548	*	3,880	9,505	*	4,300	10,075	*
Hispanic	780	8,245	*	520	6,061	*	700	7,813	*	720	6,897	*	800	7,260	*
Asian	420	4,762	*	520	5,274	*	460	5,169	*	640	6,154	*	660	6,134	*
Other	840	6,287	*	860	5,695	*	720	4,832	*	1,040	7,182	*	1,160	7,223	*
Region	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Northeast	10,940	8,113	*	11,040	8,007	*	10,500	7,356	*	11,600	7,832	*	13,100	8,463	*
South	16,860	7,289	*	19,020	7,642	*	18,000	6,787	*	19,760	7,038	*	24,100	7,998	*
Midwest	13,620	10,378	R	13,760	10,019	R	13,520	9,488	R	14,700	9,758	R	15,300	10,047	R
West	6,040	7,366	*	6,580	7,403	*	6,640	6,934	*	7,320	7,206	*	9,520	8,404	*
Gender	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female	21,260	9,041	R	23,100	9,283	R	22,040	8,492	R	24,900	9,102	R	29,100	10,073	R
Male	26,200	7,611	*	27,300	7,497	*	26,620	6,884	*	28,480	6,988	*	32,920	7,606	*

SOURCE: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

NOTE: X represents the number of Medicare beneficiaries diagnosed with any diagnosis of urolithiasis

NOTE: Unweighted counts have been multiplied by 20 to arrive at counts in table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 adult Medicare beneficiaries with a primary diagnosis of urolithiasis.

NOTE: * indicates that the rate is significantly different from reference group (R)

NOTE: R indicates the reference group