

Table 1 - Prevalence of Kidney Stones in Individuals Ages 18 and Older, 2007-2012

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)	Refused to Answer/Don't Know (N)	Refused to Answer/Don't Know (%)
	219,353,204	19,225,012	8.76%	8.12	9.44	199,625,322	91.01%	502,871	0.23%
<b>Gender</b>	-	-	-	-	-	-	-	-	-
Male	105,550,501	10,517,819	9.96%	8.98	11.01	94,710,731	89.73%	321,951	0.31%
Female	113,802,703	8,707,192	7.65%	6.82	8.51	104,914,590	92.19%	180,920	0.16%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-
18-24	13,181,779	517,927	3.93%	2.51	5.36	12,648,290	95.95%	15,561	0.12%
25-34	36,140,335	1,867,949	5.17%	4.15	6.19	34,256,345	94.79%	16,041	0.04%
35-44	41,347,113	3,206,346	7.75%	6.29	9.29	37,959,238	91.81%	181,529	0.44%
45-54	48,718,584	4,549,844	9.34%	7.87	10.85	44,070,298	90.46%	98,441	0.20%
55-59	20,076,793	1,887,311	9.40%	7.54	11.30	18,153,424	90.42%	36,059	0.18%
60-64	17,853,013	2,146,007	12.02%	9.78	14.34	15,649,754	87.66%	57,252	0.32%
65-69	13,486,545	1,486,413	11.02%	8.72	13.38	11,968,170	88.74%	31,962	0.24%
70-74	10,311,146	1,162,716	11.28%	7.59	15.02	9,122,990	88.48%	25,442	0.25%
75-79	7,879,650	928,032	11.78%	8.97	14.62	6,940,371	88.08%	11,246	0.14%
80+	10,358,248	1,472,467	14.22%	11.06	17.45	8,856,443	85.50%	29,339	0.28%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-
Mexican American	18,027,042	1,154,898	6.41%	5.27	7.59	16,811,694	93.26%	60,452	0.34%
Other Hispanic	12,074,051	994,475	8.24%	6.65	9.89	11,029,629	91.35%	49,948	0.41%
Non-Hispanic White	148,950,560	15,003,061	10.07%	9.22	10.96	133,676,253	89.75%	271,246	0.18%
Non-Hispanic Black	24,987,691	1,100,233	4.40%	3.66	5.16	23,852,354	95.46%	35,105	0.14%
Other race - including multi-racial	15,313,860	972,346	6.35%	4.84	7.93	14,255,392	93.09%	86,121	0.56%
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-
Poverty income ratio less than 1	31,655,073	2,386,576	7.54%	6.39	8.75	29,138,589	92.05%	129,907	0.41%
Poverty income ratio 1 or greater	171,608,969	15,352,905	8.95%	8.25	9.68	155,915,368	90.86%	340,696	0.20%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-
Normal: less than 25	68,544,152	4,255,152	6.21%	5.28	7.16	64,157,488	93.60%	131,512	0.19%
Overweight: 25.0 - 29.9	72,695,180	6,470,774	8.90%	7.85	10.00	66,034,529	90.84%	189,876	0.26%
Obese: 30 or higher	75,619,061	8,225,998	10.88%	9.70	12.11	67,225,957	88.90%	167,105	0.22%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-
Yes	98,989,721	9,559,357	9.66%	8.45	10.92	89,132,061	90.04%	298,303	0.30%
No	120,277,342	9,660,759	8.03%	7.36	8.73	110,424,626	91.81%	191,958	0.16%
Refused	*	*	*	*	*	*	*	*	*
<b>Self-reported weight 10 years ago</b>	-	-	-	-	-	-	-	-	-
No clinically significant weight loss	189,289,170	15,620,556	8.25%	7.63	8.90	173,326,791	91.57%	341,824	0.18%
Clinically significant weight loss	30,064,035	3,604,457	11.99%	10.39	13.72	26,298,531	87.48%	161,047	0.54%
<b>Most ever weighed, excluding pregnancy</b>	-	-	-	-	-	-	-	-	-
No clinically significant weight loss	103,240,126	8,493,782	8.23%	7.42	9.06	94,552,652	91.59%	193,691	0.19%
Clinically significant weight loss	116,113,079	10,731,230	9.24%	8.50	10.03	105,072,670	90.49%	309,180	0.27%
<b>Regular milk intake</b>	-	-	-	-	-	-	-	-	-
Regular milk drinker for most or all of life	95,572,997	8,391,745	8.78%	7.93	9.67	86,938,661	90.97%	242,592	0.25%
Have never been a regular milk drinker	48,010,562	4,368,320	9.10%	7.34	10.89	43,561,243	90.73%	81,000	0.17%
Milk drinking has varied over life	75,695,551	6,464,947	8.54%	7.48	9.64	69,063,934	91.24%	166,670	0.22%
Refused/Don't know	*	*	*	*	*	*	*	*	*
<b>Number of meals not home-prepared</b>	-	-	-	-	-	-	-	-	-
0-4 meals	152,720,132	13,923,126	9.12%	8.51	9.76	138,427,777	90.64%	369,227	0.24%
More than 5 meals	66,594,939	5,294,016	7.95%	6.66	9.28	61,167,279	91.85%	133,644	0.20%
Refused/Don't know	*	*	*	*	*	*	*	*	*

SOURCE: National Health and Nutrition Examination Survey, 2007-2012.

NOTE: CI stands for confidence interval.

NOTE: \* represents that figure does not meet the standard for reliability or precision.

NOTE: Data is for all individuals ages 18 and older.

NOTE: The data in this table are based on question KI.Q.026: {Have you/Has SP} ever had kidney stones?

Table 1 - Prevalence of Kidney Stones in Individuals Ages 18 and Older, 2007-2012

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)	Refused to Answer/Don't Know (N)	Refused to Answer/Don't Know (%)
<b>Total</b>	219,353,204	19,225,012	8.76%	8.12	9.44	199,625,322	91.01%	502,871	0.23%
<b>Number of meals from fast food or pizza place</b>	-	-	-	-	-	-	-	-	-
0-4 meals	156,413,270	13,404,940	8.57%	7.81	9.37	142,657,313	91.21%	351,017	0.22%
More than 5 meals	23,779,932	2,153,302	9.06%	7.46	10.68	21,586,246	90.78%	40,385	0.17%
Refused/Don't know	*	*	*	*	*	*	*	*	*
<b>Ever told had gout</b>	-	-	-	-	-	-	-	-	-
Yes	8,210,519	1,611,362	19.63%	16.37	22.93	6,588,518	80.24%	10,639	0.13%
No	211,024,933	17,583,825	8.33%	7.68	9.03	192,963,761	91.44%	477,347	0.23%
Don't know	*	*	*	*	*	*	*	*	*

SOURCE: National Health and Nutrition Examination Survey, 2007-2012.

NOTE: CI stands for confidence interval.

NOTE: \* represents that figure does not meet the standard for reliability or precision.

NOTE: Data is for all individuals ages 18 and older.

NOTE: The data in this table are based on question KIQ.026: {Have you/Has SP} ever had kidney stones?