

Table 2 - Number of Kidney Stones Passed among Those Reporting Having Had Kidney Stone, 2007-2012

| | Total | 1 stone (N) | 1 stone (%) | 2 stones (N) | 2 stones (%) | 3+ stones (N) | 3+ stones (%) |
|--|------------|-------------|-------------|--------------|--------------|---------------|---------------|
| Total | 19,225,012 | 12,416,079 | 64.58% | 2,841,253 | 14.78% | 3,713,587 | 19.32% |
| Gender | - | - | - | - | - | - | - |
| Male | 10,517,819 | 6,558,211 | 62.35% | 1,765,179 | 16.78% | 2,088,584 | 19.86% |
| Female | 8,707,192 | 5,857,866 | 67.28% | 1,076,073 | 12.36% | 1,625,001 | 18.66% |
| Age at screening | - | - | - | - | - | - | - |
| 18-24 | 517,927 | 309,931 | 59.84% | 74,549 | 14.39% | 105,577 | 20.38% |
| 25-34 | 1,867,949 | 1,198,403 | 64.16% | 241,624 | 12.94% | 399,858 | 21.41% |
| 35-44 | 3,206,346 | 2,104,265 | 65.63% | 339,425 | 10.59% | 706,309 | 22.03% |
| 45-54 | 4,549,844 | 2,886,760 | 63.45% | 786,811 | 17.29% | 819,412 | 18.01% |
| 55-59 | 1,887,311 | 1,244,777 | 65.96% | 347,662 | 18.42% | 256,401 | 13.59% |
| 60-64 | 2,146,007 | 1,358,306 | 63.29% | 251,645 | 11.73% | 530,585 | 24.72% |
| 65-69 | 1,486,413 | 939,764 | 63.22% | 191,515 | 12.88% | 347,392 | 23.37% |
| 70-74 | 1,162,716 | 602,405 | 51.81% | 275,136 | 23.66% | 279,852 | 24.07% |
| 75-79 | 928,032 | 662,233 | 71.36% | 127,924 | 13.78% | 134,015 | 14.44% |
| 80+ | 1,472,467 | 1,109,234 | 75.33% | 204,963 | 13.92% | 134,189 | 9.11% |
| Race/ethnicity | - | - | - | - | - | - | - |
| Mexican American | 1,154,898 | 827,252 | 71.63% | 155,275 | 13.44% | 122,001 | 10.56% |
| Other Hispanic | 994,475 | 604,555 | 60.79% | 227,456 | 22.87% | 159,996 | 16.09% |
| Non-Hispanic White | 15,003,061 | 9,478,631 | 63.18% | 2,147,131 | 14.31% | 3,212,906 | 21.42% |
| Non-Hispanic Black | 1,100,233 | 856,838 | 77.88% | 124,894 | 11.35% | 94,048 | 8.55% |
| Other race - including multi-racial | 972,346 | 648,804 | 66.73% | 186,497 | 19.18% | 124,637 | 12.82% |
| Poverty income ratio | - | - | - | - | - | - | - |
| Poverty income ratio less than 1 | 2,386,576 | 1,554,066 | 65.12% | 350,707 | 14.69% | 365,378 | 15.31% |
| Poverty income ratio 1 or greater | 15,352,905 | 9,998,757 | 65.13% | 2,187,856 | 14.25% | 3,083,756 | 20.09% |
| Body mass index | - | - | - | - | - | - | - |
| Normal: less than 25 | 4,255,152 | 2,960,246 | 69.57% | 579,549 | 13.62% | 693,581 | 16.30% |
| Overweight: 25.0 - 29.9 | 6,470,774 | 4,260,334 | 65.84% | 924,409 | 14.29% | 1,161,631 | 17.95% |
| Obese: 30 or higher | 8,225,998 | 4,971,091 | 60.43% | 1,308,252 | 15.90% | 1,838,736 | 22.35% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - |
| Yes | 9,559,357 | 6,113,492 | 63.95% | 1,406,710 | 14.72% | 1,960,824 | 20.51% |
| No | 9,660,759 | 6,297,691 | 65.19% | 1,434,543 | 14.85% | 1,752,762 | 18.14% |
| Don't know | * | * | * | * | * | * | * |
| Self-reported weight 10 years ago | - | - | - | - | - | - | - |
| No clinically significant weight loss | 15,620,556 | 9,883,466 | 63.27% | 2,382,771 | 15.25% | 3,120,382 | 19.98% |
| Clinically significant weight loss | 3,604,457 | 2,532,613 | 70.26% | 458,482 | 12.72% | 593,205 | 16.46% |
| Most ever weighed, excluding pregnancy | - | - | - | - | - | - | - |
| No clinically significant weight loss | 8,493,782 | 5,492,932 | 64.67% | 1,398,476 | 16.46% | 1,455,482 | 17.14% |
| Clinically significant weight loss | 10,731,230 | 6,923,146 | 64.51% | 1,442,777 | 13.44% | 2,258,105 | 21.04% |
| Regular milk intake | - | - | - | - | - | - | - |
| Regular milk drinker for most or all of life | 8,391,745 | 5,373,857 | 64.04% | 1,439,582 | 17.15% | 1,449,194 | 17.27% |
| Have never been a regular milk drinker | 4,368,320 | 2,886,616 | 66.08% | 457,028 | 10.46% | 999,202 | 22.87% |
| Milk drinking has varied over life | 6,464,947 | 4,155,605 | 64.28% | 944,642 | 14.61% | 1,265,191 | 19.57% |
| Number of meals not home-prepared | - | - | - | - | - | - | - |
| 0-4 meals | 13,923,126 | 8,797,542 | 63.19% | 2,116,720 | 15.20% | 2,846,376 | 20.44% |
| More than 5 meals | 5,294,016 | 3,618,536 | 68.35% | 722,277 | 13.64% | 861,602 | 16.28% |
| Refused/Don't know | * | * | * | * | * | * | * |
| Number of meals from fast food or pizza place | - | - | - | - | - | - | - |
| 0-4 meals | 13,404,940 | 8,330,415 | 62.14% | 2,123,816 | 15.84% | 2,817,448 | 21.02% |
| More than 5 meals | 2,153,302 | 1,529,621 | 71.04% | 265,970 | 12.35% | 276,178 | 12.83% |
| Ever told had gout | - | - | - | - | - | - | - |
| Yes | 1,611,362 | 1,090,870 | 67.70% | 198,217 | 12.30% | 308,223 | 19.13% |
| No | 17,583,825 | 11,295,385 | 64.24% | 2,643,036 | 15.03% | 3,405,364 | 19.37% |
| Don't know | * | * | * | * | * | * | * |

SOURCE: National Health and Nutrition Examination Survey, 2007-2012.

NOTE: Data is for all individuals ages 18 and older.

NOTE: The data in this table are based on question KID.028: How many times {have you/has SP} passed a kidney stone?

NOTE: * represents that figure does not meet the standard for reliability or precision.