

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*
20-29	12,825,480	3,572,600	27.86%	19.33%	36.39%	9,252,880	72.14%
30-39	18,943,946	7,678,692	40.53%	31.59%	49.48%	11,265,254	59.47%
40-49	22,500,594	11,180,818	49.69%	42.49%	56.90%	11,319,776	50.31%
50-59	17,533,327	10,205,687	58.21%	53.30%	63.12%	7,327,640	41.79%
60-64	5,474,602	2,894,093	52.86%	42.27%	63.46%	2,580,509	47.14%
65-69	5,240,754	3,113,945	59.42%	48.07%	70.77%	2,126,809	40.58%
70-74	4,310,387	2,496,191	57.91%	50.64%	65.18%	1,814,196	42.09%
75-79	4,281,324	2,948,350	68.87%	58.39%	79.34%	1,332,974	31.13%
80-84	2,888,291	1,888,506	65.38%	51.67%	79.10%	999,785	34.62%
85+	2,160,116	1,436,839	66.52%	46.36%	86.68%	723,277	33.48%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,044,522	2,718,243	44.97%	41.55%	48.39%	3,326,279	55.03%
Other Hispanic	5,721,649	2,222,528	38.84%	25.66%	52.03%	3,499,121	61.16%
Non-Hispanic White	69,130,204	36,773,932	53.20%	50.31%	56.08%	32,356,272	46.80%
Non-Hispanic Black	10,606,305	3,451,308	32.54%	26.17%	38.91%	7,154,998	67.46%
Other Race - Including Multi-Racial	4,925,900	2,298,987	46.67%	28.01%	65.33%	2,626,913	53.33%
Education	-	-	-	-	-	-	-
Missing	2,566,250	1,147,523	44.72%	28.47%	60.96%	1,418,727	55.28%
Less Than 9th Grade	7,001,060	3,648,026	52.11%	45.15%	59.06%	3,353,033	47.89%
9-11th Grade	11,493,555	5,689,145	49.50%	42.70%	56.30%	5,804,410	50.50%
High School Grad/GED or Equivalent	75,160,644	36,955,169	49.17%	45.75%	52.59%	38,205,474	50.83%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	606,658	132,015	21.76%	0.30%	43.22%	474,643	78.24%
Married	56,394,435	28,943,639	51.32%	46.92%	55.73%	27,450,796	48.68%
Widowed	9,963,613	6,231,408	62.54%	54.41%	70.67%	3,732,205	37.46%
Divorced	9,586,940	4,912,696	51.24%	44.79%	57.70%	4,674,244	48.76%
Separated	3,664,817	1,496,357	40.83%	32.12%	49.54%	2,168,460	59.17%
Never Married	10,695,115	4,174,795	39.03%	32.42%	45.65%	6,520,321	60.97%
Living with Partner	5,458,082	1,537,737	28.17%	14.59%	41.75%	3,920,344	71.83%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,237,721	2,840,140	45.53%	34.65%	56.41%	3,397,580	54.47%
Poverty income ratio less than 1	14,133,124	6,126,491	43.35%	36.56%	50.14%	8,006,633	56.65%
Poverty income ratio 1 or greater	76,057,736	38,498,366	50.62%	47.73%	53.50%	37,559,370	49.38%
Body mass index	-	-	-	-	-	-	-
Missing	3,885,282	2,457,815	63.26%	52.48%	74.04%	1,427,467	36.74%
Normal: less than 25	36,617,802	14,775,941	40.35%	34.83%	45.87%	21,841,860	59.65%
Overweight: 25.0 - 29.9	25,513,375	13,204,493	51.76%	46.68%	56.83%	12,308,882	48.24%
Obese: 30 or higher	30,412,122	17,026,748	55.99%	52.10%	59.87%	13,385,375	44.01%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,329,836	1,136,003	34.12%	19.54%	48.69%	2,193,832	65.88%
Glycohemoglobin < 5.7%	75,071,657	35,757,787	47.63%	44.69%	50.58%	39,313,870	52.37%
Glycohemoglobin 5.7% - 6.4%	13,338,158	8,147,720	61.09%	54.89%	67.28%	5,190,438	38.91%
Glycohemoglobin 6.5% or higher	4,688,929	2,423,486	51.69%	40.40%	62.97%	2,265,443	48.31%
Diabetes	-	-	-	-	-	-	-
Yes	6,416,102	3,400,429	53.00%	45.38%	60.62%	3,015,674	47.00%
No	90,012,478	44,064,568	48.95%	45.54%	52.37%	45,947,910	51.05%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,796,498	22,046,460	52.75%	47.52%	57.97%	19,750,038	47.25%
No	54,539,318	25,349,918	46.48%	42.53%	50.43%	29,189,399	53.52%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	27,336,826	16,482,833	60.30%	56.60%	64.00%	10,853,993	39.70%
No	68,701,804	30,754,128	44.76%	40.59%	48.94%	37,947,676	55.24%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,316,505	4,054,628	64.19%	53.98%	74.40%	2,261,877	35.81%
Mobility limitations	13,215,083	8,924,360	67.53%	60.98%	74.08%	4,290,723	32.47%
No mobility limitations	17,642,704	9,994,909	56.65%	50.32%	62.98%	7,647,795	43.35%
N/A - Respondent not asked mobility limitation questions	59,254,288	24,491,100	41.33%	36.56%	46.11%	34,763,189	58.67%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	3,837,297	2,390,395	62.29%	46.86%	77.73%	1,446,903	37.71%
30 to 34	3,723,416	2,413,729	64.83%	54.76%	74.89%	1,309,687	35.17%
35 to 39	4,731,533	3,044,250	64.34%	55.08%	73.60%	1,687,283	35.66%
40 to 44	5,719,731	3,362,062	58.78%	49.28%	68.28%	2,357,669	41.22%
45 to 49	10,167,552	5,551,790	54.60%	48.57%	60.64%	4,615,762	45.40%
50 to 54	9,442,873	6,200,160	65.66%	56.62%	74.70%	3,242,713	34.34%
55 or older	3,441,979	2,036,340	59.16%	48.52%	69.80%	1,405,639	40.84%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	55,153,584	22,330,686	40.49%	36.08%	44.90%	32,822,898	59.51%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,404,672	676,346	48.15%	32.06%	64.24%	728,327	51.85%
None	15,771,213	4,509,228	28.59%	24.45%	32.73%	11,261,985	71.41%
1	10,550,872	5,200,552	49.29%	39.29%	59.29%	5,350,321	50.71%
2	18,612,397	9,422,571	50.63%	43.83%	57.42%	9,189,826	49.37%
More than 2	50,089,425	27,656,300	55.21%	50.93%	59.49%	22,433,126	44.79%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,107,547	938,708	44.54%	28.60%	60.48%	1,168,839	55.46%
29 years old or younger	56,147,820	31,053,427	55.31%	52.09%	58.52%	25,094,393	44.69%
30 - 40 years old	3,449,954	1,745,958	50.61%	38.25%	62.96%	1,703,996	49.39%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	4,509,228	28.59%	24.45%	32.73%	11,261,985	71.41%
Only 1 live birth - see age at last live birth	15,349,454	7,354,891	47.92%	41.35%	54.48%	7,994,563	52.08%
No pregnancies resulting in live births	3,535,523	1,846,854	52.24%	37.64%	66.83%	1,688,669	47.76%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	7,125,329	3,773,094	52.95%	44.14%	61.76%	3,352,235	47.05%
29 years old or younger	41,926,311	20,591,782	49.11%	45.23%	53.00%	21,334,529	50.89%
30 - 40 years old	25,851,442	15,131,027	58.53%	53.86%	63.20%	10,720,415	41.47%
41 years or older	2,115,817	1,570,123	74.21%	55.59%	92.83%	545,694	25.79%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	4,509,228	28.59%	24.45%	32.73%	11,261,985	71.41%
No pregnancies resulting in live births	3,535,523	1,846,854	52.24%	37.64%	66.83%	1,688,669	47.76%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	42,912,018	17,039,045	39.71%	34.73%	44.69%	25,872,974	60.29%
Yes	20,960,048	13,907,890	66.35%	62.31%	70.40%	7,052,157	33.65%
No	32,527,622	16,518,062	50.78%	46.58%	54.99%	16,009,560	49.22%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	42,973,654	17,071,788	39.73%	34.82%	44.63%	25,901,866	60.27%
35 years old or younger	7,522,654	5,168,523	68.71%	58.69%	78.72%	2,354,132	31.29%
36 - 50 years old	10,843,595	6,881,436	63.46%	59.39%	67.53%	3,962,159	36.54%
51 years or older	2,336,250	1,655,099	70.84%	59.75%	81.94%	681,151	29.16%
Refused/Don't know	224,806	170,090	75.66%	46.32%	105.00%	54,716	24.34%
N/A - Did not have procedure	32,527,622	16,518,062	50.78%	46.58%	54.99%	16,009,560	49.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,428,581	47,464,997	49.22%	46.19%	52.25%	48,963,584	50.78%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	8,390,140	4,011,915	47.82%	37.63%	58.00%	4,378,225	52.18%
Yes	20,960,048	13,907,890	66.35%	62.31%	70.40%	7,052,157	33.65%
No	67,049,500	29,545,192	44.06%	40.73%	47.40%	37,504,309	55.94%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Age at screening	-	-	-	-	-	-	-
10-19	696,430	292,416	41.99%	12.35%	71.63%	404,014	58.01%
20-29	12,963,783	3,608,846	27.84%	22.09%	33.58%	9,354,937	72.16%
30-39	17,035,030	7,277,402	42.72%	36.57%	48.87%	9,757,628	57.28%
40-49	20,812,873	10,952,929	52.63%	48.69%	56.57%	9,859,944	47.37%
50-59	18,372,452	10,674,979	58.10%	51.21%	65.00%	7,697,474	41.90%
60-64	5,613,252	2,899,455	51.65%	40.64%	62.67%	2,713,796	48.35%
65-69	6,373,059	4,074,747	63.94%	55.00%	72.88%	2,298,312	36.06%
70-74	5,295,769	3,191,932	60.27%	47.70%	72.84%	2,103,838	39.73%
75-79	5,606,134	3,607,529	64.35%	53.89%	74.81%	1,998,605	35.65%
80-84	3,339,588	1,806,355	54.09%	41.90%	66.28%	1,533,233	45.91%
85+	2,149,983	1,282,311	59.64%	46.16%	73.13%	867,673	40.36%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,695,203	3,320,375	49.59%	43.31%	55.88%	3,374,827	50.41%
Other Hispanic	3,490,447	1,612,394	46.19%	22.54%	69.85%	1,878,053	53.81%
Non-Hispanic White	71,803,772	37,628,646	52.40%	49.21%	55.60%	34,175,126	47.60%
Non-Hispanic Black	11,514,225	4,998,893	43.41%	39.36%	47.47%	6,515,333	56.59%
Other Race - Including Multi-Racial	4,754,706	2,108,592	44.35%	32.92%	55.77%	2,646,114	55.65%
Education	-	-	-	-	-	-	-
Missing	3,272,227	1,565,375	47.84%	24.23%	71.45%	1,706,852	52.16%
Less Than 9th Grade	7,084,184	3,808,148	53.76%	47.13%	60.38%	3,276,036	46.24%
9-11th Grade	10,222,174	5,416,849	52.99%	44.47%	61.51%	4,805,325	47.01%
High School Grad/GED or Equivalent	77,555,073	38,831,527	50.07%	47.12%	53.02%	38,723,546	49.93%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	538,429	125,408	23.29%	(2.05%)	48.63%	413,021	76.71%
Married	55,195,562	28,441,613	51.53%	48.16%	54.89%	26,753,949	48.47%
Widowed	11,477,397	7,023,421	61.19%	52.42%	69.96%	4,453,975	38.81%
Divorced	11,142,010	5,843,281	52.44%	42.83%	62.06%	5,298,729	47.56%
Separated	2,410,096	988,558	41.02%	30.77%	51.26%	1,421,538	58.98%
Never Married	12,255,856	4,823,731	39.36%	32.50%	46.22%	7,432,125	60.64%
Living with Partner	5,186,193	2,422,887	46.72%	37.10%	56.34%	2,763,306	53.28%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	5,148,576	2,464,445	47.87%	40.18%	55.55%	2,684,131	52.13%
Poverty income ratio less than 1	13,079,770	6,757,628	51.66%	44.21%	59.12%	6,322,141	48.34%
Poverty income ratio 1 or greater	80,030,008	40,446,827	50.54%	47.94%	53.14%	39,583,180	49.46%
Body mass index	-	-	-	-	-	-	-
Missing	1,422,592	746,870	52.50%	34.16%	70.84%	675,721	47.50%
Normal: less than 25	36,283,873	14,598,962	40.24%	36.92%	43.55%	21,684,911	59.76%
Overweight: 25.0 - 29.9	27,727,409	14,607,828	52.68%	47.11%	58.26%	13,119,581	47.32%
Obese: 30 or higher	32,824,479	19,715,241	60.06%	55.35%	64.78%	13,109,239	39.94%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,306,134	1,123,877	33.99%	24.42%	43.57%	2,182,258	66.01%
Glycohemoglobin < 5.7%	75,274,178	36,705,283	48.76%	45.94%	51.59%	38,568,895	51.24%
Glycohemoglobin 5.7% - 6.4%	13,399,092	7,967,296	59.46%	53.10%	65.83%	5,431,796	40.54%
Glycohemoglobin 6.5% or higher	6,278,949	3,872,445	61.67%	49.39%	73.96%	2,406,503	38.33%
Diabetes	-	-	-	-	-	-	-
Yes	8,142,154	5,238,254	64.33%	55.02%	73.65%	2,903,901	35.67%
No	90,116,199	44,430,647	49.30%	46.88%	51.73%	45,685,552	50.70%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	43,300,708	23,427,363	54.10%	50.83%	57.37%	19,873,345	45.90%
No	54,921,840	26,236,835	47.77%	44.15%	51.39%	28,685,005	52.23%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,594,115	18,880,048	61.71%	58.13%	65.29%	11,714,067	38.29%
No	67,091,553	30,466,498	45.41%	43.17%	47.65%	36,625,055	54.59%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,807,229	7,973,085	67.53%	60.09%	74.96%	3,834,144	32.47%
Mobility limitations	13,102,646	8,334,329	63.61%	57.53%	69.69%	4,768,317	36.39%
No mobility limitations	20,850,571	11,263,149	54.02%	47.72%	60.31%	9,587,422	45.98%
N/A - Respondent not asked mobility limitation questions	52,497,907	22,098,338	42.09%	39.08%	45.11%	30,399,569	57.91%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	2,539,111	1,398,757	55.09%	37.65%	72.52%	1,140,354	44.91%
Younger than 30	2,891,152	1,813,301	62.72%	44.42%	81.02%	1,077,850	37.28%
30 to 34	4,500,186	3,352,427	74.50%	62.23%	86.76%	1,147,759	25.50%
35 to 39	5,581,951	3,917,954	70.19%	59.54%	80.84%	1,663,996	29.81%
40 to 44	6,448,152	3,852,572	59.75%	52.63%	66.87%	2,595,580	40.25%
45 to 49	10,508,053	6,311,824	60.07%	53.42%	66.72%	4,196,230	39.93%
50 to 54	9,282,520	5,279,756	56.88%	49.19%	64.57%	4,002,764	43.12%
55 or older	4,542,021	2,726,474	60.03%	46.51%	73.54%	1,815,547	39.97%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	51,728,762	20,937,698	40.48%	36.67%	44.28%	30,791,064	59.52%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,870,938	958,973	51.26%	33.08%	69.43%	911,965	48.74%
None	17,857,228	5,982,167	33.50%	25.14%	41.86%	11,875,062	66.50%
1	13,698,789	6,362,208	46.44%	39.05%	53.84%	7,336,582	53.56%
2	22,340,303	10,801,210	48.35%	42.14%	54.55%	11,539,093	51.65%
More than 2	42,491,094	25,564,343	60.16%	56.71%	63.62%	16,926,751	39.84%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Age of patient when she had 9 lb or more baby
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,503,448	1,076,075	42.98%	28.66%	57.31%	1,427,373	57.02%
29 years old or younger	52,997,355	30,909,649	58.32%	55.65%	61.00%	22,087,706	41.68%
30 - 40 years old	4,126,416	2,008,576	48.68%	35.89%	61.46%	2,117,841	51.32%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	5,982,167	33.50%	25.14%	41.86%	11,875,062	66.50%
Only 1 live birth - see age at last live birth	16,014,683	7,964,471	49.73%	43.58%	55.89%	8,050,212	50.27%
No pregnancies resulting in live births	4,581,496	1,665,033	36.34%	21.83%	50.85%	2,916,463	63.66%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,856,024	3,574,328	52.13%	42.86%	61.41%	3,281,695	47.87%
29 years old or younger	42,675,691	23,736,444	55.62%	52.27%	58.97%	18,939,246	44.38%
30 - 40 years old	24,326,556	13,701,788	56.32%	51.96%	60.69%	10,624,768	43.68%
41 years or older	1,773,198	960,358	54.16%	38.39%	69.93%	812,840	45.84%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	5,982,167	33.50%	25.14%	41.86%	11,875,062	66.50%
No pregnancies resulting in live births	4,581,496	1,665,033	36.34%	21.83%	50.85%	2,916,463	63.66%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	41,299,409	17,066,555	41.32%	37.19%	45.46%	24,232,854	58.68%
Yes	21,148,808	14,431,796	68.24%	62.82%	73.66%	6,717,012	31.76%
No	35,742,140	18,102,553	50.65%	46.04%	55.26%	17,639,587	49.35%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	41,367,405	17,134,551	41.42%	37.26%	45.58%	24,232,854	58.58%
35 years old or younger	7,565,019	5,505,373	72.77%	64.64%	80.90%	2,059,646	27.23%
36 - 50 years old	10,954,863	6,930,919	63.27%	55.49%	71.05%	4,023,944	36.73%
51 years or older	2,435,004	1,886,583	77.48%	67.93%	87.02%	548,421	22.52%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	35,742,140	18,102,553	50.65%	46.04%	55.26%	17,639,587	49.35%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,258,353	49,668,900	50.55%	48.21%	52.89%	48,589,453	49.45%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	1,940,125	1,019,234	52.53%	33.69%	71.38%	920,891	47.47%
Yes	21,148,808	14,431,796	68.24%	62.82%	73.66%	6,717,012	31.76%
No	75,101,424	34,149,874	45.47%	42.79%	48.16%	40,951,550	54.53%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Age at screening	-	-	-	-	-	-	-
10-19	398,858	95,437	23.93%	(2.54%)	50.40%	303,420	76.07%
20-29	12,756,001	3,941,809	30.90%	22.89%	38.92%	8,814,191	69.10%
30-39	17,627,319	7,894,514	44.79%	37.95%	51.62%	9,732,805	55.21%
40-49	23,800,262	11,905,894	50.02%	44.94%	55.10%	11,894,369	49.98%
50-59	19,102,084	11,076,998	57.99%	53.98%	62.00%	8,025,086	42.01%
60-64	6,594,843	4,306,761	65.30%	56.11%	74.50%	2,288,082	34.70%
65-69	5,449,338	4,027,952	73.92%	65.12%	82.71%	1,421,386	26.08%
70-74	5,009,301	3,416,773	68.21%	59.13%	77.28%	1,592,528	31.79%
75-79	4,097,250	2,778,416	67.81%	56.09%	79.53%	1,318,834	32.19%
80-84	2,886,255	1,929,806	66.86%	54.62%	79.11%	956,449	33.14%
85+	2,013,172	1,352,603	67.19%	52.49%	81.88%	660,569	32.81%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,995,209	3,728,425	53.30%	45.58%	61.02%	3,266,784	46.70%
Other Hispanic	3,345,447	1,850,588	55.32%	44.72%	65.91%	1,494,859	44.68%
Non-Hispanic White	72,367,356	39,062,749	53.98%	50.19%	57.77%	33,304,607	46.02%
Non-Hispanic Black	11,495,230	5,487,746	47.74%	43.47%	52.01%	6,007,484	52.26%
Other Race - Including Multi-Racial	5,531,440	2,597,454	46.96%	34.95%	58.97%	2,933,986	53.04%
Education	-	-	-	-	-	-	-
Missing	2,515,028	987,475	39.26%	22.42%	56.10%	1,527,554	60.74%
Less Than 9th Grade	5,421,258	3,249,541	59.94%	51.49%	68.39%	2,171,717	40.06%
9-11th Grade	11,067,500	6,419,605	58.00%	52.86%	63.15%	4,647,895	42.00%
High School Grad/GED or Equivalent	80,520,157	42,033,567	52.20%	48.72%	55.68%	38,486,590	47.80%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	745,740	49,746	6.67%	(1.70%)	15.05%	695,994	93.33%
Married	57,232,510	30,891,097	53.97%	49.71%	58.24%	26,341,413	46.03%
Widowed	9,023,315	5,387,335	59.70%	52.23%	67.18%	3,635,980	40.30%
Divorced	12,389,946	6,926,457	55.90%	43.81%	68.00%	5,463,489	44.10%
Separated	3,088,185	1,602,315	51.89%	40.99%	62.78%	1,485,870	48.11%
Never Married	10,375,951	4,816,185	46.42%	35.97%	56.87%	5,559,766	53.58%
Living with Partner	6,702,495	2,963,262	44.21%	37.59%	50.83%	3,739,233	55.79%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	3,202,329	2,060,738	64.35%	58.46%	70.24%	1,141,591	35.65%
Poverty income ratio less than 1	10,921,963	5,439,550	49.80%	40.63%	58.98%	5,482,413	50.20%
Poverty income ratio 1 or greater	85,610,391	45,226,675	52.83%	49.60%	56.05%	40,383,716	47.17%
Body mass index	-	-	-	-	-	-	-
Missing	922,023	564,713	61.25%	38.94%	83.55%	357,310	38.75%
Normal: less than 25	38,372,806	16,462,167	42.90%	37.99%	47.81%	21,910,639	57.10%
Overweight: 25.0 - 29.9	25,231,035	13,604,590	53.92%	47.91%	59.93%	11,626,445	46.08%
Obese: 30 or higher	35,208,819	22,095,493	62.76%	58.86%	66.65%	13,113,326	37.24%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,122,733	1,130,605	36.21%	21.83%	50.58%	1,992,128	63.79%
Glycohemoglobin < 5.7%	75,099,753	37,712,559	50.22%	45.91%	54.52%	37,387,194	49.78%
Glycohemoglobin 5.7% - 6.4%	15,578,112	9,679,205	62.13%	56.43%	67.84%	5,898,907	37.87%
Glycohemoglobin 6.5% or higher	5,934,085	4,204,593	70.85%	60.48%	81.23%	1,729,492	29.15%
Diabetes	-	-	-	-	-	-	-
Yes	8,498,641	5,890,347	69.31%	63.19%	75.43%	2,608,294	30.69%
No	91,236,042	46,836,616	51.34%	48.10%	54.57%	44,399,426	48.66%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	42,045,365	23,259,740	55.32%	50.61%	60.03%	18,785,625	44.68%
No	57,680,831	29,458,736	51.07%	46.45%	55.70%	28,222,095	48.93%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	31,114,886	20,316,113	65.29%	60.17%	70.42%	10,798,773	34.71%
No	68,351,937	32,204,718	47.12%	43.19%	51.04%	36,147,220	52.88%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	10,829,266	7,607,759	70.25%	63.80%	76.70%	3,221,507	29.75%
Mobility limitations	8,792,109	6,347,095	72.19%	62.77%	81.61%	2,445,015	27.81%
No mobility limitations	20,606,777	12,359,990	59.98%	55.36%	64.60%	8,246,786	40.02%
N/A - Respondent not asked mobility limitation questions	59,506,530	26,412,119	44.39%	40.81%	47.96%	33,094,412	55.61%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,664,983	4,500,857	67.53%	52.99%	82.07%	2,164,126	32.47%
No	92,991,453	48,213,491	51.85%	48.76%	54.93%	44,777,962	48.15%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	707,756	509,435	71.98%	44.41%	99.55%	198,321	28.02%
2 or more times a day	11,136,447	6,903,066	61.99%	54.63%	69.34%	4,233,381	38.01%
Once a day	9,094,735	4,756,242	52.30%	43.88%	60.71%	4,338,493	47.70%
2 or more time a week	10,514,091	5,982,021	56.90%	48.53%	65.26%	4,532,070	43.10%
Once a week	6,191,809	3,547,839	57.30%	49.84%	64.76%	2,643,970	42.70%
1-3 times a month	13,475,962	8,401,712	62.35%	53.13%	71.56%	5,074,250	37.65%
Never	48,527,765	22,545,806	46.46%	42.10%	50.82%	25,981,959	53.54%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	707,756	509,435	71.98%	44.41%	99.55%	198,321	28.02%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	10,514,091	5,982,021	56.90%	17.40%	62.80%	4,532,070	43.10%
Once a week	*	*	*	*	*	*	*
1-3 times a month	13,475,962	8,401,712	62.35%	81.83%	101.10%	5,074,250	37.65%
Never	48,527,765	22,545,806	46.46%	49.00%	55.56%	25,981,959	53.54%
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	480,340	235,260	48.98%	13.78%	84.17%	245,080	51.02%
Younger than 30	3,249,527	1,824,552	56.15%	39.12%	73.17%	1,424,975	43.85%
30 to 34	3,113,296	1,997,986	64.18%	44.32%	84.04%	1,115,311	35.82%
35 to 39	5,187,920	3,238,424	62.42%	51.74%	73.11%	1,949,495	37.58%
40 to 44	6,708,828	4,893,379	72.94%	61.60%	84.28%	1,815,449	27.06%
45 to 49	10,986,256	6,724,172	61.21%	51.76%	70.65%	4,262,084	38.79%
50 to 54	11,102,718	7,899,526	71.15%	64.07%	78.22%	3,203,191	28.85%
55 or older	3,656,590	2,465,597	67.43%	56.39%	78.47%	1,190,992	32.57%
Don't know	491,202	152,551	31.06%	(0.60%)	62.71%	338,651	68.94%
N/A - Had period in last 12 months	54,758,007	23,295,515	42.54%	37.38%	47.71%	31,462,491	57.46%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,268,216	4,625,041	28.43%	20.54%	36.32%	11,643,174	71.57%
1	12,829,048	6,051,577	47.17%	40.23%	54.11%	6,777,471	52.83%
2	20,557,226	11,218,022	54.57%	48.72%	60.42%	9,339,204	45.43%
More than 2	50,026,214	30,832,322	61.63%	58.44%	64.83%	19,193,892	38.37%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	993,141	261,752	26.36%	12.48%	40.24%	731,389	73.64%
None	32,053,075	11,854,360	36.98%	30.75%	43.21%	20,198,715	63.02%
1	14,883,888	7,825,805	52.58%	47.64%	57.52%	7,058,083	47.42%
2	21,245,034	12,790,074	60.20%	54.44%	65.97%	8,454,960	39.80%
More than 2	30,536,516	19,994,971	65.48%	61.21%	69.75%	10,541,544	34.52%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	1,016,170	261,752	25.76%	12.26%	39.26%	754,417	74.24%
None	29,973,541	17,652,821	58.89%	56.11%	61.68%	12,320,720	41.11%
1	9,918,573	4,695,588	47.34%	36.87%	57.81%	5,222,985	52.66%
2	5,209,552	2,784,155	53.44%	37.86%	69.02%	2,425,397	46.56%
More than 2	1,827,082	776,865	42.52%	30.03%	55.01%	1,050,216	57.48%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	51,608,282	26,447,433	51.25%	46.55%	55.94%	25,160,849	48.75%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	3,595,632	1,659,608	46.16%	31.35%	60.96%	1,936,024	53.84%
29 years old or younger	55,749,247	34,383,269	61.67%	58.02%	65.33%	21,365,978	38.33%
30 - 40 years old	3,767,991	2,208,454	58.61%	45.75%	71.47%	1,559,537	41.39%
Refused/Don't know	288,861	154,571	53.51%	21.32%	85.70%	134,291	46.49%
N/A - Never pregnant	16,268,216	4,625,041	28.43%	20.54%	36.32%	11,643,174	71.57%
Only 1 live birth - see age at last live birth	16,009,516	8,001,034	49.98%	44.48%	55.47%	8,008,482	50.02%
No pregnancies resulting in live births	4,055,220	1,694,986	41.80%	27.65%	55.95%	2,360,234	58.20%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	3,639,681	1,659,608	45.60%	30.45%	60.74%	1,980,073	54.40%
29 years old or younger	42,817,703	23,327,162	54.48%	49.55%	59.41%	19,490,540	45.52%
30 - 40 years old	31,135,985	20,224,723	64.96%	60.79%	69.12%	10,911,262	35.04%
41 years or older	1,331,628	892,812	67.05%	52.61%	81.49%	438,817	32.95%
Refused/Don't know	486,249	302,631	62.24%	33.17%	91.31%	183,619	37.76%
N/A - Never pregnant	16,268,216	4,625,041	28.43%	20.54%	36.32%	11,643,174	71.57%
No pregnancies resulting in live births	4,055,220	1,694,986	41.80%	27.65%	55.95%	2,360,234	58.20%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	44,487,469	18,596,543	41.80%	37.02%	46.59%	25,890,926	58.20%
Yes	21,019,000	13,722,942	65.29%	58.22%	72.36%	7,296,058	34.71%
No	34,194,845	20,374,110	59.58%	54.26%	64.91%	13,820,736	40.42%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	44,520,837	18,629,911	41.85%	37.12%	46.57%	25,890,926	58.15%
35 years old or younger	6,450,871	3,946,323	61.18%	50.93%	71.42%	2,504,548	38.82%
36 - 50 years old	11,551,731	7,781,942	67.37%	58.36%	76.37%	3,769,789	32.63%
51 years or older	2,278,795	1,747,855	76.70%	67.50%	85.90%	530,940	23.30%
Refused/Don't know	737,604	246,822	33.46%	11.86%	55.07%	490,782	66.54%
N/A - Did not have procedure	34,194,845	20,374,110	59.58%	54.26%	64.91%	13,820,736	40.42%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,734,683	52,726,963	52.87%	49.63%	56.11%	47,007,720	47.13%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,019,000	13,722,942	65.29%	58.22%	72.36%	7,296,058	34.71%
No	78,682,314	38,970,652	49.53%	45.90%	53.16%	39,711,662	50.47%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for “no” or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
Age at screening	-	-	-	-	-	-	-
10-19	723,602	219,056	30.27%	(0.99%)	61.53%	504,547	69.73%
20-29	12,424,789	3,182,912	25.62%	18.28%	32.95%	9,241,877	74.38%
30-39	17,279,284	6,918,439	40.04%	36.45%	43.63%	10,360,844	59.96%
40-49	20,998,655	12,492,517	59.49%	51.29%	67.69%	8,506,137	40.51%
50-59	20,124,734	12,594,999	62.58%	57.02%	68.15%	7,529,735	37.42%
60-64	7,669,158	4,552,494	59.36%	46.08%	72.64%	3,116,664	40.64%
65-69	6,578,135	4,705,139	71.53%	62.03%	81.03%	1,872,996	28.47%
70-74	5,342,587	3,894,677	72.90%	64.05%	81.75%	1,447,910	27.10%
75-79	4,498,310	3,060,873	68.04%	59.82%	76.27%	1,437,437	31.96%
80+	5,603,300	3,633,887	64.85%	55.84%	73.86%	1,969,414	35.15%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,557,995	4,071,673	53.87%	49.72%	58.03%	3,486,323	46.13%
Other Hispanic	4,965,594	2,405,234	48.44%	43.11%	53.76%	2,560,360	51.56%
Non-Hispanic White	71,500,163	40,870,046	57.16%	53.18%	61.14%	30,630,116	42.84%
Non-Hispanic Black	11,993,229	5,876,222	49.00%	44.31%	53.68%	6,117,007	51.00%
Other Race - Including Multi-Racial	5,225,573	2,031,819	38.88%	26.59%	51.18%	3,193,755	61.12%
Education	-	-	-	-	-	-	-
Missing	2,863,940	1,407,816	49.16%	32.70%	65.61%	1,456,124	50.84%
Less Than 9th Grade	6,386,684	4,092,738	64.08%	56.31%	71.85%	2,293,945	35.92%
9-11th Grade	13,278,989	7,310,642	55.05%	46.04%	64.06%	5,968,347	44.95%
High School Grad/GED or Equivalent	78,701,325	42,432,181	53.92%	49.98%	57.85%	36,269,145	46.08%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,342,297	452,036	33.68%	17.23%	50.13%	890,260	66.32%
Married	57,555,671	32,317,208	56.15%	52.75%	59.55%	25,238,463	43.85%
Widowed	9,350,724	6,068,669	64.90%	57.58%	72.22%	3,282,055	35.10%
Divorced	12,590,987	7,535,511	59.85%	52.07%	67.63%	5,055,476	40.15%
Separated	2,378,944	1,434,023	60.28%	47.92%	72.64%	944,921	39.72%
Never Married	11,834,836	4,683,803	39.58%	30.93%	48.22%	7,151,033	60.42%
Living with Partner	6,162,712	2,737,359	44.42%	30.82%	58.01%	3,425,353	55.58%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	7,045,954	3,595,462	51.03%	38.30%	63.76%	3,450,492	48.97%
Poverty income ratio less than 1	13,770,564	7,088,380	51.47%	44.34%	58.61%	6,682,184	48.53%
Poverty income ratio 1 or greater	80,426,036	44,571,151	55.42%	51.20%	59.64%	35,854,885	44.58%
Body mass index	-	-	-	-	-	-	-
Missing	963,961	646,049	67.02%	51.63%	82.41%	317,912	32.98%
Normal: less than 25	35,580,675	16,148,519	45.39%	38.82%	51.95%	19,432,157	54.61%
Overweight: 25.0 - 29.9	29,297,628	15,692,750	53.56%	46.71%	60.42%	13,604,877	46.44%
Obese: 30 or higher	35,400,290	22,767,675	64.31%	59.61%	69.02%	12,632,616	35.69%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,619,352	1,536,929	42.46%	32.40%	52.53%	2,082,423	57.54%
Glycohemoglobin < 5.7%	68,219,894	33,413,569	48.98%	44.32%	53.64%	34,806,325	51.02%
Glycohemoglobin 5.7% - 6.4%	22,416,735	15,105,548	67.39%	60.78%	73.99%	7,311,187	32.61%
Glycohemoglobin 6.5% or higher	6,986,572	5,198,947	74.41%	69.74%	79.09%	1,787,625	25.59%
Diabetes	-	-	-	-	-	-	-
Yes	9,437,447	6,970,907	73.86%	66.60%	81.13%	2,466,539	26.14%
No	91,805,107	48,284,086	52.59%	48.56%	56.63%	43,521,022	47.41%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,244,498	24,409,633	59.18%	55.09%	63.28%	16,834,865	40.82%
No	59,933,874	30,845,360	51.47%	46.44%	56.50%	29,088,514	48.53%
Refused	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	32,857,231	22,774,299	69.31%	64.27%	74.35%	10,082,932	30.69%
No	68,301,740	32,480,694	47.55%	43.87%	51.24%	35,821,045	52.45%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,674,366	7,426,918	63.62%	58.53%	68.71%	4,247,448	36.38%
Mobility limitations	10,730,823	7,358,320	68.57%	61.37%	75.78%	3,372,504	31.43%
No mobility limitations	20,388,040	12,909,086	63.32%	56.94%	69.69%	7,478,954	36.68%
N/A - Respondent not asked mobility limitation questions	58,449,325	27,560,669	47.15%	43.94%	50.37%	30,888,656	52.85%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,361,300	4,870,121	66.16%	52.36%	79.95%	2,491,179	33.84%
No	93,791,876	50,295,493	53.62%	50.32%	56.93%	43,496,382	46.38%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	627,295	382,322	60.95%	39.89%	82.01%	244,972	39.05%
2 or more times a day	9,899,256	6,277,306	63.41%	51.68%	75.15%	3,621,949	36.59%
Once a day	9,682,887	4,903,241	50.64%	44.41%	56.86%	4,779,646	49.36%
2 or more time a week	7,646,398	4,905,793	64.16%	55.98%	72.34%	2,740,605	35.84%
Once a week	7,022,920	4,625,573	65.86%	54.11%	77.61%	2,397,347	34.14%
1-3 times a month	14,356,537	8,831,478	61.52%	55.18%	67.85%	5,525,059	38.48%
Never	51,780,823	25,216,053	48.70%	45.59%	51.81%	26,564,770	51.30%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	627,295	382,322	60.95%	39.89%	82.01%	244,972	39.05%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	7,646,398	4,905,793	64.16%	42.54%	102.40%	2,740,605	35.84%
Once a week	*	*	*	*	*	*	*
1-3 times a month	14,356,537	8,831,478	61.52%	84.58%	98.64%	5,525,059	38.48%
Never	51,780,823	25,216,053	48.70%	50.20%	57.26%	26,564,770	51.30%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	432,110	116,800	27.03%	(9.83%)	63.89%	315,310	72.97%
Younger than 30	3,069,584	1,553,160	50.60%	39.96%	61.23%	1,516,424	49.40%
30 to 34	3,711,338	1,880,705	50.67%	35.42%	65.93%	1,830,632	49.33%
35 to 39	5,097,417	3,535,465	69.36%	61.08%	77.64%	1,561,952	30.64%
40 to 44	7,562,363	5,262,758	69.59%	58.88%	80.31%	2,299,605	30.41%
45 to 49	11,650,306	7,787,025	66.84%	61.60%	72.08%	3,863,282	33.16%
50 to 54	10,341,790	6,764,134	65.41%	55.51%	75.31%	3,577,656	34.59%
55 or older	5,131,554	3,975,633	77.47%	66.55%	88.39%	1,155,922	22.53%
Don't know	224,390	150,201	66.94%	39.98%	93.89%	74,189	33.06%
N/A - Had period in last 12 months	54,021,700	24,229,111	44.85%	40.52%	49.18%	29,792,589	55.15%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,936,586	6,302,817	37.21%	31.03%	43.40%	10,633,770	62.79%
1	13,395,607	6,110,529	45.62%	39.81%	51.43%	7,285,078	54.38%
2	22,728,804	12,563,352	55.28%	48.16%	62.39%	10,165,452	44.72%
More than 2	47,901,009	30,220,672	63.09%	59.51%	66.67%	17,680,337	36.91%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	674,220	327,229	48.53%	37.29%	59.78%	346,991	51.47%
None	32,086,410	13,145,375	40.97%	37.28%	44.66%	18,941,035	59.03%
1	17,758,915	9,225,483	51.95%	46.72%	57.18%	8,533,433	48.05%
2	23,038,524	14,193,456	61.61%	55.10%	68.11%	8,845,068	38.39%
More than 2	27,670,164	18,349,130	66.31%	61.20%	71.42%	9,321,034	33.69%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	964,977	364,303	37.75%	20.05%	55.46%	600,675	62.25%
None	25,250,769	15,099,244	59.80%	53.70%	65.89%	10,151,525	40.20%
1	9,965,596	5,006,103	50.23%	41.45%	59.02%	4,959,493	49.77%
2	5,549,270	2,687,078	48.42%	38.28%	58.56%	2,862,192	51.58%
More than 2	2,981,345	1,662,174	55.75%	45.97%	65.53%	1,319,171	44.25%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	56,516,277	30,421,771	53.83%	49.02%	58.64%	26,094,506	46.17%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	21,511,495	8,490,183	39.47%	32.90%	46.04%	13,021,312	60.53%
Yes	13,130,418	8,532,270	64.98%	54.61%	75.35%	4,598,148	35.02%
No	66,420,922	38,178,210	57.48%	54.50%	60.46%	28,242,712	42.52%
Don't know	179,719	54,330	30.23%	(4.69%)	65.15%	125,388	69.77%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	21,691,214	8,544,513	39.39%	32.90%	45.88%	13,146,701	60.61%
35 years old or younger	12,104,834	7,821,671	64.62%	53.55%	75.68%	4,283,163	35.38%
36 years or older	932,800	702,686	75.33%	57.98%	92.68%	230,114	24.67%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	66,420,922	38,178,210	57.48%	54.50%	60.46%	28,242,712	42.52%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	4,574,909	2,187,367	47.81%	35.60%	60.02%	2,387,543	52.19%
29 years old or younger	55,881,085	34,586,909	61.89%	57.41%	66.38%	21,294,176	38.11%
30 - 40 years old	5,156,343	3,268,792	63.39%	50.16%	76.62%	1,887,551	36.61%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	16,936,586	6,302,817	37.21%	31.03%	43.40%	10,633,770	62.79%
Only 1 live birth - see age at last live birth	18,125,337	8,746,482	48.26%	43.75%	52.76%	9,378,854	51.74%
No pregnancies resulting in live births	491,126	102,973	20.97%	(2.61%)	44.55%	388,153	79.03%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	5,807,618	2,875,976	49.52%	37.61%	61.43%	2,931,642	50.48%
29 years old or younger	46,352,347	26,428,836	57.02%	52.93%	61.11%	19,923,511	42.98%
30 - 40 years old	29,486,973	18,150,685	61.55%	56.95%	66.16%	11,336,289	38.45%
41 years or older	1,806,507	1,135,527	62.86%	40.57%	85.14%	670,980	37.14%
Refused/Don't know	361,396	258,179	71.44%	56.15%	86.73%	103,216	28.56%
N/A - Never pregnant	16,936,586	6,302,817	37.21%	31.03%	43.40%	10,633,770	62.79%
No pregnancies resulting in live births	491,126	102,973	20.97%	(2.61%)	44.55%	388,153	79.03%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,906,168	14,948,548	68.24%	63.08%	73.39%	6,957,620	31.76%
No	79,178,587	40,268,951	50.86%	46.81%	54.90%	38,909,636	49.14%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	6,125,811	3,434,280	56.06%	47.41%	64.71%	2,691,531	43.94%
36 - 50 years old	12,738,019	9,424,613	73.99%	68.77%	79.20%	3,313,406	26.01%
51 years or older	2,470,853	1,775,979	71.88%	61.43%	82.33%	694,874	28.12%
Refused/Don't know	571,486	313,677	54.89%	23.27%	86.51%	257,809	45.11%
N/A - Did not have procedure	79,178,587	40,268,951	50.86%	46.81%	54.90%	38,909,636	49.14%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,803,240	834,063	46.25%	24.45%	68.06%	969,177	53.75%
Yes	11,812,694	8,428,117	71.35%	62.75%	79.94%	3,384,577	28.65%
No	86,965,881	45,666,685	52.51%	48.17%	56.86%	41,299,196	47.49%
Don't know	660,739	326,128	49.36%	19.23%	79.49%	334,611	50.64%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,463,979	1,160,191	47.09%	25.58%	68.60%	1,303,788	52.91%
35 years old or younger	2,315,265	1,147,976	49.58%	34.40%	64.76%	1,167,289	50.42%
36 - 50 years old	6,648,470	5,268,764	79.25%	70.52%	87.98%	1,379,706	20.75%
51 years or older	2,584,749	1,814,685	70.21%	55.79%	84.63%	770,063	29.79%
Refused/Don't know	264,211	196,692	74.45%	44.43%	104.50%	67,519	25.55%
N/A - Did not have procedure	86,965,881	45,666,685	52.51%	48.17%	56.86%	41,299,196	47.49%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,242,554	55,254,993	54.58%	51.07%	58.08%	45,987,561	45.42%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,329,357	14,751,624	69.16%	63.66%	74.66%	6,577,732	30.84%
No	79,755,399	40,465,875	50.74%	46.68%	54.80%	39,289,523	49.26%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for “no” or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,818,595	53,120,267	54.30%	52.05%	56.56%	44,698,328	45.70%
Age at screening	-	-	-	-	-	-	-
10-19	353,109	262,709	74.40%	50.50%	98.29%	90,401	25.60%
20-29	12,072,924	4,065,787	33.68%	28.31%	39.04%	8,007,137	66.32%
30-39	15,950,025	6,839,684	42.88%	37.55%	48.21%	9,110,341	57.12%
40-49	19,772,641	10,250,083	51.84%	46.42%	57.26%	9,522,558	48.16%
50-59	20,729,616	12,355,481	59.60%	51.60%	67.60%	8,374,135	40.40%
60-64	7,767,115	5,026,629	64.72%	58.10%	71.34%	2,740,486	35.28%
65-69	6,636,364	3,883,820	58.52%	48.19%	68.85%	2,752,544	41.48%
70-74	5,133,707	3,780,684	73.64%	66.52%	80.77%	1,353,024	26.36%
75-79	4,119,631	2,933,187	71.20%	62.13%	80.27%	1,186,444	28.80%
80+	5,283,463	3,722,204	70.45%	61.80%	79.10%	1,561,259	29.55%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,365,117	4,069,163	55.25%	51.96%	58.53%	3,295,954	44.75%
Other Hispanic	4,864,518	2,732,072	56.16%	49.18%	63.15%	2,132,447	43.84%
Non-Hispanic White	67,963,077	37,988,402	55.90%	52.99%	58.80%	29,974,675	44.10%
Non-Hispanic Black	11,442,102	5,747,915	50.23%	43.59%	56.88%	5,694,187	49.77%
Other Race - Including Multi-Racial	6,183,781	2,582,715	41.77%	29.09%	54.44%	3,601,065	58.23%
Education	-	-	-	-	-	-	-
Missing	2,266,489	1,313,679	57.96%	37.77%	78.15%	952,810	42.04%
Less Than 9th Grade	5,812,537	3,531,136	60.75%	53.07%	68.43%	2,281,401	39.25%
9-11th Grade	11,569,166	6,443,993	55.70%	48.52%	62.88%	5,125,173	44.30%
High School Grad/GED or Equivalent	77,969,048	41,666,441	53.44%	51.51%	55.37%	36,302,608	46.56%
Refused	*	*	*	*	*	*	*
Don't Know	156,713	120,376	76.81%	50.11%	103.50%	36,337	23.19%
Marital status	-	-	-	-	-	-	-
Missing	353,109	262,709	74.40%	50.50%	98.29%	90,401	25.60%
Married	55,985,946	31,320,097	55.94%	52.92%	58.96%	24,665,848	44.06%
Widowed	8,890,620	5,725,689	64.40%	58.74%	70.07%	3,164,931	35.60%
Divorced	11,116,682	5,876,548	52.86%	46.03%	59.70%	5,240,134	47.14%
Separated	2,211,003	1,156,763	52.32%	33.27%	71.37%	1,054,240	47.68%
Never Married	11,937,752	5,312,504	44.50%	38.00%	51.01%	6,625,248	55.50%
Living with Partner	7,120,836	3,263,310	45.83%	37.31%	54.35%	3,857,526	54.17%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,365,743	3,461,346	54.37%	47.61%	61.14%	2,904,397	45.63%
Poverty income ratio less than 1	13,921,006	7,420,707	53.31%	49.71%	56.90%	6,500,299	46.69%
Poverty income ratio 1 or greater	77,531,847	42,238,215	54.48%	51.97%	56.99%	35,293,632	45.52%
Body mass index	-	-	-	-	-	-	-
Missing	432,161	293,341	67.88%	34.22%	101.50%	138,820	32.12%
Normal: less than 25	34,146,620	14,584,986	42.71%	38.27%	47.16%	19,561,634	57.29%
Overweight: 25.0 - 29.9	27,289,741	14,911,333	54.64%	51.14%	58.14%	12,378,409	45.36%
Obese: 30 or higher	35,950,074	23,330,608	64.90%	61.57%	68.22%	12,619,466	35.10%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,818,595	53,120,267	54.30%	52.05%	56.56%	44,698,328	45.70%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,219,727	1,668,099	51.81%	41.51%	62.11%	1,551,628	48.19%
Glycohemoglobin < 5.7%	62,621,445	30,602,649	48.87%	46.58%	51.16%	32,018,796	51.13%
Glycohemoglobin 5.7% - 6.4%	25,204,944	16,073,315	63.77%	59.27%	68.27%	9,131,629	36.23%
Glycohemoglobin 6.5% or higher	6,772,479	4,776,203	70.52%	64.99%	76.06%	1,996,276	29.48%
Diabetes	-	-	-	-	-	-	-
Yes	8,795,160	6,380,242	72.54%	65.80%	79.29%	2,414,917	27.46%
No	89,023,436	46,740,025	52.50%	50.36%	54.64%	42,283,411	47.50%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	39,121,940	21,778,096	55.67%	51.93%	59.40%	17,343,844	44.33%
No	58,696,655	31,342,171	53.40%	49.87%	56.92%	27,354,484	46.60%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,233,702	20,351,258	67.31%	63.79%	70.83%	9,882,444	32.69%
No	67,523,526	32,736,151	48.48%	46.57%	50.40%	34,787,375	51.52%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,418,591	7,647,500	66.97%	62.11%	71.84%	3,771,091	33.03%
Mobility limitations	9,660,238	6,972,640	72.18%	67.10%	77.26%	2,687,598	27.82%
No mobility limitations	20,668,087	12,437,902	60.18%	55.61%	64.75%	8,230,185	39.82%
N/A - Respondent not asked mobility limitation questions	56,071,679	26,062,226	46.48%	43.97%	48.99%	30,009,454	53.52%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,026,447	5,143,753	73.21%	67.09%	79.32%	1,882,694	26.79%
No	90,716,210	47,900,575	52.80%	50.19%	55.42%	42,815,634	47.20%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	539,983	307,782	57.00%	23.60%	90.40%	232,201	43.00%
2 or more times a day	9,005,246	5,686,523	63.15%	55.22%	71.07%	3,318,723	36.85%
Once a day	8,275,588	5,070,887	61.28%	56.12%	66.43%	3,204,701	38.72%
2 or more time a week	8,108,175	5,295,409	65.31%	55.73%	74.89%	2,812,766	34.69%
Once a week	6,154,232	3,199,487	51.99%	40.89%	63.09%	2,954,745	48.01%
1-3 times a month	13,260,263	7,825,512	59.01%	48.21%	69.82%	5,434,751	40.99%
Never	52,378,637	25,655,477	48.98%	46.11%	51.85%	26,723,160	51.02%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	539,983	307,782	57.00%	29.78%	92.44%	232,201	43.00%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	8,108,175	5,295,409	65.31%	50.71%	100.80%	2,812,766	34.69%
Once a week	6,154,232	3,199,487	51.99%	42.95%	109.10%	2,954,745	48.01%
1-3 times a month	13,260,263	7,825,512	59.01%	56.98%	92.75%	5,434,751	40.99%
Never	52,378,637	25,655,477	48.98%	51.59%	55.85%	26,723,160	51.02%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,818,595	53,120,267	54.30%	52.05%	56.56%	44,698,328	45.70%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	624,425	248,765	39.84%	17.32%	62.36%	375,661	60.16%
Younger than 30	2,882,766	1,713,383	59.44%	47.72%	71.15%	1,169,384	40.56%
30 to 34	3,254,943	1,802,726	55.38%	45.03%	65.74%	1,452,217	44.62%
35 to 39	4,541,323	2,569,168	56.57%	42.55%	70.60%	1,972,155	43.43%
40 to 44	7,479,695	5,071,108	67.80%	59.61%	75.98%	2,408,587	32.20%
45 to 49	9,847,084	6,526,244	66.28%	59.25%	73.31%	3,320,840	33.72%
50 to 54	13,715,632	9,200,275	67.08%	61.28%	72.88%	4,515,357	32.92%
55 or older	5,142,489	3,421,372	66.53%	57.98%	75.08%	1,721,117	33.47%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	50,043,739	22,444,285	44.85%	41.91%	47.79%	27,599,454	55.15%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	17,565,210	5,537,317	31.52%	27.50%	35.54%	12,027,892	68.48%
1	12,038,159	5,977,406	49.65%	42.46%	56.85%	6,060,753	50.35%
2	23,714,828	13,552,903	57.15%	50.58%	63.72%	10,161,926	42.85%
More than 2	44,396,044	27,978,806	63.02%	59.63%	66.41%	16,417,239	36.98%
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	310,780	239,111	76.94%	58.18%	95.70%	71,669	23.06%
None	32,911,743	12,945,501	39.33%	36.78%	41.89%	19,966,242	60.67%
1	14,225,353	8,115,599	57.05%	50.36%	63.74%	6,109,754	42.95%
2	23,367,578	14,716,051	62.98%	57.27%	68.68%	8,651,527	37.02%
More than 2	27,003,142	17,104,005	63.34%	58.86%	67.82%	9,899,137	36.66%
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	779,924	568,872	72.94%	53.63%	92.25%	211,053	27.06%
None	25,135,681	15,677,793	62.37%	58.46%	66.28%	9,457,888	37.63%
1	9,753,274	5,360,090	54.96%	45.94%	63.98%	4,393,185	45.04%
2	5,288,336	2,754,921	52.09%	44.94%	59.25%	2,533,415	47.91%
More than 2	2,314,074	1,252,718	54.13%	42.91%	65.36%	1,061,356	45.87%
N/A - Vaginal delivery/deliveries only	54,547,307	27,505,875	50.43%	47.13%	53.72%	27,041,432	49.57%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	22,992,485	8,351,508	36.32%	33.02%	39.62%	14,640,977	63.68%
Yes	14,458,970	8,918,445	61.68%	54.92%	68.44%	5,540,526	38.32%
No	59,974,609	35,572,114	59.31%	56.91%	61.71%	24,402,494	40.69%
Don't know	392,531	278,200	70.87%	41.88%	99.86%	114,331	29.13%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	23,385,016	8,629,708	36.90%	33.53%	40.28%	14,755,308	63.10%
35 years old or younger	13,118,409	8,040,593	61.29%	53.89%	68.70%	5,077,816	38.71%
36 years or older	1,284,225	859,571	66.93%	49.14%	84.72%	424,654	33.07%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	59,974,609	35,572,114	59.31%	56.91%	61.71%	24,402,494	40.69%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,818,595	53,120,267	54.30%	52.05%	56.56%	44,698,328	45.70%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	5,427,276	2,814,191	51.85%	39.65%	64.06%	2,613,085	48.15%
29 years old or younger	54,043,097	33,379,414	61.76%	58.81%	64.72%	20,663,684	38.24%
30 - 40 years old	5,231,391	3,223,916	61.63%	52.93%	70.33%	2,007,474	38.37%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,565,210	5,537,317	31.52%	27.50%	35.54%	12,027,892	68.48%
Only 1 live birth - see age at last live birth	15,122,928	7,853,343	51.93%	46.62%	57.24%	7,269,584	48.07%
No pregnancies resulting in live births	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,912,927	3,438,662	49.74%	38.30%	61.18%	3,474,265	50.26%
29 years old or younger	41,075,647	23,775,404	57.88%	55.43%	60.34%	17,300,243	42.12%
30 - 40 years old	29,514,813	18,682,784	63.30%	59.61%	66.99%	10,832,029	36.70%
41 years or older	2,166,264	1,231,955	56.87%	41.68%	72.06%	934,309	43.13%
Refused/Don't know	254,181	142,059	55.89%	11.19%	100.60%	112,122	44.11%
N/A - Never pregnant	17,565,210	5,537,317	31.52%	27.50%	35.54%	12,027,892	68.48%
No pregnancies resulting in live births	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,537,624	13,766,152	67.03%	61.86%	72.20%	6,771,472	32.97%
No	77,202,671	39,326,063	50.94%	48.29%	53.58%	37,876,607	49.06%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	5,391,230	3,420,124	63.44%	54.88%	72.00%	1,971,106	36.56%
36 - 50 years old	11,048,011	7,582,176	68.63%	61.32%	75.94%	3,465,835	31.37%
51 years or older	3,681,428	2,488,392	67.59%	58.73%	76.46%	1,193,036	32.41%
Refused/Don't know	416,955	275,460	66.06%	40.89%	91.24%	141,495	33.94%
N/A - Did not have procedure	77,202,671	39,326,063	50.94%	48.29%	53.58%	37,876,607	49.06%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,954,499	1,291,214	66.06%	51.63%	80.49%	663,285	33.94%
Yes	10,117,822	6,929,756	68.49%	61.75%	75.23%	3,188,066	31.51%
No	85,010,794	44,393,514	52.22%	49.65%	54.79%	40,617,280	47.78%
Refused	*	*	*	*	*	*	*
Don't know	723,930	505,783	69.87%	42.34%	97.39%	218,146	30.13%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,689,979	1,796,997	66.80%	53.41%	80.20%	892,982	33.20%
35 years old or younger	2,196,357	1,438,027	65.47%	50.95%	80.00%	758,330	34.53%
36 - 50 years old	5,245,669	3,788,018	72.21%	61.25%	83.18%	1,457,652	27.79%
51 years or older	2,452,490	1,523,012	62.10%	53.34%	70.87%	929,478	37.90%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	85,010,794	44,393,514	52.22%	49.65%	54.79%	40,617,280	47.78%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F01 - Prevalence of Any Urinary Incontinence Among Females, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,818,595	53,120,267	54.30%	52.05%	56.56%	44,698,328	45.70%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,338,065	13,714,084	67.43%	62.34%	72.52%	6,623,981	32.57%
No	77,402,229	39,378,131	50.87%	48.22%	53.53%	38,024,098	49.13%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for “no” or missing hysterectomy data responses.