

Table F02 - Prevalence of stress urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*
20-29	12,825,480	2,482,684	19.36%	12.57%	26.14%	10,342,796	80.64%
30-39	18,943,946	6,288,693	33.20%	24.71%	41.68%	12,655,253	66.80%
40-49	22,500,594	9,216,571	40.96%	34.21%	47.72%	13,284,023	59.04%
50-59	17,533,327	8,577,959	48.92%	43.92%	53.92%	8,955,368	51.08%
60-64	5,474,602	2,586,198	47.24%	37.43%	57.05%	2,888,404	52.76%
65-69	5,240,754	2,596,196	49.54%	37.69%	61.38%	2,644,557	50.46%
70-74	4,310,387	1,994,382	46.27%	39.70%	52.83%	2,316,005	53.73%
75-79	4,298,734	2,333,591	54.29%	42.05%	66.52%	1,965,143	45.71%
80-84	2,880,661	1,564,983	54.33%	41.81%	66.85%	1,315,678	45.67%
85+	2,160,116	1,020,671	47.25%	32.61%	61.89%	1,139,445	52.75%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,036,892	2,179,964	36.11%	32.20%	40.02%	3,856,928	63.89%
Other Hispanic	5,721,649	1,550,701	27.10%	18.81%	35.39%	4,170,948	72.90%
Non-Hispanic White	69,130,204	30,783,407	44.53%	41.17%	47.89%	38,346,797	55.47%
Non-Hispanic Black	10,623,716	2,438,829	22.96%	16.11%	29.80%	8,184,887	77.04%
Other Race - Including Multi-Racial	4,925,900	1,736,672	35.26%	17.52%	52.99%	3,189,228	64.74%
Education	-	-	-	-	-	-	-
Missing	2,566,250	1,025,184	39.95%	23.98%	55.91%	1,541,066	60.05%
Less Than 9th Grade	6,993,429	2,732,725	39.08%	32.47%	45.68%	4,260,704	60.92%
9-11th Grade	11,493,555	4,552,708	39.61%	30.96%	48.27%	6,940,847	60.39%
High School Grad/GED or Equivalent	75,178,055	30,361,756	40.39%	36.58%	44.19%	44,816,298	59.61%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	606,658	74,621	12.30%	(2.43%)	27.03%	532,037	87.70%
Married	56,394,435	24,615,627	43.65%	39.08%	48.21%	31,778,808	56.35%
Widowed	9,955,983	4,910,232	49.32%	40.81%	57.82%	5,045,750	50.68%
Divorced	9,604,351	3,732,044	38.86%	30.71%	47.00%	5,872,306	61.14%
Separated	3,664,817	1,142,774	31.18%	19.51%	42.86%	2,522,042	68.82%
Never Married	10,695,115	2,968,369	27.75%	20.29%	35.22%	7,726,746	72.25%
Living with Partner	5,458,082	1,209,554	22.16%	10.59%	33.73%	4,248,528	77.84%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,237,721	2,009,449	32.21%	20.76%	43.67%	4,228,272	67.79%
Poverty income ratio less than 1	14,125,494	4,714,007	33.37%	26.70%	40.05%	9,411,487	66.63%
Poverty income ratio 1 or greater	76,075,146	31,966,117	42.02%	38.39%	45.65%	44,109,029	57.98%
Body mass index	-	-	-	-	-	-	-
Missing	3,877,651	2,025,765	52.24%	40.88%	63.61%	1,851,887	47.76%
Normal: less than 25	36,617,802	11,696,016	31.94%	26.01%	37.87%	24,921,786	68.06%
Overweight: 25.0 - 29.9	25,530,785	10,505,442	41.15%	35.47%	46.82%	15,025,343	58.85%
Obese: 30 or higher	30,412,122	14,462,350	47.55%	42.45%	52.66%	15,949,772	52.45%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,329,836	891,814	26.78%	12.49%	41.08%	2,438,022	73.22%
Glycohemoglobin < 5.7%	75,089,068	29,047,153	38.68%	35.21%	42.16%	46,041,915	61.32%
Glycohemoglobin 5.7% - 6.4%	13,338,158	6,822,228	51.15%	44.64%	57.65%	6,515,930	48.85%
Glycohemoglobin 6.5% or higher	4,681,299	1,928,378	41.19%	31.19%	51.20%	2,752,921	58.81%
Diabetes	-	-	-	-	-	-	-
Yes	6,408,472	2,675,310	41.75%	36.52%	46.97%	3,733,162	58.25%
No	90,029,889	36,014,263	40.00%	36.21%	43.80%	54,015,626	60.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,796,498	18,772,437	44.91%	39.47%	50.35%	23,024,062	55.09%
No	54,549,098	19,848,518	36.39%	31.61%	41.16%	34,700,580	63.61%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	27,329,196	14,053,704	51.42%	47.51%	55.34%	13,275,492	48.58%
No	68,719,214	24,498,583	35.65%	31.30%	40.00%	44,220,631	64.35%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,308,875	3,379,061	53.56%	43.42%	63.70%	2,929,814	46.44%
Mobility limitations	13,232,494	7,232,538	54.66%	47.64%	61.68%	5,999,955	45.34%
No mobility limitations	17,642,704	7,921,631	44.90%	38.71%	51.09%	9,721,073	55.10%
N/A - Respondent not asked mobility limitation questions	59,254,288	20,156,342	34.02%	29.55%	38.49%	39,097,946	65.98%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	3,837,297	2,120,605	55.26%	39.81%	70.72%	1,716,692	44.74%
30 to 34	3,723,416	1,957,558	52.57%	41.31%	63.84%	1,765,857	47.43%
35 to 39	4,731,533	2,663,202	56.29%	44.01%	68.57%	2,068,330	43.71%
40 to 44	5,719,731	2,951,461	51.60%	42.63%	60.57%	2,768,270	48.40%
45 to 49	10,184,963	4,380,859	43.01%	36.94%	49.09%	5,804,104	56.99%
50 to 54	9,435,243	5,225,821	55.39%	48.67%	62.10%	4,209,421	44.61%
55 or older	3,441,979	1,555,049	45.18%	36.55%	53.81%	1,886,930	54.82%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	55,153,584	17,756,974	32.20%	27.40%	36.99%	37,396,610	67.80%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,404,672	524,616	37.35%	18.94%	55.76%	880,056	62.65%
None	15,771,213	2,989,695	18.96%	14.40%	23.51%	12,781,517	81.04%
1	10,550,872	4,079,139	38.66%	29.20%	48.12%	6,471,734	61.34%
2	18,612,397	7,965,461	42.80%	36.76%	48.84%	10,646,936	57.20%
More than 2	50,099,206	23,130,661	46.17%	41.69%	50.65%	26,968,544	53.83%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,107,547	674,807	32.02%	18.38%	45.66%	1,432,740	67.98%
29 years old or younger	56,157,600	26,583,303	47.34%	43.95%	50.72%	29,574,297	52.66%
30 - 40 years old	3,449,954	1,448,759	41.99%	29.08%	54.91%	2,001,195	58.01%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	2,989,695	18.96%	14.40%	23.51%	12,781,517	81.04%
Only 1 live birth - see age at last live birth	15,349,454	5,609,434	36.54%	28.94%	44.15%	9,740,020	63.46%
No pregnancies resulting in live births	3,535,523	1,383,575	39.13%	23.66%	54.61%	2,151,948	60.87%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	7,125,329	2,991,677	41.99%	34.16%	49.82%	4,133,652	58.01%
29 years old or younger	41,943,721	17,519,528	41.77%	36.99%	46.55%	24,424,193	58.23%
30 - 40 years old	25,851,442	12,818,285	49.58%	44.88%	54.29%	13,033,157	50.42%
41 years or older	2,108,186	943,923	44.77%	32.45%	57.10%	1,164,263	55.23%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	2,989,695	18.96%	14.40%	23.51%	12,781,517	81.04%
No pregnancies resulting in live births	3,535,523	1,383,575	39.13%	23.66%	54.61%	2,151,948	60.87%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	42,912,018	13,253,555	30.89%	26.01%	35.76%	29,658,463	69.11%
Yes	20,977,458	12,073,527	57.55%	52.19%	62.92%	8,903,931	42.45%
No	32,519,991	13,362,490	41.09%	36.72%	45.46%	19,157,501	58.91%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	42,973,654	13,286,298	30.92%	26.09%	35.75%	29,687,356	69.08%
35 years old or younger	7,522,654	4,460,670	59.30%	48.38%	70.21%	3,061,984	40.70%
36 - 50 years old	10,861,005	6,031,417	55.53%	50.68%	60.38%	4,829,588	44.47%
51 years or older	2,336,250	1,404,236	60.11%	44.33%	75.88%	932,014	39.89%
Refused/Don't know	224,806	144,461	64.26%	40.57%	87.95%	80,345	35.74%
N/A - Did not have procedure	32,519,991	13,362,490	41.09%	36.72%	45.46%	19,157,501	58.91%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,438,361	38,689,573	40.12%	36.43%	43.80%	57,748,788	59.88%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	8,390,140	3,428,486	40.86%	30.42%	51.31%	4,961,654	59.14%
Yes	20,977,458	12,073,527	57.55%	52.19%	62.92%	8,903,931	42.45%
No	67,041,870	23,187,559	34.59%	30.59%	38.59%	43,854,310	65.41%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Age at screening	-	-	-	-	-	-	-
10-19	696,430	165,372	23.75%	(0.75%)	48.24%	531,058	76.25%
20-29	12,963,783	3,210,843	24.77%	18.97%	30.56%	9,752,940	75.23%
30-39	17,120,651	6,117,018	35.73%	28.53%	42.93%	11,003,633	64.27%
40-49	20,812,873	9,374,366	45.04%	40.85%	49.23%	11,438,507	54.96%
50-59	18,372,452	8,687,563	47.29%	40.34%	54.23%	9,684,889	52.71%
60-64	5,613,252	2,250,785	40.10%	32.01%	48.18%	3,362,467	59.90%
65-69	6,373,059	3,369,560	52.87%	45.47%	60.28%	3,003,499	47.13%
70-74	5,295,769	2,454,011	46.34%	35.77%	56.91%	2,841,759	53.66%
75-79	5,606,134	3,007,491	53.65%	41.82%	65.47%	2,598,643	46.35%
80-84	3,339,588	1,373,047	41.11%	29.52%	52.70%	1,966,541	58.89%
85+	2,149,983	927,747	43.15%	29.57%	56.73%	1,222,236	56.85%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,695,203	2,779,841	41.52%	36.36%	46.68%	3,915,362	58.48%
Other Hispanic	3,490,447	1,077,720	30.88%	6.96%	54.80%	2,412,727	69.12%
Non-Hispanic White	71,889,393	31,886,450	44.35%	41.74%	46.97%	40,002,943	55.65%
Non-Hispanic Black	11,514,225	3,421,700	29.72%	24.80%	34.63%	8,092,525	70.28%
Other Race - Including Multi-Racial	4,754,706	1,772,091	37.27%	26.03%	48.51%	2,982,615	62.73%
Education	-	-	-	-	-	-	-
Missing	3,272,227	1,258,930	38.47%	18.56%	58.38%	2,013,297	61.53%
Less Than 9th Grade	7,084,184	3,139,070	44.31%	34.94%	53.68%	3,945,114	55.69%
9-11th Grade	10,222,174	4,198,384	41.07%	33.38%	48.76%	6,023,790	58.93%
High School Grad/GED or Equivalent	77,640,695	32,294,417	41.59%	39.00%	44.19%	45,346,278	58.41%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	538,429	110,673	20.55%	(2.79%)	43.90%	427,756	79.45%
Married	55,281,184	24,069,512	43.54%	40.70%	46.39%	31,211,672	56.46%
Widowed	11,477,397	5,501,185	47.93%	39.11%	56.75%	5,976,212	52.07%
Divorced	11,142,010	4,333,056	38.89%	30.80%	46.97%	6,808,953	61.11%
Separated	2,410,096	898,880	37.30%	27.39%	47.20%	1,511,216	62.70%
Never Married	12,255,856	3,964,174	32.35%	25.17%	39.52%	8,291,682	67.65%
Living with Partner	5,186,193	2,060,322	39.73%	29.10%	50.35%	3,125,871	60.27%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	5,148,576	2,119,563	41.17%	33.19%	49.14%	3,029,013	58.83%
Poverty income ratio less than 1	13,079,770	5,171,207	39.54%	31.74%	47.33%	7,908,563	60.46%
Poverty income ratio 1 or greater	80,115,629	33,647,032	42.00%	39.52%	44.48%	46,468,597	58.00%
Body mass index	-	-	-	-	-	-	-
Missing	1,422,592	611,364	42.98%	25.80%	60.15%	811,228	57.02%
Normal: less than 25	36,369,495	11,818,706	32.50%	29.13%	35.87%	24,550,789	67.50%
Overweight: 25.0 - 29.9	27,727,409	12,104,924	43.66%	39.08%	48.23%	15,622,484	56.34%
Obese: 30 or higher	32,824,479	16,402,809	49.97%	45.34%	54.60%	16,421,671	50.03%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,306,134	1,020,006	30.85%	20.96%	40.74%	2,286,128	69.15%
Glycohemoglobin < 5.7%	75,359,799	30,297,213	40.20%	37.35%	43.06%	45,062,587	59.80%
Glycohemoglobin 5.7% - 6.4%	13,399,092	6,341,434	47.33%	40.35%	54.30%	7,057,658	52.67%
Glycohemoglobin 6.5% or higher	6,278,949	3,279,150	52.22%	41.43%	63.02%	2,999,799	47.78%
Diabetes	-	-	-	-	-	-	-
Yes	8,142,154	4,332,499	53.21%	44.95%	61.47%	3,809,655	46.79%
No	90,201,820	36,605,303	40.58%	38.19%	42.97%	53,596,517	59.42%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	43,300,708	19,253,688	44.47%	41.52%	47.41%	24,047,019	55.53%
No	55,007,462	21,679,412	39.41%	36.67%	42.15%	33,328,050	60.59%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,594,115	15,763,719	51.53%	47.23%	55.83%	14,830,396	48.47%
No	67,177,174	25,009,797	37.23%	35.18%	39.28%	42,167,377	62.77%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,807,229	6,065,226	51.37%	42.28%	60.46%	5,742,003	48.63%
Mobility limitations	13,102,646	6,789,352	51.82%	45.28%	58.35%	6,313,294	48.18%
No mobility limitations	20,850,571	9,254,130	44.38%	38.60%	50.17%	11,596,441	55.62%
N/A - Respondent not asked mobility limitation questions	52,583,529	18,829,095	35.81%	32.93%	38.69%	33,754,434	64.19%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	2,624,733	1,212,097	46.18%	30.10%	62.26%	1,412,636	53.82%
Younger than 30	2,891,152	1,504,079	52.02%	36.15%	67.90%	1,387,073	47.98%
30 to 34	4,500,186	2,658,334	59.07%	48.96%	69.18%	1,841,852	40.93%
35 to 39	5,581,951	2,986,470	53.50%	41.79%	65.22%	2,595,481	46.50%
40 to 44	6,448,152	2,804,191	43.49%	36.53%	50.45%	3,643,961	56.51%
45 to 49	10,508,053	4,913,679	46.76%	38.04%	55.49%	5,594,374	53.24%
50 to 54	9,282,520	4,409,693	47.51%	41.17%	53.84%	4,872,827	52.49%
55 or older	4,542,021	2,240,211	49.32%	38.00%	60.65%	2,301,811	50.68%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	51,728,762	18,136,080	35.06%	31.76%	38.36%	33,592,682	64.94%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,956,559	772,313	39.47%	25.41%	53.53%	1,184,247	60.53%
None	17,857,228	4,779,210	26.76%	20.68%	32.84%	13,078,018	73.24%
1	13,698,789	5,270,778	38.48%	31.64%	45.31%	8,428,011	61.52%
2	22,340,303	9,127,200	40.86%	34.38%	47.33%	13,213,103	59.14%
More than 2	42,491,094	20,988,301	49.39%	45.28%	53.51%	21,502,793	50.61%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,589,069	875,220	33.80%	21.56%	46.05%	1,713,849	66.20%
29 years old or younger	52,997,355	25,690,010	48.47%	46.15%	50.80%	27,307,345	51.53%
30 - 40 years old	4,126,416	1,780,259	43.14%	30.50%	55.79%	2,346,157	56.86%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	4,779,210	26.76%	20.68%	32.84%	13,078,018	73.24%
Only 1 live birth - see age at last live birth	16,014,683	6,503,725	40.61%	34.81%	46.41%	9,510,958	59.39%
No pregnancies resulting in live births	4,581,496	1,246,447	27.21%	17.20%	37.21%	3,335,049	72.79%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,941,645	2,708,022	39.01%	29.89%	48.13%	4,233,623	60.99%
29 years old or younger	42,675,691	20,155,726	47.23%	44.67%	49.79%	22,519,965	52.77%
30 - 40 years old	24,326,556	11,359,884	46.70%	41.83%	51.57%	12,966,672	53.30%
41 years or older	1,773,198	639,731	36.08%	16.98%	55.18%	1,133,467	63.92%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	4,779,210	26.76%	20.68%	32.84%	13,078,018	73.24%
No pregnancies resulting in live births	4,581,496	1,246,447	27.21%	17.20%	37.21%	3,335,049	72.79%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	41,385,030	14,683,175	35.48%	32.21%	38.75%	26,701,855	64.52%
Yes	21,148,808	11,601,040	54.85%	51.03%	58.68%	9,547,768	45.15%
No	35,742,140	14,585,591	40.81%	37.18%	44.44%	21,156,549	59.19%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	41,453,026	14,751,171	35.59%	32.26%	38.91%	26,701,855	64.41%
35 years old or younger	7,565,019	4,421,643	58.45%	51.16%	65.73%	3,143,376	41.55%
36 - 50 years old	10,954,863	5,592,360	51.05%	43.27%	58.83%	5,362,504	48.95%
51 years or older	2,435,004	1,550,288	63.67%	52.81%	74.52%	884,715	36.33%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	35,742,140	14,585,591	40.81%	37.18%	44.44%	21,156,549	59.19%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,343,974	40,937,802	41.63%	39.47%	43.79%	57,406,172	58.37%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	2,025,746	832,574	41.10%	25.14%	57.06%	1,193,172	58.90%
Yes	21,148,808	11,601,040	54.85%	51.03%	58.68%	9,547,768	45.15%
No	75,101,424	28,436,191	37.86%	35.39%	40.34%	46,665,233	62.14%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Age at screening	-	-	-	-	-	-	-
10-19	398,858	77,918	19.54%	(5.98%)	45.05%	320,939	80.46%
20-29	12,756,001	3,239,453	25.40%	18.68%	32.11%	9,516,547	74.60%
30-39	17,627,319	6,960,817	39.49%	32.48%	46.50%	10,666,502	60.51%
40-49	23,800,262	9,920,393	41.68%	38.17%	45.19%	13,879,870	58.32%
50-59	19,102,084	9,123,564	47.76%	42.20%	53.33%	9,978,520	52.24%
60-64	6,594,843	3,555,336	53.91%	44.45%	63.37%	3,039,507	46.09%
65-69	5,433,285	3,222,284	59.31%	48.67%	69.94%	2,211,002	40.69%
70-74	5,062,672	2,368,953	46.79%	35.27%	58.32%	2,693,719	53.21%
75-79	4,097,250	1,890,669	46.14%	33.21%	59.08%	2,206,581	53.86%
80-84	2,919,334	1,215,250	41.63%	28.92%	54.33%	1,704,083	58.37%
85+	2,013,172	989,972	49.17%	34.37%	63.98%	1,023,200	50.83%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,995,209	3,157,732	45.14%	38.13%	52.15%	3,837,477	54.86%
Other Hispanic	3,345,447	1,661,885	49.68%	41.30%	58.05%	1,683,562	50.32%
Non-Hispanic White	72,453,806	31,928,636	44.07%	41.14%	47.00%	40,525,170	55.93%
Non-Hispanic Black	11,479,177	3,517,078	30.64%	26.75%	34.52%	7,962,099	69.36%
Other Race - Including Multi-Racial	5,531,440	2,299,278	41.57%	29.31%	53.83%	3,232,162	58.43%
Education	-	-	-	-	-	-	-
Missing	2,515,028	831,195	33.05%	15.59%	50.51%	1,683,834	66.95%
Less Than 9th Grade	5,421,258	2,556,698	47.16%	35.94%	58.38%	2,864,560	52.84%
9-11th Grade	11,100,579	5,319,906	47.92%	40.74%	55.11%	5,780,673	52.08%
High School Grad/GED or Equivalent	80,557,475	33,852,812	42.02%	39.35%	44.69%	46,704,664	57.98%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	745,740	20,608	2.76%	0.37%	5.15%	725,133	97.24%
Married	57,265,589	25,849,680	45.14%	41.60%	48.68%	31,415,909	54.86%
Widowed	9,060,633	3,465,651	38.25%	28.69%	47.81%	5,594,982	61.75%
Divorced	12,389,946	5,705,638	46.05%	34.89%	57.21%	6,684,308	53.95%
Separated	3,088,185	1,366,068	44.24%	34.40%	54.07%	1,722,117	55.76%
Never Married	10,375,951	3,591,518	34.61%	26.01%	43.22%	6,784,433	65.39%
Living with Partner	6,702,495	2,565,446	38.28%	30.89%	45.66%	4,137,049	61.72%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	3,235,408	1,425,556	44.06%	31.58%	56.54%	1,809,852	55.94%
Poverty income ratio less than 1	10,905,910	4,166,444	38.20%	31.96%	44.44%	6,739,466	61.80%
Poverty income ratio 1 or greater	85,663,762	36,972,609	43.16%	40.78%	45.54%	48,691,153	56.84%
Body mass index	-	-	-	-	-	-	-
Missing	922,023	357,091	38.73%	16.58%	60.88%	564,932	61.27%
Normal: less than 25	38,459,256	13,169,925	34.24%	29.68%	38.81%	25,289,331	65.76%
Overweight: 25.0 - 29.9	25,231,035	10,934,070	43.34%	37.70%	48.98%	14,296,966	56.66%
Obese: 30 or higher	35,192,766	18,103,524	51.44%	48.73%	54.16%	17,089,242	48.56%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,122,733	731,777	23.43%	9.41%	37.46%	2,390,956	76.57%
Glycohemoglobin < 5.7%	75,153,124	30,755,342	40.92%	37.63%	44.22%	44,397,782	59.08%
Glycohemoglobin 5.7% - 6.4%	15,611,191	7,742,768	49.60%	43.22%	55.98%	7,868,423	50.40%
Glycohemoglobin 6.5% or higher	5,918,032	3,334,722	56.35%	44.59%	68.10%	2,583,310	43.65%
Diabetes	-	-	-	-	-	-	-
Yes	8,498,641	4,725,468	55.60%	48.41%	62.79%	3,773,173	44.40%
No	91,306,439	37,839,141	41.44%	39.33%	43.56%	53,467,298	58.56%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	42,029,312	18,903,740	44.98%	41.59%	48.36%	23,125,572	55.02%
No	57,767,281	23,652,382	40.94%	37.42%	44.47%	34,114,899	59.06%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	31,131,911	16,330,392	52.46%	47.77%	57.14%	14,801,519	47.54%
No	68,405,309	26,047,905	38.08%	35.26%	40.90%	42,357,404	61.92%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	10,862,345	6,163,308	56.74%	50.15%	63.33%	4,699,037	43.26%
Mobility limitations	8,776,056	5,172,690	58.94%	51.02%	66.86%	3,603,366	41.06%
No mobility limitations	20,660,148	9,430,103	45.64%	40.98%	50.31%	11,230,045	54.36%
N/A - Respondent not asked mobility limitation questions	59,506,530	21,798,508	36.63%	34.02%	39.24%	37,708,022	63.37%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,664,983	4,060,439	60.92%	46.12%	75.72%	2,604,544	39.08%
No	93,061,850	38,491,555	41.36%	39.47%	43.25%	54,570,295	58.64%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	761,127	481,077	63.21%	33.89%	92.53%	280,050	36.79%
2 or more times a day	11,136,447	5,468,718	49.11%	41.66%	56.56%	5,667,729	50.89%
Once a day	9,078,682	3,871,921	42.65%	35.95%	49.35%	5,206,760	57.35%
2 or more time a week	10,514,091	4,939,864	46.98%	37.85%	56.12%	5,574,227	53.02%
Once a week	6,191,809	3,002,113	48.49%	38.12%	58.85%	3,189,696	51.51%
1-3 times a month	13,509,041	6,764,106	50.07%	42.21%	57.93%	6,744,935	49.93%
Never	48,527,765	18,036,809	37.17%	33.95%	40.39%	30,490,955	62.83%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	761,127	481,077	63.21%	33.89%	92.53%	280,050	36.79%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	10,514,091	4,939,864	46.98%	(7.29%)	56.17%	5,574,227	53.02%
Once a week	*	*	*	*	*	*	*
1-3 times a month	13,509,041	6,764,106	50.07%	51.10%	85.41%	6,744,935	49.93%
Never	48,527,765	18,036,809	37.17%	39.91%	44.53%	30,490,955	62.83%
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	480,340	194,118	40.41%	4.34%	76.48%	286,223	59.59%
Younger than 30	3,249,527	1,205,729	37.10%	23.87%	50.34%	2,043,798	62.90%
30 to 34	3,113,296	1,734,615	55.72%	35.91%	75.52%	1,378,681	44.28%
35 to 39	5,187,920	2,404,562	46.35%	36.22%	56.47%	2,783,358	53.65%
40 to 44	6,708,828	3,761,116	56.06%	45.44%	66.68%	2,947,712	43.94%
45 to 49	11,019,335	4,930,199	44.74%	36.07%	53.41%	6,089,135	55.26%
50 to 54	11,086,665	6,302,197	56.84%	48.25%	65.44%	4,784,467	43.16%
55 or older	3,656,590	2,066,011	56.50%	42.94%	70.06%	1,590,579	43.50%
Don't know	544,573	152,551	28.01%	(0.64%)	56.67%	392,022	71.99%
N/A - Had period in last 12 months	54,758,007	19,813,511	36.18%	31.85%	40.52%	34,944,496	63.82%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,268,216	3,530,858	21.70%	13.43%	29.98%	12,737,358	78.30%
1	12,829,048	4,765,152	37.14%	29.95%	44.33%	8,063,896	62.86%
2	20,643,676	9,163,179	44.39%	39.25%	49.52%	11,480,498	55.61%
More than 2	50,010,161	25,105,420	50.20%	47.27%	53.14%	24,904,741	49.80%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	993,141	217,115	21.86%	7.89%	35.84%	776,026	78.14%
None	32,053,075	9,387,622	29.29%	23.23%	35.35%	22,665,453	70.71%
1	14,883,888	5,851,319	39.31%	34.19%	44.44%	9,032,569	60.69%
2	21,331,484	11,250,354	52.74%	47.20%	58.28%	10,081,131	47.26%
More than 2	30,520,463	15,858,199	51.96%	46.76%	57.16%	14,662,263	48.04%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	1,016,170	217,115	21.37%	7.58%	35.15%	799,054	78.63%
None	29,957,488	14,058,919	46.93%	42.95%	50.91%	15,898,570	53.07%
1	9,918,573	3,944,506	39.77%	31.45%	48.08%	5,974,068	60.23%
2	5,209,552	2,355,885	45.22%	29.52%	60.92%	2,853,667	54.78%
More than 2	1,827,082	708,758	38.79%	26.65%	50.93%	1,118,324	61.21%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	51,694,732	21,171,079	40.95%	36.69%	45.22%	30,523,653	59.05%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	3,595,632	1,213,152	33.74%	16.33%	51.15%	2,382,479	66.26%
29 years old or younger	55,786,566	28,478,414	51.05%	47.59%	54.50%	27,308,151	48.95%
30 - 40 years old	3,767,991	1,797,224	47.70%	32.14%	63.26%	1,970,767	52.30%
Refused/Don't know	288,861	109,890	38.04%	(4.99%)	81.08%	178,971	61.96%
N/A - Never pregnant	16,268,216	3,530,858	21.70%	13.43%	29.98%	12,737,358	78.30%
Only 1 live birth - see age at last live birth	16,042,594	6,285,223	39.18%	33.96%	44.40%	9,757,371	60.82%
No pregnancies resulting in live births	4,055,220	1,149,846	28.35%	14.29%	42.42%	2,905,374	71.65%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	3,639,681	1,213,152	33.33%	15.88%	50.78%	2,426,529	66.67%
29 years old or younger	42,871,074	19,293,999	45.00%	41.02%	48.99%	23,577,075	55.00%
30 - 40 years old	31,169,064	16,382,986	52.56%	48.10%	57.03%	14,786,078	47.44%
41 years or older	1,315,575	775,489	58.95%	44.83%	73.07%	540,086	41.05%
Refused/Don't know	486,249	218,278	44.89%	20.90%	68.89%	267,971	55.11%
N/A - Never pregnant	16,268,216	3,530,858	21.70%	13.43%	29.98%	12,737,358	78.30%
No pregnancies resulting in live births	4,055,220	1,149,846	28.35%	14.29%	42.42%	2,905,374	71.65%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	44,487,469	15,720,625	35.34%	31.45%	39.22%	28,766,843	64.66%
Yes	21,002,947	10,838,859	51.61%	45.09%	58.12%	10,164,088	48.39%
No	34,281,296	15,971,756	46.59%	41.34%	51.84%	18,309,540	53.41%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	44,520,837	15,753,994	35.39%	31.57%	39.20%	28,766,843	64.61%
35 years old or younger	6,450,871	3,055,791	47.37%	39.01%	55.73%	3,395,080	52.63%
36 - 50 years old	11,535,678	6,133,724	53.17%	43.92%	62.42%	5,401,954	46.83%
51 years or older	2,278,795	1,453,457	63.78%	49.10%	78.46%	825,338	36.22%
Refused/Don't know	737,604	195,887	26.56%	6.62%	46.50%	541,716	73.44%
N/A - Did not have procedure	34,281,296	15,971,756	46.59%	41.34%	51.84%	18,309,540	53.41%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,805,080	42,564,609	42.65%	40.49%	44.81%	57,240,471	57.35%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,002,947	10,838,859	51.61%	45.09%	58.12%	10,164,088	48.39%
No	78,768,764	31,692,381	40.23%	37.42%	43.04%	47,076,383	59.77%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,304,896	43,574,138	43.01%	39.89%	46.14%	57,730,757	56.99%
Age at screening	-	-	-	-	-	-	-
10-19	723,602	123,866	17.12%	(0.59%)	34.82%	599,736	82.88%
20-29	12,424,789	2,389,987	19.24%	13.80%	24.67%	10,034,802	80.76%
30-39	17,279,284	5,874,771	34.00%	30.77%	37.23%	11,404,513	66.00%
40-49	20,998,655	10,542,388	50.21%	44.43%	55.98%	10,456,266	49.79%
50-59	20,151,118	10,190,538	50.57%	44.89%	56.26%	9,960,579	49.43%
60-64	7,675,019	3,682,691	47.98%	34.97%	61.00%	3,992,327	52.02%
65-69	6,578,135	3,515,083	53.44%	45.09%	61.78%	3,063,052	46.56%
70-74	5,342,587	2,808,468	52.57%	43.79%	61.34%	2,534,119	47.43%
75-79	4,498,310	1,945,073	43.24%	32.06%	54.42%	2,553,238	56.76%
80+	5,633,398	2,501,274	44.40%	37.76%	51.04%	3,132,124	55.60%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,563,856	3,535,216	46.74%	42.37%	51.11%	4,028,640	53.26%
Other Hispanic	4,965,594	1,835,340	36.96%	28.53%	45.39%	3,130,254	63.04%
Non-Hispanic White	71,500,163	32,645,762	45.66%	41.85%	49.46%	38,854,401	54.34%
Non-Hispanic Black	12,049,710	3,863,130	32.06%	29.14%	34.98%	8,186,580	67.94%
Other Race - Including Multi-Racial	5,225,573	1,694,691	32.43%	20.90%	43.96%	3,530,882	67.57%
Education	-	-	-	-	-	-	-
Missing	2,863,940	950,237	33.18%	18.74%	47.62%	1,913,703	66.82%
Less Than 9th Grade	6,416,781	3,263,040	50.85%	42.41%	59.30%	3,153,741	49.15%
9-11th Grade	13,305,373	5,919,271	44.49%	35.88%	53.09%	7,386,102	55.51%
High School Grad/GED or Equivalent	78,707,186	33,429,975	42.47%	39.47%	45.48%	45,277,211	57.53%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,342,297	314,106	23.40%	11.35%	35.45%	1,028,190	76.60%
Married	57,596,410	26,289,858	45.64%	41.92%	49.37%	31,306,552	54.36%
Widowed	9,356,585	4,751,627	50.78%	45.67%	55.90%	4,604,958	49.22%
Divorced	12,606,729	5,862,932	46.51%	39.51%	53.50%	6,743,797	53.49%
Separated	2,378,944	1,146,885	48.21%	33.86%	62.56%	1,232,059	51.79%
Never Married	11,834,836	3,193,357	26.98%	21.15%	32.82%	8,641,479	73.02%
Living with Partner	6,162,712	1,988,990	32.27%	20.50%	44.05%	4,173,722	67.73%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	7,061,696	2,666,791	37.76%	29.48%	46.05%	4,394,905	62.24%
Poverty income ratio less than 1	13,770,564	5,577,123	40.50%	34.46%	46.54%	8,193,441	59.50%
Poverty income ratio 1 or greater	80,472,636	35,330,225	43.90%	40.12%	47.69%	45,142,411	56.10%
Body mass index	-	-	-	-	-	-	-
Missing	963,961	537,545	55.76%	36.34%	75.19%	426,416	44.24%
Normal: less than 25	35,622,801	12,125,743	34.04%	28.74%	39.34%	23,497,058	65.96%
Overweight: 25.0 - 29.9	29,311,983	12,310,041	42.00%	35.84%	48.16%	17,001,942	58.00%
Obese: 30 or higher	35,406,151	18,600,809	52.54%	47.46%	57.61%	16,805,342	47.46%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,619,352	1,150,933	31.80%	20.66%	42.94%	2,468,419	68.20%
Glycohemoglobin < 5.7%	68,246,278	27,395,711	40.14%	35.76%	44.53%	40,850,567	59.86%
Glycohemoglobin 5.7% - 6.4%	22,438,338	11,037,175	49.19%	43.31%	55.07%	11,401,163	50.81%
Glycohemoglobin 6.5% or higher	7,000,928	3,990,320	57.00%	48.19%	65.80%	3,010,608	43.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,304,896	43,574,138	43.01%	39.89%	46.14%	57,730,757	56.99%
Diabetes	-	-	-	-	-	-	-
Yes	9,453,189	5,550,032	58.71%	50.48%	66.94%	3,903,157	41.29%
No	91,851,707	38,024,106	41.40%	37.72%	45.07%	53,827,601	58.60%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,244,498	19,612,973	47.55%	43.84%	51.27%	21,631,525	52.45%
No	59,996,215	23,961,165	39.94%	35.52%	44.35%	36,035,050	60.06%
Refused	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	32,871,586	17,323,777	52.70%	47.93%	57.47%	15,547,809	47.30%
No	68,349,726	26,250,362	38.41%	34.87%	41.95%	42,099,364	61.59%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,674,366	6,171,749	52.87%	47.62%	58.11%	5,502,617	47.13%
Mobility limitations	10,745,179	5,628,671	52.38%	44.14%	60.63%	5,116,507	47.62%
No mobility limitations	20,436,026	9,028,727	44.18%	37.53%	50.83%	11,407,300	55.82%
N/A - Respondent not asked mobility limitation questions	58,449,325	22,744,991	38.91%	35.40%	42.43%	35,704,334	61.09%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,361,300	3,720,848	50.55%	39.22%	61.87%	3,640,452	49.45%
No	93,854,217	39,763,911	42.37%	39.28%	45.46%	54,090,306	57.63%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	627,295	324,720	51.77%	35.76%	67.77%	302,575	48.23%
2 or more times a day	9,905,116	5,231,083	52.81%	43.47%	62.15%	4,674,034	47.19%
Once a day	9,682,887	3,913,332	40.41%	34.95%	45.88%	5,769,555	59.59%
2 or more time a week	7,646,398	3,844,144	50.27%	40.45%	60.10%	3,802,254	49.73%
Once a week	7,022,920	3,608,143	51.38%	39.95%	62.81%	3,414,777	48.62%
1-3 times a month	14,356,537	7,135,871	49.70%	43.53%	55.88%	7,220,666	50.30%
Never	51,821,562	19,466,051	37.56%	33.36%	41.77%	32,355,511	62.44%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	627,295	324,720	51.77%	35.76%	67.77%	302,575	48.23%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	7,646,398	3,844,144	50.27%	32.46%	94.33%	3,802,254	49.73%
Once a week	*	*	*	*	*	*	*
1-3 times a month	14,356,537	7,135,871	49.70%	58.28%	90.01%	7,220,666	50.30%
Never	51,821,562	19,466,051	37.56%	39.15%	45.33%	32,355,511	62.44%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,304,896	43,574,138	43.01%	39.89%	46.14%	57,730,757	56.99%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	432,110	99,537	23.04%	(13.15%)	59.22%	332,573	76.96%
Younger than 30	3,069,584	1,125,672	36.67%	26.23%	47.11%	1,943,912	63.33%
30 to 34	3,711,338	1,246,661	33.59%	26.81%	40.37%	2,464,677	66.41%
35 to 39	5,113,159	2,595,857	50.77%	38.36%	63.17%	2,517,302	49.23%
40 to 44	7,568,224	4,075,874	53.86%	41.67%	66.04%	3,492,350	46.14%
45 to 49	11,650,306	5,902,598	50.66%	44.55%	56.78%	5,747,709	49.34%
50 to 54	10,341,790	5,183,409	50.12%	41.27%	58.97%	5,158,381	49.88%
55 or older	5,131,554	2,615,254	50.96%	41.45%	60.48%	2,516,300	49.04%
Don't know	238,745	103,276	43.26%	10.59%	75.92%	135,469	56.74%
N/A - Had period in last 12 months	54,048,084	20,626,000	38.16%	34.29%	42.03%	33,422,083	61.84%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,962,970	4,300,728	25.35%	18.51%	32.19%	12,662,242	74.65%
1	13,401,467	4,324,010	32.27%	24.99%	39.54%	9,077,457	67.73%
2	22,728,804	10,327,253	45.44%	38.69%	52.19%	12,401,551	54.56%
More than 2	47,931,106	24,622,147	51.37%	48.11%	54.63%	23,308,959	48.63%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	674,220	231,660	34.36%	18.20%	50.51%	442,560	65.64%
None	32,118,654	9,434,212	29.37%	25.02%	33.72%	22,684,442	70.63%
1	17,758,915	7,482,869	42.14%	36.76%	47.52%	10,276,046	57.86%
2	23,038,524	11,944,848	51.85%	45.40%	58.30%	11,093,677	48.15%
More than 2	27,700,262	14,466,230	52.22%	47.83%	56.62%	13,234,031	47.78%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	964,977	302,685	31.37%	15.88%	46.86%	662,293	68.63%
None	25,256,630	12,459,932	49.33%	42.77%	55.90%	12,796,697	50.67%
1	9,965,596	4,166,993	41.81%	35.09%	48.54%	5,798,602	58.19%
2	5,549,270	2,055,989	37.05%	28.11%	45.99%	3,493,282	62.95%
More than 2	2,981,345	1,399,863	46.95%	35.26%	58.65%	1,581,482	53.05%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	56,572,758	23,174,357	40.96%	36.97%	44.96%	33,398,401	59.04%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	21,543,739	5,902,879	27.40%	20.20%	34.60%	15,640,860	72.60%
Yes	13,130,418	6,751,371	51.42%	40.64%	62.20%	6,379,047	48.58%
No	66,451,019	30,875,242	46.46%	43.42%	49.50%	35,575,777	53.54%
Don't know	179,719	44,645	24.84%	(11.54%)	61.22%	135,073	75.16%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	21,723,458	5,947,525	27.38%	20.25%	34.51%	15,775,933	72.62%
35 years old or younger	12,104,834	6,233,281	51.49%	40.47%	62.52%	5,871,553	48.51%
36 years or older	932,800	518,090	55.54%	30.05%	81.03%	414,709	44.46%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	66,451,019	30,875,242	46.46%	43.42%	49.50%	35,575,777	53.54%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,304,896	43,574,138	43.01%	39.89%	46.14%	57,730,757	56.99%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	4,580,770	1,602,152	34.98%	18.66%	51.29%	2,978,618	65.02%
29 years old or younger	55,911,182	27,873,639	49.85%	46.21%	53.50%	28,037,543	50.15%
30 - 40 years old	5,156,343	2,875,571	55.77%	45.52%	66.02%	2,280,772	44.23%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	16,962,970	4,300,728	25.35%	18.51%	32.19%	12,662,242	74.65%
Only 1 live birth - see age at last live birth	18,125,337	6,822,202	37.64%	30.98%	44.30%	11,303,134	62.36%
No pregnancies resulting in live births	491,126	53,593	10.91%	(7.18%)	29.01%	437,533	89.09%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	5,813,479	2,121,717	36.50%	23.25%	49.75%	3,691,762	63.50%
29 years old or younger	46,368,089	21,522,115	46.42%	42.79%	50.05%	24,845,974	53.58%
30 - 40 years old	29,501,329	14,499,324	49.15%	45.06%	53.24%	15,002,005	50.85%
41 years or older	1,806,507	885,016	48.99%	30.51%	67.47%	921,491	51.01%
Refused/Don't know	361,396	191,646	53.03%	27.19%	78.87%	169,749	46.97%
N/A - Never pregnant	16,962,970	4,300,728	25.35%	18.51%	32.19%	12,662,242	74.65%
No pregnancies resulting in live births	491,126	53,593	10.91%	(7.18%)	29.01%	437,533	89.09%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,906,168	11,219,082	51.21%	44.80%	57.63%	10,687,086	48.79%
No	79,240,929	32,317,563	40.78%	37.01%	44.56%	46,923,366	59.22%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	6,125,811	2,525,405	41.23%	32.50%	49.96%	3,600,406	58.77%
36 - 50 years old	12,738,019	7,127,515	55.95%	47.34%	64.57%	5,610,504	44.05%
51 years or older	2,470,853	1,364,112	55.21%	40.29%	70.12%	1,106,741	44.79%
Refused/Don't know	571,486	202,051	35.36%	5.95%	64.76%	369,435	64.64%
N/A - Did not have procedure	79,240,929	32,317,563	40.78%	37.01%	44.56%	46,923,366	59.22%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,803,240	772,445	42.84%	20.13%	65.55%	1,030,795	57.16%
Yes	11,812,694	6,713,473	56.83%	46.88%	66.79%	5,099,221	43.17%
No	87,028,222	35,893,923	41.24%	37.60%	44.89%	51,134,299	58.76%
Don't know	660,739	194,297	29.41%	8.96%	49.85%	466,442	70.59%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,463,979	966,742	39.23%	19.56%	58.91%	1,497,238	60.77%
35 years old or younger	2,315,265	864,424	37.34%	25.48%	49.19%	1,450,841	62.66%
36 - 50 years old	6,648,470	4,375,715	65.82%	52.83%	78.80%	2,272,755	34.18%
51 years or older	2,584,749	1,370,647	53.03%	36.16%	69.89%	1,214,101	46.97%
Refused/Don't know	264,211	102,687	38.87%	4.34%	73.39%	161,524	61.13%
N/A - Did not have procedure	87,028,222	35,893,923	41.24%	37.60%	44.89%	51,134,299	58.76%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,304,896	43,574,138	43.01%	39.89%	46.14%	57,730,757	56.99%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,329,357	11,058,207	51.85%	45.03%	58.66%	10,271,149	48.15%
No	79,817,740	32,478,438	40.69%	36.92%	44.47%	47,339,303	59.31%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,709,924	41,565,171	42.54%	40.44%	44.64%	56,144,753	57.46%
Age at screening	-	-	-	-	-	-	-
10-19	353,109	199,400	56.47%	31.66%	81.28%	153,709	43.53%
20-29	12,072,924	3,150,713	26.10%	21.48%	30.71%	8,922,211	73.90%
30-39	15,950,025	5,460,004	34.23%	28.80%	39.66%	10,490,021	65.77%
40-49	19,772,641	8,461,428	42.79%	36.74%	48.84%	11,311,213	57.21%
50-59	20,729,616	10,595,004	51.11%	44.03%	58.19%	10,134,612	48.89%
60-64	7,767,115	3,593,050	46.26%	37.30%	55.22%	4,174,065	53.74%
65-69	6,636,364	2,846,674	42.90%	33.74%	52.05%	3,789,690	57.10%
70-74	5,133,707	2,783,570	54.22%	45.68%	62.76%	2,350,138	45.78%
75-79	4,039,012	2,155,087	53.36%	45.48%	61.24%	1,883,925	46.64%
80+	5,255,411	2,320,241	44.15%	37.20%	51.10%	2,935,170	55.85%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,365,117	3,258,435	44.24%	40.07%	48.41%	4,106,682	55.76%
Other Hispanic	4,864,518	2,286,436	47.00%	41.27%	52.73%	2,578,082	53.00%
Non-Hispanic White	67,854,406	30,321,787	44.69%	41.62%	47.75%	37,532,619	55.31%
Non-Hispanic Black	11,442,102	3,376,104	29.51%	25.12%	33.89%	8,065,999	70.49%
Other Race - Including Multi-Racial	6,183,781	2,322,409	37.56%	26.17%	48.95%	3,861,371	62.44%
Education	-	-	-	-	-	-	-
Missing	2,266,489	1,124,284	49.60%	32.87%	66.34%	1,142,205	50.40%
Less Than 9th Grade	5,812,537	2,766,828	47.60%	39.55%	55.65%	3,045,709	52.40%
9-11th Grade	11,488,547	5,087,699	44.28%	38.45%	50.12%	6,400,847	55.72%
High School Grad/GED or Equivalent	77,940,997	32,476,919	41.67%	39.48%	43.85%	45,464,078	58.33%
Refused	*	*	*	*	*	*	*
Don't Know	156,713	64,799	41.35%	(0.61%)	83.31%	91,914	58.65%
Marital status	-	-	-	-	-	-	-
Missing	353,109	199,400	56.47%	31.66%	81.28%	153,709	43.53%
Married	55,939,626	24,874,488	44.47%	41.38%	47.55%	31,065,138	55.53%
Widowed	8,828,269	4,090,678	46.34%	40.40%	52.27%	4,737,591	53.66%
Divorced	11,116,682	4,648,101	41.81%	34.96%	48.67%	6,468,582	58.19%
Separated	2,211,003	812,753	36.76%	17.75%	55.77%	1,398,249	63.24%
Never Married	11,937,752	4,118,534	34.50%	28.02%	40.98%	7,819,217	65.50%
Living with Partner	7,120,836	2,657,020	37.31%	27.82%	46.81%	4,463,817	62.69%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,365,743	2,647,636	41.59%	34.18%	49.00%	3,718,107	58.41%
Poverty income ratio less than 1	13,921,006	5,923,419	42.55%	37.83%	47.27%	7,997,586	57.45%
Poverty income ratio 1 or greater	77,423,176	32,994,116	42.62%	40.19%	45.04%	44,429,060	57.38%
Body mass index	-	-	-	-	-	-	-
Missing	432,161	104,005	24.07%	(1.55%)	49.68%	328,156	75.93%
Normal: less than 25	34,112,320	11,270,152	33.04%	29.55%	36.52%	22,842,169	66.96%
Overweight: 25.0 - 29.9	27,215,370	11,673,193	42.89%	39.29%	46.50%	15,542,176	57.11%
Obese: 30 or higher	35,950,074	18,517,821	51.51%	48.07%	54.95%	17,432,252	48.49%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,219,727	1,376,114	42.74%	33.38%	52.10%	1,843,612	57.26%
Glycohemoglobin < 5.7%	62,621,445	23,931,482	38.22%	35.87%	40.56%	38,689,964	61.78%
Glycohemoglobin 5.7% - 6.4%	25,124,325	12,584,363	50.09%	45.77%	54.40%	12,539,962	49.91%
Glycohemoglobin 6.5% or higher	6,744,427	3,673,212	54.46%	46.51%	62.41%	3,071,215	45.54%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,709,924	41,565,171	42.54%	40.44%	44.64%	56,144,753	57.46%
Diabetes	-	-	-	-	-	-	-
Yes	8,795,160	4,941,122	56.18%	49.10%	63.26%	3,854,037	43.82%
No	88,914,765	36,624,049	41.19%	39.06%	43.32%	52,290,716	58.81%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	39,087,641	17,189,244	43.98%	41.21%	46.75%	21,898,397	56.02%
No	58,622,284	24,375,928	41.58%	38.59%	44.58%	34,246,356	58.42%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,125,031	15,658,321	51.98%	47.72%	56.23%	14,466,710	48.02%
No	67,523,526	25,889,369	38.34%	36.21%	40.47%	41,634,158	61.66%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,344,219	5,512,998	48.60%	42.65%	54.55%	5,831,221	51.40%
Mobility limitations	9,625,939	5,546,085	57.62%	52.72%	62.52%	4,079,854	42.38%
No mobility limitations	20,668,087	9,189,061	44.46%	40.02%	48.90%	11,479,026	55.54%
N/A - Respondent not asked mobility limitation questions	56,071,679	21,317,028	38.02%	35.51%	40.53%	34,754,652	61.98%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,026,447	4,034,403	57.42%	47.50%	67.33%	2,992,044	42.58%
No	90,607,539	37,461,743	41.35%	39.10%	43.59%	53,145,796	58.65%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	465,612	210,458	45.20%	8.84%	81.56%	255,153	54.80%
2 or more times a day	9,005,246	4,953,414	55.01%	46.24%	63.77%	4,051,832	44.99%
Once a day	8,275,588	3,644,732	44.04%	37.53%	50.55%	4,630,856	55.96%
2 or more time a week	8,108,175	4,467,470	55.10%	43.17%	67.03%	3,640,705	44.90%
Once a week	6,154,232	2,649,682	43.05%	32.08%	54.03%	3,504,551	56.95%
1-3 times a month	13,225,964	6,040,827	45.67%	38.47%	52.88%	7,185,137	54.33%
Never	52,378,637	19,557,241	37.34%	34.24%	40.43%	32,821,397	62.66%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	465,612	210,458	45.20%	15.91%	86.47%	255,153	54.80%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	8,108,175	4,467,470	55.10%	29.95%	99.21%	3,640,705	44.90%
Once a week	6,154,232	2,649,682	43.05%	38.79%	106.00%	3,504,551	56.95%
1-3 times a month	13,225,964	6,040,827	45.67%	40.67%	79.85%	7,185,137	54.33%
Never	52,378,637	19,557,241	37.34%	39.83%	44.09%	32,821,397	62.66%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,709,924	41,565,171	42.54%	40.44%	44.64%	56,144,753	57.46%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	624,425	182,138	29.17%	3.60%	54.74%	442,288	70.83%
Younger than 30	2,882,766	1,390,166	48.22%	35.99%	60.46%	1,492,600	51.78%
30 to 34	3,254,943	1,667,940	51.24%	41.45%	61.04%	1,587,003	48.76%
35 to 39	4,541,323	2,132,891	46.97%	32.60%	61.33%	2,408,432	53.03%
40 to 44	7,445,396	3,834,630	51.50%	43.55%	59.45%	3,610,766	48.50%
45 to 49	9,847,084	5,016,951	50.95%	44.73%	57.16%	4,830,133	49.05%
50 to 54	13,669,312	7,197,564	52.65%	45.77%	59.54%	6,471,748	47.35%
55 or older	5,142,489	2,247,737	43.71%	31.24%	56.18%	2,894,752	56.29%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	50,043,739	17,811,880	35.59%	33.78%	37.40%	32,231,859	64.41%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	17,565,210	4,012,962	22.85%	18.97%	26.73%	13,552,247	77.15%
1	12,038,159	4,743,482	39.40%	30.78%	48.03%	7,294,677	60.60%
2	23,714,828	10,672,802	45.00%	39.81%	50.20%	13,042,027	55.00%
More than 2	44,287,373	22,062,090	49.82%	46.49%	53.14%	22,225,284	50.18%
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	310,780	147,392	47.43%	9.87%	84.98%	163,388	52.57%
None	32,911,743	9,409,713	28.59%	26.17%	31.02%	23,502,029	71.41%
1	14,225,353	6,701,306	47.11%	38.61%	55.60%	7,524,047	52.89%
2	23,367,578	12,237,832	52.37%	47.21%	57.53%	11,129,745	47.63%
More than 2	26,894,471	13,068,928	48.59%	43.94%	53.25%	13,825,543	51.41%
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	779,924	477,152	61.18%	39.27%	83.09%	302,772	38.82%
None	25,135,681	12,464,366	49.59%	44.11%	55.07%	12,671,315	50.41%
1	9,753,274	4,283,563	43.92%	35.24%	52.60%	5,469,712	56.08%
2	5,288,336	2,177,127	41.17%	33.47%	48.87%	3,111,208	58.83%
More than 2	2,314,074	878,903	37.98%	25.29%	50.67%	1,435,170	62.02%
N/A - Vaginal delivery/deliveries only	54,438,636	21,284,061	39.10%	36.26%	41.93%	33,154,575	60.90%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	22,992,485	5,961,702	25.93%	22.40%	29.46%	17,030,783	74.07%
Yes	14,458,970	6,966,434	48.18%	41.77%	54.59%	7,492,537	51.82%
No	59,893,990	28,431,631	47.47%	45.06%	49.88%	31,462,359	52.53%
Don't know	364,479	205,405	56.36%	23.97%	88.74%	159,074	43.64%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	23,356,964	6,167,107	26.40%	22.90%	29.91%	17,189,857	73.60%
35 years old or younger	13,118,409	6,388,489	48.70%	42.40%	54.99%	6,729,919	51.30%
36 years or older	1,284,225	577,944	45.00%	27.86%	62.15%	706,280	55.00%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	59,893,990	28,431,631	47.47%	45.06%	49.88%	31,462,359	52.53%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,709,924	41,565,171	42.54%	40.44%	44.64%	56,144,753	57.46%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	5,427,276	1,948,740	35.91%	21.86%	49.96%	3,478,536	64.09%
29 years old or younger	53,934,426	26,595,475	49.31%	46.11%	52.51%	27,338,951	50.69%
30 - 40 years old	5,231,391	2,256,437	43.13%	31.14%	55.12%	2,974,954	56.87%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,565,210	4,012,962	22.85%	18.97%	26.73%	13,552,247	77.15%
Only 1 live birth - see age at last live birth	15,122,928	6,481,179	42.86%	36.27%	49.45%	8,641,748	57.14%
No pregnancies resulting in live births	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,912,927	2,512,359	36.34%	23.23%	49.45%	4,400,568	63.66%
29 years old or younger	41,075,647	18,932,870	46.09%	42.85%	49.34%	22,142,776	53.91%
30 - 40 years old	29,406,142	14,850,293	50.50%	48.07%	52.93%	14,555,850	49.50%
41 years or older	2,166,264	862,532	39.82%	23.77%	55.86%	1,303,733	60.18%
Refused/Don't know	254,181	123,778	48.70%	5.53%	91.87%	130,404	51.30%
N/A - Never pregnant	17,565,210	4,012,962	22.85%	18.97%	26.73%	13,552,247	77.15%
No pregnancies resulting in live births	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,503,325	11,106,953	54.17%	48.13%	60.21%	9,396,371	45.83%
No	77,156,351	30,458,218	39.48%	37.45%	41.50%	46,698,132	60.52%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	5,391,230	2,928,483	54.32%	43.60%	65.04%	2,462,747	45.68%
36 - 50 years old	11,013,712	6,136,538	55.72%	47.92%	63.51%	4,877,174	44.28%
51 years or older	3,681,428	1,791,666	48.67%	39.98%	57.36%	1,889,762	51.33%
Refused/Don't know	416,955	250,266	60.02%	34.68%	85.37%	166,689	39.98%
N/A - Did not have procedure	77,156,351	30,458,218	39.48%	37.45%	41.50%	46,698,132	60.52%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,954,499	1,167,397	59.73%	45.06%	74.40%	787,103	40.27%
Yes	10,117,822	5,650,720	55.85%	50.32%	61.38%	4,467,102	44.15%
No	84,902,123	34,370,000	40.48%	37.96%	43.01%	50,532,123	59.52%
Refused	*	*	*	*	*	*	*
Don't know	723,930	377,055	52.08%	27.50%	76.67%	346,874	47.92%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,689,979	1,544,452	57.42%	46.87%	67.96%	1,145,528	42.59%
35 years old or younger	2,196,357	1,140,620	51.93%	34.81%	69.05%	1,055,737	48.07%
36 - 50 years old	5,245,669	3,156,611	60.18%	49.84%	70.51%	2,089,059	39.82%
51 years or older	2,452,490	1,210,967	49.38%	37.73%	61.02%	1,241,522	50.62%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	84,902,123	34,370,000	40.48%	37.96%	43.01%	50,532,123	59.52%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02 - Prevalence of stress urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,709,924	41,565,171	42.54%	40.44%	44.64%	56,144,753	57.46%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,303,766	11,070,262	54.52%	48.55%	60.50%	9,233,504	45.48%
No	77,355,910	30,494,910	39.42%	37.38%	41.46%	46,861,000	60.58%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.