

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	2,482,684	144,110	5.80%	416,824	16.79%	527,134	21.23%	1,394,616	56.17%	0	0.00%	0	0.00%
30-39	6,288,693	851,562	13.54%	978,988	15.57%	1,585,285	25.21%	2,866,676	45.58%	6,183	0.10%	0	0.00%
40-49	9,216,571	992,470	10.77%	1,742,472	18.91%	2,261,626	24.54%	4,220,003	45.79%	0	0.00%	0	0.00%
50-59	8,577,959	1,542,606	17.98%	2,084,006	24.29%	1,993,637	23.24%	2,957,710	34.48%	0	0.00%	0	0.00%
60-64	2,586,198	625,809	24.20%	691,412	26.73%	701,612	27.13%	567,366	21.94%	0	0.00%	0	0.00%
65-69	2,596,196	734,341	28.29%	696,334	26.82%	583,166	22.46%	582,355	22.43%	0	0.00%	0	0.00%
70-74	1,994,382	573,432	28.75%	351,793	17.64%	630,007	31.59%	439,149	22.02%	0	0.00%	0	0.00%
75-79	2,333,591	792,621	33.97%	547,418	23.46%	633,718	27.16%	359,834	15.42%	0	0.00%	0	0.00%
80-84	1,564,983	508,121	32.47%	279,624	17.87%	321,442	20.54%	431,921	27.60%	23,874	1.53%	0	0.00%
85+	1,020,671	384,905	37.71%	221,945	21.75%	259,397	25.41%	130,468	12.78%	23,956	2.35%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	2,179,964	268,710	12.33%	413,675	18.98%	505,646	23.20%	991,933	45.50%	0	0.00%	0	0.00%
Other Hispanic	1,550,701	445,892	28.75%	393,186	25.36%	349,340	22.53%	338,410	21.82%	23,874	1.54%	0	0.00%
Non-Hispanic White	30,783,407	5,627,374	18.28%	6,352,950	20.64%	7,562,373	24.57%	11,210,572	36.42%	30,139	0.10%	0	0.00%
Non-Hispanic Black	2,438,829	494,374	20.27%	518,931	21.28%	685,993	28.13%	739,531	30.32%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	1,736,672	333,948	19.23%	332,075	19.12%	400,997	23.09%	669,652	38.56%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,025,184	137,950	13.46%	330,438	32.23%	267,900	26.13%	288,896	28.18%	0	0.00%	0	0.00%
Less Than 9th Grade	2,732,725	783,732	28.68%	626,318	22.92%	943,958	34.54%	378,718	13.86%	0	0.00%	0	0.00%
9-11th Grade	4,552,708	1,045,450	22.96%	1,032,389	22.68%	1,082,937	23.79%	1,391,931	30.57%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	30,361,756	5,203,166	17.14%	6,017,247	19.82%	7,209,553	23.75%	11,877,777	39.12%	54,013	0.18%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	24,615,627	3,557,339	14.45%	5,279,613	21.45%	6,201,485	25.19%	9,571,007	38.88%	6,183	0.03%	0	0.00%
Widowed	4,910,232	1,806,261	36.79%	1,135,139	23.12%	1,180,603	24.04%	764,274	15.56%	23,956	0.49%	0	0.00%
Divorced	3,732,044	903,909	24.22%	675,584	18.10%	802,156	21.49%	1,326,521	35.54%	23,874	0.64%	0	0.00%
Separated	1,142,774	255,364	22.35%	141,965	12.42%	293,459	25.68%	451,987	39.55%	0	0.00%	0	0.00%
Never Married	2,968,369	566,254	19.08%	485,096	16.34%	691,506	23.30%	1,225,514	41.29%	0	0.00%	0	0.00%
Living with Partner	1,209,554	58,031	4.80%	293,419	24.26%	298,133	24.65%	559,972	46.30%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,009,449	592,231	29.47%	409,066	20.36%	414,788	20.64%	569,408	28.34%	23,956	1.19%	0	0.00%
Poverty income ratio less than 1	4,714,007	1,379,265	29.26%	1,173,354	24.89%	967,154	20.52%	1,194,234	25.33%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	31,966,117	5,198,801	16.26%	6,428,396	20.11%	8,122,406	25.41%	12,186,457	38.12%	30,057	0.09%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,025,765	1,138,085	56.18%	148,668	7.34%	337,177	16.64%	401,834	19.84%	0	0.00%	0	0.00%
Normal: less than 25	11,696,016	1,238,179	10.59%	1,979,962	16.93%	2,859,573	24.45%	5,618,302	48.04%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	10,505,442	1,711,672	16.29%	2,108,680	20.07%	2,741,486	26.10%	3,889,592	37.02%	54,013	0.51%	0	0.00%
Obese: 30 or higher	14,462,350	3,082,361	21.31%	3,773,506	26.09%	3,566,113	24.66%	4,040,370	27.94%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	891,814	358,700	40.22%	30,077	3.37%	176,018	19.74%	327,019	36.67%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	29,047,153	4,467,989	15.38%	5,727,491	19.72%	7,215,766	24.84%	11,605,767	39.95%	30,139	0.10%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	6,822,228	1,762,559	25.84%	1,716,899	25.17%	1,603,237	23.50%	1,715,660	25.15%	23,874	0.35%	0	0.00%
Glycohemoglobin 6.5% or higher	1,928,378	581,049	30.13%	536,349	27.81%	509,327	26.41%	301,652	15.64%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,675,310	923,366	34.51%	679,789	25.41%	692,091	25.87%	380,063	14.21%	0	0.00%	0	0.00%
No	36,014,263	6,246,931	17.35%	7,331,027	20.36%	8,812,257	24.47%	13,570,035	37.68%	54,013	0.15%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	18,772,437	3,952,623	21.06%	4,160,106	22.16%	3,898,531	20.77%	6,731,038	35.86%	30,139	0.16%	0	0.00%
No	19,848,518	3,217,674	16.21%	3,850,709	19.40%	5,547,332	27.95%	7,208,929	36.32%	23,874	0.12%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	14,053,704	3,438,868	24.47%	3,367,969	23.96%	3,391,208	24.13%	3,801,646	27.05%	54,013	0.38%	0	0.00%
No	24,498,583	3,731,429	15.23%	4,642,846	18.95%	6,022,465	24.58%	10,101,843	41.23%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,379,061	1,337,912	39.59%	741,459	21.94%	941,107	27.85%	358,582	10.61%	0	0.00%	0	0.00%
Mobility limitations	7,232,538	2,451,522	33.90%	2,062,106	28.51%	1,516,519	20.97%	1,178,436	16.29%	23,956	0.33%	0	0.00%
No mobility limitations	7,921,631	1,167,913	14.74%	1,490,095	18.81%	2,262,834	28.57%	2,976,916	37.58%	23,874	0.30%	0	0.00%
N/A - Respondent not asked mobility limitation questions	20,156,342	2,212,950	10.98%	3,717,156	18.44%	4,783,888	23.73%	9,436,165	46.81%	6,183	0.03%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	2,120,605	176,233	8.31%	658,020	31.03%	362,720	17.10%	923,632	43.56%	0	0.00%	0	0.00%
30 to 34	1,957,558	465,436	23.78%	486,380	24.85%	587,249	30.00%	418,492	21.38%	0	0.00%	0	0.00%
35 to 39	2,663,202	785,594	29.50%	689,682	25.90%	864,775	32.47%	323,151	12.13%	0	0.00%	0	0.00%
40 to 44	2,951,461	804,350	27.25%	356,564	12.08%	857,424	29.05%	933,124	31.62%	0	0.00%	0	0.00%
45 to 49	4,380,859	1,167,188	26.64%	1,033,239	23.59%	957,159	21.85%	1,223,273	27.92%	0	0.00%	0	0.00%
50 to 54	5,225,821	1,358,465	26.00%	1,270,604	24.31%	1,151,715	22.04%	1,445,038	27.65%	0	0.00%	0	0.00%
55 or older	1,555,049	418,095	26.89%	275,297	17.70%	280,933	18.07%	532,894	34.27%	47,830	3.08%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	17,756,974	1,994,937	11.23%	3,180,030	17.91%	4,442,373	25.02%	8,133,451	45.80%	6,183	0.03%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	524,616	91,501	17.44%	119,074	22.70%	152,629	29.09%	161,412	30.77%	0	0.00%	0	0.00%
None	2,989,695	390,260	13.05%	510,345	17.07%	546,245	18.27%	1,542,845	51.61%	0	0.00%	0	0.00%
1	4,079,139	221,156	5.42%	864,370	21.19%	1,120,550	27.47%	1,873,063	45.92%	0	0.00%	0	0.00%
2	7,965,461	1,389,731	17.45%	1,349,991	16.95%	2,249,855	28.25%	2,975,884	37.36%	0	0.00%	0	0.00%
More than 2	23,130,661	5,077,649	21.95%	5,167,036	22.34%	5,435,070	23.50%	7,396,893	31.98%	54,013	0.23%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	674,807	99,128	14.69%	189,449	28.07%	177,321	26.28%	208,909	30.96%	0	0.00%	0	0.00%
29 years old or younger	26,583,303	5,696,656	21.43%	5,579,112	20.99%	6,511,596	24.50%	8,741,926	32.89%	54,013	0.20%	0	0.00%
30 - 40 years old	1,448,759	276,307	19.07%	274,932	18.98%	567,632	39.18%	329,888	22.77%	0	0.00%	0	0.00%
N/A - Never pregnant	2,989,695	390,260	13.05%	510,345	17.07%	546,245	18.27%	1,542,845	51.61%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	5,609,434	596,375	10.63%	1,278,439	22.79%	1,530,847	27.29%	2,203,773	39.29%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,383,575	111,570	8.06%	178,539	12.90%	170,708	12.34%	922,758	66.69%	0	0.00%	0	0.00%
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,991,677	810,464	27.09%	725,820	24.26%	689,681	23.05%	765,712	25.59%	0	0.00%	0	0.00%
29 years old or younger	17,519,528	2,988,388	17.06%	3,867,641	22.08%	4,543,611	25.93%	6,113,705	34.90%	6,183	0.04%	0	0.00%
30 - 40 years old	12,818,285	2,641,420	20.61%	2,452,612	19.13%	3,319,012	25.89%	4,381,367	34.18%	23,874	0.19%	0	0.00%
41 years or older	943,923	228,195	24.18%	275,859	29.22%	215,602	22.84%	200,312	21.22%	23,956	2.54%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,989,695	390,260	13.05%	510,345	17.07%	546,245	18.27%	1,542,845	51.61%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,383,575	111,570	8.06%	178,539	12.90%	170,708	12.34%	922,758	66.69%	0	0.00%	0	0.00%
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	13,253,555	1,306,082	9.85%	2,260,930	17.06%	3,324,670	25.09%	6,361,874	48.00%	0	0.00%	0	0.00%
Yes	12,073,527	3,184,426	26.38%	2,943,722	24.38%	3,198,514	26.49%	2,722,910	22.55%	23,956	0.20%	0	0.00%
No	13,362,490	2,679,790	20.05%	2,806,164	21.00%	2,981,165	22.31%	4,865,315	36.41%	30,057	0.22%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	13,286,298	1,306,082	9.83%	2,260,930	17.02%	3,357,413	25.27%	6,361,874	47.88%	0	0.00%	0	0.00%
35 years old or younger	4,460,670	873,317	19.58%	1,275,878	28.60%	1,024,435	22.97%	1,287,040	28.85%	0	0.00%	0	0.00%
36 - 50 years old	6,031,417	1,858,006	30.81%	1,162,823	19.28%	1,801,949	29.88%	1,208,640	20.04%	0	0.00%	0	0.00%
51 years or older	1,404,236	453,103	32.27%	480,032	34.18%	268,446	19.12%	178,699	12.73%	23,956	1.71%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	13,362,490	2,679,790	20.05%	2,806,164	21.00%	2,981,165	22.31%	4,865,315	36.41%	30,057	0.22%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2001-2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	38,689,573	7,170,297	18.53%	8,010,816	20.71%	9,504,348	24.57%	13,950,099	36.06%	54,013	0.14%	0	0.0
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,428,486	717,543	20.93%	386,245	11.27%	870,569	25.39%	1,454,130	42.41%	0	0.00%	0	0%
Yes	12,073,527	3,184,426	26.38%	2,943,722	24.38%	3,198,514	26.49%	2,722,910	22.55%	23,956	0.20%	0	0%
No	23,187,559	3,268,329	14.10%	4,680,849	20.19%	5,435,266	23.44%	9,773,059	42.15%	30,057	0.13%	0	0%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	3,210,843	294,653	9.18%	568,614	17.71%	700,006	21.80%	1,647,570	51.31%	0	0.00%	0	0.00%
30-39	6,117,018	746,248	12.20%	1,362,815	22.28%	1,969,277	32.19%	2,038,678	33.33%	0	0.00%	0	0.00%
40-49	9,374,366	1,466,048	15.64%	1,525,299	16.27%	2,840,078	30.30%	3,542,941	37.79%	0	0.00%	0	0.00%
50-59	8,687,563	1,268,963	14.61%	2,118,877	24.39%	1,856,841	21.37%	3,442,881	39.63%	0	0.00%	0	0.00%
60-64	2,250,785	380,270	16.89%	440,375	19.57%	464,022	20.62%	966,117	42.92%	0	0.00%	0	0.00%
65-69	3,369,560	874,107	25.94%	717,301	21.29%	701,573	20.82%	1,076,578	31.95%	0	0.00%	0	0.00%
70-74	2,454,011	732,234	29.84%	808,649	32.95%	194,321	7.92%	718,807	29.29%	0	0.00%	0	0.00%
75-79	3,007,491	913,451	30.37%	657,768	21.87%	744,852	24.77%	691,420	22.99%	0	0.00%	0	0.00%
80-84	1,373,047	419,997	30.59%	415,947	30.29%	370,309	26.97%	166,793	12.15%	0	0.00%	0	0.00%
85+	927,747	489,610	52.77%	204,425	22.03%	86,793	9.36%	146,919	15.84%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	2,779,841	535,641	19.27%	647,641	23.30%	820,176	29.50%	776,383	27.93%	0	0.00%	0	0.00%
Other Hispanic	1,077,720	70,488	6.54%	179,514	16.66%	314,998	29.23%	512,719	47.57%	0	0.00%	0	0.00%
Non-Hispanic White	31,886,450	6,158,662	19.31%	7,080,327	22.20%	7,520,840	23.59%	11,126,622	34.89%	0	0.00%	0	0.00%
Non-Hispanic Black	3,421,700	533,486	15.59%	661,025	19.32%	994,136	29.05%	1,233,053	36.04%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	1,772,091	371,824	20.98%	251,562	14.20%	351,959	19.86%	796,746	44.96%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,258,930	244,197	19.40%	380,848	30.25%	244,110	19.39%	389,774	30.96%	0	0.00%	0	0.00%
Less Than 9th Grade	3,139,070	1,011,318	32.22%	672,487	21.42%	690,744	22.00%	764,520	24.35%	0	0.00%	0	0.00%
9-11th Grade	4,198,384	1,187,209	28.28%	1,167,003	27.80%	869,383	20.71%	974,788	23.22%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	32,294,417	5,192,124	16.08%	6,596,373	20.43%	8,189,478	25.36%	12,316,441	38.14%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	24,069,512	3,876,597	16.11%	5,082,960	21.12%	6,226,374	25.87%	8,883,580	36.91%	0	0.00%	0	0.00%
Widowed	5,501,185	1,842,132	33.49%	1,316,835	23.94%	867,498	15.77%	1,474,720	26.81%	0	0.00%	0	0.00%
Divorced	4,333,056	863,275	19.92%	999,090	23.06%	992,594	22.91%	1,478,098	34.11%	0	0.00%	0	0.00%
Separated	898,880	209,679	23.33%	278,705	31.01%	217,613	24.21%	192,883	21.46%	0	0.00%	0	0.00%
Never Married	3,964,174	602,091	15.19%	764,656	19.29%	877,390	22.13%	1,720,036	43.39%	0	0.00%	0	0.00%
Living with Partner	2,060,322	243,170	11.80%	371,700	18.04%	749,247	36.37%	696,206	33.79%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,119,563	706,644	33.34%	612,110	28.88%	388,690	18.34%	412,119	19.44%	0	0.00%	0	0.00%
Poverty income ratio less than 1	5,171,207	1,031,533	19.95%	1,024,378	19.81%	1,289,370	24.93%	1,825,926	35.31%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	33,647,032	5,931,923	17.63%	7,183,582	21.35%	8,324,049	24.74%	12,207,478	36.28%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	611,364	292,284	47.81%	248,650	40.67%	44,604	7.30%	25,825	4.22%	0	0.00%	0	0.00%
Normal: less than 25	11,818,706	1,840,819	15.58%	2,117,518	17.92%	2,983,149	25.24%	4,877,219	41.27%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	12,104,924	1,992,419	16.46%	2,606,687	21.53%	3,249,411	26.84%	4,256,408	35.16%	0	0.00%	0	0.00%
Obese: 30 or higher	16,402,809	3,544,578	21.61%	3,847,214	23.45%	3,724,945	22.71%	5,286,071	32.23%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,020,006	128,057	12.55%	411,685	40.36%	120,409	11.80%	359,855	35.28%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	30,297,213	5,200,498	17.16%	5,684,835	18.76%	7,845,352	25.89%	11,566,528	38.18%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	6,341,434	1,509,159	23.80%	1,605,889	25.32%	1,378,529	21.74%	1,847,858	29.14%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	3,279,150	832,386	25.38%	1,117,660	34.08%	657,820	20.06%	671,283	20.47%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,332,499	1,229,835	28.39%	1,187,693	27.41%	1,028,364	23.74%	886,608	20.46%	0	0.00%	0	0.00%
No	36,605,303	6,440,266	17.59%	7,632,376	20.85%	8,973,745	24.51%	13,558,916	37.04%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	19,253,688	3,914,672	20.33%	4,336,091	22.52%	4,450,836	23.12%	6,552,089	34.03%	0	0.00%	0	0.00%
No	21,679,412	3,750,726	17.30%	4,483,978	20.68%	5,551,273	25.61%	7,893,435	36.41%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	15,763,719	4,203,768	26.67%	3,762,573	23.87%	3,568,906	22.64%	4,228,473	26.82%	0	0.00%	0	0.00%
No	25,009,797	3,390,409	13.56%	5,029,098	20.11%	6,373,239	25.48%	10,217,051	40.85%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,065,226	2,087,224	34.41%	1,413,972	23.31%	1,185,608	19.55%	1,378,421	22.73%	0	0.00%	0	0.00%
Mobility limitations	6,789,352	1,777,716	26.18%	1,816,876	26.76%	1,172,432	17.27%	2,022,328	29.79%	0	0.00%	0	0.00%
No mobility limitations	9,254,130	1,611,506	17.41%	2,013,956	21.76%	2,457,344	26.55%	3,171,324	34.27%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	18,829,095	2,193,654	11.65%	3,575,264	18.99%	5,186,725	27.55%	7,873,451	41.82%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,212,097	347,554	28.67%	270,087	22.28%	267,552	22.07%	326,904	26.97%	0	0.00%	0	0.00%
Younger than 30	1,504,079	351,855	23.39%	220,866	14.68%	498,564	33.15%	432,794	28.77%	0	0.00%	0	0.00%
30 to 34	2,658,334	580,798	21.85%	769,095	28.93%	463,072	17.42%	845,368	31.80%	0	0.00%	0	0.00%
35 to 39	2,986,470	1,129,289	37.81%	586,101	19.63%	344,387	11.53%	926,693	31.03%	0	0.00%	0	0.00%
40 to 44	2,804,191	651,693	23.24%	540,235	19.27%	778,212	27.75%	834,051	29.74%	0	0.00%	0	0.00%
45 to 49	4,913,679	775,773	15.79%	1,460,852	29.73%	1,079,149	21.96%	1,597,906	32.52%	0	0.00%	0	0.00%
50 to 54	4,409,693	1,292,104	29.30%	716,640	16.25%	1,107,317	25.11%	1,293,632	29.34%	0	0.00%	0	0.00%
55 or older	2,240,211	479,170	21.39%	800,615	35.74%	535,265	23.89%	425,162	18.98%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	18,136,080	2,007,924	11.07%	3,436,549	18.95%	4,928,591	27.18%	7,763,016	42.80%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	772,313	275,749	35.70%	244,953	31.72%	97,721	12.65%	153,890	19.93%	0	0.00%	0	0.00%
None	4,779,210	649,884	13.60%	636,411	13.32%	911,477	19.07%	2,581,438	54.01%	0	0.00%	0	0.00%
1	5,270,778	693,438	13.16%	1,153,008	21.88%	1,363,979	25.88%	2,060,353	39.09%	0	0.00%	0	0.00%
2	9,127,200	1,226,815	13.44%	2,664,348	29.19%	2,127,940	23.31%	3,108,096	34.05%	0	0.00%	0	0.00%
More than 2	20,988,301	4,824,215	22.99%	4,121,349	19.64%	5,500,991	26.21%	6,541,747	31.17%	0	0.00%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	875,220	275,749	31.51%	327,184	37.38%	194,148	22.18%	78,140	8.93%	0	0.00%	0	0.00%
29 years old or younger	25,690,010	5,538,387	21.56%	5,664,650	22.05%	6,629,627	25.81%	7,857,346	30.59%	0	0.00%	0	0.00%
30 - 40 years old	1,780,259	230,504	12.95%	521,402	29.29%	301,632	16.94%	726,722	40.82%	0	0.00%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	4,779,210	649,884	13.60%	636,411	13.32%	911,477	19.07%	2,581,438	54.01%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	6,503,725	760,926	11.70%	1,477,951	22.72%	1,718,757	26.43%	2,546,091	39.15%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,246,447	206,415	16.56%	192,471	15.44%	246,468	19.77%	601,093	48.22%	0	0.00%	0	0.00%
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,708,022	554,836	20.49%	924,409	34.14%	662,956	24.48%	565,821	20.89%	0	0.00%	0	0.00%
29 years old or younger	20,155,726	3,692,954	18.32%	4,377,939	21.72%	5,085,969	25.23%	6,998,863	34.72%	0	0.00%	0	0.00%
30 - 40 years old	11,359,884	2,402,754	21.15%	2,546,052	22.41%	2,856,559	25.15%	3,554,520	31.29%	0	0.00%	0	0.00%
41 years or older	639,731	119,700	18.71%	137,564	21.50%	238,680	37.31%	143,788	22.48%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	4,779,210	649,884	13.60%	636,411	13.32%	911,477	19.07%	2,581,438	54.01%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,246,447	206,415	16.56%	192,471	15.44%	246,468	19.77%	601,093	48.22%	0	0.00%	0	0.00%
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	14,683,175	1,686,527	11.49%	2,842,771	19.36%	3,734,456	25.43%	6,419,422	43.72%	0	0.00%	0	0.00%
Yes	11,601,040	3,403,594	29.34%	2,428,396	20.93%	2,411,319	20.79%	3,357,731	28.94%	0	0.00%	0	0.00%
No	14,585,591	2,575,277	17.66%	3,548,902	24.33%	3,793,040	26.01%	4,668,371	32.01%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%)	8,820,069	21.55%)	10,002,109	24.43%)	14,445,524	35.29%)	0	0.00%)	0	0.00%)
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	14,751,171	1,691,229	11.47%	2,842,771	19.27%	3,797,749	25.75%	6,419,422	43.52%	0	0.00%	0	0.00%
35 years old or younger	4,421,643	1,232,108	27.87%	1,038,946	23.50%	857,091	19.38%	1,293,498	29.25%	0	0.00%	0	0.00%
36 - 50 years old	5,592,360	1,335,506	23.88%	1,145,225	20.48%	1,225,983	21.92%	1,885,645	33.72%	0	0.00%	0	0.00%
51 years or older	1,550,288	799,230	51.55%	244,225	15.75%	328,246	21.17%	178,588	11.52%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	14,585,591	2,575,277	17.66%	3,548,902	24.33%	3,793,040	26.01%	4,668,371	32.01%	0	0.00%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	40,937,802	7,670,100	18.74%	8,820,069	21.55%	10,002,109	24.43%	14,445,524	35.29%	0	0.00%	0	0%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	832,574	275,749	33.12%	244,953	29.42%	166,493	20.00%	145,379	17.46%	0	0.00%	0	0%
Yes	11,601,040	3,403,594	29.34%	2,428,396	20.93%	2,411,319	20.79%	3,357,731	28.94%	0	0.00%	0	0%
No	28,436,191	3,986,056	14.02%	6,146,720	21.62%	7,361,002	25.89%	10,942,413	38.48%	0	0.00%	0	0%
Don't know	*	*	*	*	*	*	*	*	*	*		*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	3,239,453	134,739	4.16%	175,584	5.42%	704,864	21.76%	2,224,267	68.66%	0	0.00%	0	0.00%
30-39	6,960,817	622,423	8.94%	793,966	11.41%	1,667,666	23.96%	3,869,576	55.59%	7,185	0.10%	0	0.00%
40-49	9,920,393	1,010,291	10.18%	1,018,380	10.27%	2,775,253	27.98%	5,116,469	51.58%	0	0.00%	0	0.00%
50-59	9,123,564	501,210	5.49%	1,264,075	13.86%	3,090,123	33.87%	4,225,464	46.31%	42,693	0.47%	0	0.00%
60-64	3,555,336	523,937	14.74%	382,741	10.77%	1,041,896	29.31%	1,606,761	45.19%	0	0.00%	0	0.00%
65-69	3,222,284	147,127	4.57%	646,394	20.06%	1,324,457	41.10%	1,104,306	34.27%	0	0.00%	0	0.00%
70-74	2,368,953	305,686	12.90%	502,665	21.22%	593,193	25.04%	967,409	40.84%	0	0.00%	0	0.00%
75-79	1,890,669	152,014	8.04%	446,477	23.61%	368,743	19.50%	871,712	46.11%	51,724	2.74%	0	0.00%
80-84	1,215,250	301,634	24.82%	172,473	14.19%	281,936	23.20%	459,208	37.79%	0	0.00%	0	0.00%
85+	989,972	220,157	22.24%	218,661	22.09%	141,039	14.25%	410,115	41.43%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	3,157,732	182,709	5.79%	440,083	13.94%	841,435	26.65%	1,667,576	52.81%	25,928	0.82%	0	0.00%
Other Hispanic	1,661,885	211,073	12.70%	68,894	4.15%	490,297	29.50%	891,621	53.65%	0	0.00%	0	0.00%
Non-Hispanic White	31,928,636	2,974,822	9.32%	4,278,494	13.40%	9,497,332	29.75%	15,126,265	47.38%	51,724	0.16%	0	0.00%
Non-Hispanic Black	3,517,078	387,169	11.01%	526,971	14.98%	700,536	19.92%	1,878,453	53.41%	23,950	0.68%	0	0.00%
Other Race - Including Multi-Racial	2,299,278	163,444	7.11%	306,974	13.35%	502,239	21.84%	1,326,621	57.70%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	831,195	104,723	12.60%	92,950	11.18%	357,532	43.01%	275,989	33.20%	0	0.00%	0	0.00%
Less Than 9th Grade	2,556,698	392,194	15.34%	393,291	15.38%	757,896	29.64%	994,574	38.90%	18,743	0.73%	0	0.00%
9-11th Grade	5,319,906	850,925	16.00%	634,549	11.93%	1,443,156	27.13%	2,339,552	43.98%	51,724	0.97%	0	0.00%
High School Grad/GED or Equivalent	33,852,812	2,571,375	7.60%	4,500,626	13.29%	9,473,255	27.98%	17,276,421	51.03%	31,134	0.09%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	25,849,680	2,132,710	8.25%	3,302,982	12.78%	7,553,640	29.22%	12,801,439	49.52%	58,908	0.23%	0	0.00%
Widowed	3,465,651	651,040	18.79%	869,631	25.09%	708,260	20.44%	1,236,721	35.69%	0	0.00%	0	0.00%
Divorced	5,705,638	454,632	7.97%	654,615	11.47%	1,862,524	32.64%	2,709,918	47.50%	23,950	0.42%	0	0.00%
Separated	1,366,068	223,278	16.34%	74,192	5.43%	522,158	38.22%	527,697	38.63%	18,743	1.37%	0	0.00%
Never Married	3,591,518	99,814	2.78%	540,147	15.04%	863,891	24.05%	2,087,666	58.13%	0	0.00%	0	0.00%
Living with Partner	2,565,446	357,744	13.94%	177,558	6.92%	519,978	20.27%	1,510,167	58.87%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,425,556	189,413	13.29%	133,328	9.35%	562,378	39.45%	540,437	37.91%	0	0.00%	0	0.00%
Poverty income ratio less than 1	4,166,444	499,885	12.00%	873,113	20.96%	1,333,394	32.00%	1,434,125	34.42%	25,928	0.62%	0	0.00%
Poverty income ratio 1 or greater	36,972,609	3,229,919	8.74%	4,614,975	12.48%	10,136,068	27.42%	18,915,973	51.16%	75,673	0.20%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	357,091	96,677	27.07%	52,148	14.60%	201,324	56.38%	6,942	1.94%	0	0.00%	0	0.00%
Normal: less than 25	13,169,925	592,880	4.50%	1,485,483	11.28%	3,499,382	26.57%	7,540,455	57.26%	51,724	0.39%	0	0.00%
Overweight: 25.0 - 29.9	10,934,070	1,226,913	11.22%	1,718,025	15.71%	3,188,399	29.16%	4,800,733	43.91%	0	0.00%	0	0.00%
Obese: 30 or higher	18,103,524	2,002,747	11.06%	2,365,760	13.07%	5,142,735	28.41%	8,542,405	47.19%	49,877	0.28%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	731,777	16,245	2.22%	129,943	17.76%	146,950	20.08%	414,689	56.67%	23,950	3.27%	0	0.00%
Glycohemoglobin < 5.7%	30,755,342	2,421,366	7.87%	3,654,594	11.88%	9,015,575	29.31%	15,663,806	50.93%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	7,742,768	1,106,361	14.29%	1,322,049	17.07%	2,147,935	27.74%	3,088,771	39.89%	77,651	1.00%	0	0.00%
Glycohemoglobin 6.5% or higher	3,334,722	375,244	11.25%	514,829	15.44%	721,379	21.63%	1,723,269	51.68%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,725,468	743,928	15.74%	581,930	12.31%	1,087,159	23.01%	2,312,451	48.94%	0	0.00%	0	0.00%
No	37,839,141	3,175,289	8.39%	5,039,485	13.32%	10,944,681	28.92%	18,578,084	49.10%	101,601	0.27%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	18,903,740	1,700,166	8.99%	2,707,437	14.32%	5,225,173	27.64%	9,239,831	48.88%	31,134	0.16%	0	0.00%
No	23,652,382	2,219,051	9.38%	2,913,979	12.32%	6,798,180	28.74%	11,650,705	49.26%	70,467	0.30%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	16,330,392	1,933,864	11.84%	2,477,640	15.17%	4,753,800	29.11%	7,141,139	43.73%	23,950	0.15%	0	0.00%
No	26,047,905	1,985,353	7.62%	3,143,776	12.07%	7,246,306	27.82%	13,594,819	52.19%	77,651	0.30%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,163,308	763,502	12.39%	998,744	16.20%	2,094,147	33.98%	2,306,915	37.43%	0	0.00%	0	0.00%
Mobility limitations	5,172,690	1,125,673	21.76%	689,899	13.34%	1,321,177	25.54%	2,017,197	39.00%	18,743	0.36%	0	0.00%
No mobility limitations	9,430,103	827,650	8.78%	1,598,262	16.95%	2,823,604	29.94%	4,128,863	43.78%	51,724	0.55%	0	0.00%
N/A - Respondent not asked mobility limitation questions	21,798,508	1,202,392	5.52%	2,334,510	10.71%	5,792,911	26.57%	12,437,560	57.06%	31,134	0.14%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,060,439	365,992	9.01%	494,772	12.19%	1,426,104	35.12%	1,773,571	43.68%	0	0.00%	0	0.00%
No	38,491,555	3,540,611	9.20%	5,126,644	13.32%	10,605,736	27.55%	19,116,964	49.67%	101,601	0.26%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	481,077	45,470	9.45%	89,360	18.57%	144,346	30.00%	201,901	41.97%	0	0.00%	0	0.00%
2 or more times a day	5,468,718	538,644	9.85%	843,843	15.43%	1,577,786	28.85%	2,508,446	45.87%	0	0.00%	0	0.00%
Once a day	3,871,921	151,382	3.91%	440,713	11.38%	1,326,051	34.25%	1,929,826	49.84%	23,950	0.62%	0	0.00%
2 or more time a week	4,939,864	463,147	9.38%	933,910	18.91%	1,615,962	32.71%	1,908,100	38.63%	18,743	0.38%	0	0.00%
Once a week	3,002,113	232,843	7.76%	358,601	11.94%	938,495	31.26%	1,472,174	49.04%	0	0.00%	0	0.00%
1-3 times a month	6,764,106	534,190	7.90%	796,981	11.78%	1,657,266	24.50%	3,775,668	55.82%	0	0.00%	0	0.00%
Never	18,036,809	1,953,540	10.83%	2,158,008	11.96%	4,771,933	26.46%	9,094,421	50.42%	58,908	0.33%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	481,077	45,470	9.45%	89,360	18.57%	144,346	30.00%	201,901	41.97%	0	0.00%	0	0.00%
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	684,556	79,928	11.68%	85,545	12.50%	131,395	19.19%	387,688	56.63%	0	0.00%	0	0.00%
Never	40,949,820	3,579,435	8.74%	5,401,420	13.19%	11,658,276	28.47%	20,209,088	49.35%	101,601	0.25%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,205,729	313,321	25.99%	180,870	15.00%	260,707	21.62%	450,831	37.39%	0	0.00%	0	0.00%
30 to 34	1,734,615	323,783	18.67%	145,744	8.40%	637,138	36.73%	627,950	36.20%	0	0.00%	0	0.00%
35 to 39	2,404,562	418,955	17.42%	318,840	13.26%	962,814	40.04%	703,953	29.28%	0	0.00%	0	0.00%
40 to 44	3,761,116	475,108	12.63%	837,957	22.28%	683,245	18.17%	1,757,620	46.73%	7,185	0.19%	0	0.00%
45 to 49	4,930,199	466,576	9.46%	1,093,571	22.18%	1,591,319	32.28%	1,703,060	34.54%	75,673	1.53%	0	0.00%
50 to 54	6,302,197	365,015	5.79%	440,616	6.99%	1,926,036	30.56%	3,570,529	56.66%	0	0.00%	0	0.00%
55 or older	2,066,011	189,254	9.16%	517,021	25.03%	606,689	29.37%	753,047	36.45%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	19,813,511	1,335,199	6.74%	2,068,342	10.44%	5,250,509	26.50%	11,159,460	56.32%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
None	3,530,858	58,078	1.64%	537,219	15.21%	783,302	22.18%	2,152,259	60.96%	0	0.00%	0	0.00%
1	4,765,152	284,215	5.96%	384,244	8.06%	1,428,621	29.98%	2,668,073	55.99%	0	0.00%	0	0.00%
2	9,163,179	1,053,495	11.50%	1,066,759	11.64%	2,067,301	22.56%	4,932,930	53.83%	42,693	0.47%	0	0.00%
More than 2	25,105,420	2,523,430	10.05%	3,633,194	14.47%	7,752,615	30.88%	11,137,273	44.36%	58,908	0.23%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	217,115	0	0.00%	16,970	7.82%	101,831	46.90%	98,315	45.28%	0	0.00%	0	0.00%
None	9,387,622	363,236	3.87%	982,471	10.47%	2,313,415	24.64%	5,728,501	61.02%	0	0.00%	0	0.00%
1	5,851,319	516,638	8.83%	644,668	11.02%	1,349,797	23.07%	3,340,215	57.08%	0	0.00%	0	0.00%
2	11,250,354	1,323,025	11.76%	1,085,046	9.64%	3,540,268	31.47%	5,259,321	46.75%	42,693	0.38%	0	0.00%
More than 2	15,858,199	1,716,318	10.82%	2,892,261	18.24%	4,726,529	29.80%	6,464,184	40.76%	58,908	0.37%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	217,115	0	0.00%	16,970	7.82%	101,831	46.90%	98,315	45.28%	0	0.00%	0	0.00%
None	14,058,919	1,230,129	8.75%	1,947,511	13.85%	4,541,191	32.30%	6,340,088	45.10%	0	0.00%	0	0.00%
1	3,944,506	496,768	12.59%	184,504	4.68%	1,048,484	26.58%	2,214,749	56.15%	0	0.00%	0	0.00%
2	2,355,885	136,135	5.78%	195,096	8.28%	507,562	21.54%	1,517,092	64.40%	0	0.00%	0	0.00%
More than 2	708,758	96,071	13.55%	92,691	13.08%	103,849	14.65%	416,148	58.72%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	21,171,079	1,960,114	9.26%	3,184,644	15.04%	5,620,576	26.55%	10,304,144	48.67%	101,601	0.48%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,213,152	64,142	5.29%	87,256	7.19%	335,090	27.62%	726,664	59.90%	0	0.00%	0	0.00%
29 years old or younger	28,478,414	3,264,695	11.46%	4,074,828	14.31%	8,140,635	28.59%	12,896,656	45.29%	101,601	0.36%	0	0.00%
30 - 40 years old	1,797,224	0	0.00%	147,229	8.19%	795,514	44.26%	854,482	47.54%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	3,530,858	58,078	1.64%	537,219	15.21%	783,302	22.18%	2,152,259	60.96%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	6,285,223	512,066	8.15%	601,665	9.57%	1,745,066	27.76%	3,426,426	54.52%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,149,846	7,623	0.66%	130,181	11.32%	232,233	20.20%	779,809	67.82%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,213,152	64,142	5.29%	87,256	7.19%	335,090	27.62%	726,664	59.90%	0	0.00%	0	0.00%
29 years old or younger	19,293,999	2,349,183	12.18%	2,041,940	10.58%	5,646,390	29.27%	9,213,793	47.75%	42,693	0.22%	0	0.00%
30 - 40 years old	16,382,986	1,315,588	8.03%	2,656,224	16.21%	4,826,968	29.46%	7,525,299	45.93%	58,908	0.36%	0	0.00%
41 years or older	775,489	79,984	10.31%	152,766	19.70%	134,922	17.40%	407,817	52.59%	0	0.00%	0	0.00%
Refused/Don't know	218,278	44,620	20.44%	15,830	7.25%	72,935	33.41%	84,893	38.89%	0	0.00%	0	0.00%
N/A - Never pregnant	3,530,858	58,078	1.64%	537,219	15.21%	783,302	22.18%	2,152,259	60.96%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,149,846	7,623	0.66%	130,181	11.32%	232,233	20.20%	779,809	67.82%	0	0.00%	0	0.00%
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	15,720,625	1,052,136	6.69%	1,878,606	11.95%	3,915,751	24.91%	8,874,132	56.45%	0	0.00%	0	0.00%
Yes	10,838,859	1,812,535	16.72%	1,723,174	15.90%	3,473,055	32.04%	3,830,095	35.34%	0	0.00%	0	0.00%
No	15,971,756	1,054,545	6.60%	1,986,267	12.44%	4,643,034	29.07%	8,186,308	51.25%	101,601	0.64%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	15,753,994	1,052,136	6.68%	1,911,975	12.14%	3,915,751	24.86%	8,874,132	56.33%	0	0.00%	0	0.00%
35 years old or younger	3,055,791	657,778	21.53%	217,595	7.12%	1,119,693	36.64%	1,060,724	34.71%	0	0.00%	0	0.00%
36 - 50 years old	6,133,724	910,286	14.84%	1,326,642	21.63%	1,798,448	29.32%	2,098,348	34.21%	0	0.00%	0	0.00%
51 years or older	1,453,457	101,836	7.01%	135,899	9.35%	544,699	37.48%	671,023	46.17%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	15,971,756	1,054,545	6.60%	1,986,267	12.44%	4,643,034	29.07%	8,186,308	51.25%	101,601	0.64%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	42,564,609	3,919,217	9.21%	5,621,416	13.21%	12,031,840	28.27%	20,890,535	49.08%	101,601	0.24%	0	0.00%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	10,838,859	1,812,535	16.72%	1,723,174	15.90%	3,473,055	32.04%	3,830,095	35.34%	0	0.00%	0	0.00%
No	31,692,381	2,106,682	6.65%	3,864,874	12.19%	8,558,785	27.01%	17,060,440	53.83%	101,601	0.32%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	2,389,987	125,268	5.24%	148,956	6.23%	493,902	20.67%	1,621,862	67.86%	0	0.00%	0	0.00%
30-39	5,874,771	473,759	8.06%	462,771	7.88%	1,784,768	30.38%	3,122,998	53.16%	30,476	0.52%	0	0.00%
40-49	10,542,388	520,304	4.94%	977,507	9.27%	2,777,878	26.35%	6,266,699	59.44%	0	0.00%	0	0.00%
50-59	10,190,538	636,714	6.25%	1,381,270	13.55%	3,709,272	36.40%	4,427,530	43.45%	35,753	0.35%	0	0.00%
60-64	3,682,691	176,864	4.80%	484,841	13.17%	1,200,065	32.59%	1,820,922	49.45%	0	0.00%	0	0.00%
65-69	3,515,083	107,931	3.07%	496,868	14.14%	888,625	25.28%	2,021,658	57.51%	0	0.00%	0	0.00%
70-74	2,808,468	215,892	7.69%	410,680	14.62%	964,492	34.34%	1,146,172	40.81%	71,231	2.54%	0	0.00%
75-79	1,945,073	314,798	16.18%	131,858	6.78%	568,244	29.21%	930,173	47.82%	0	0.00%	0	0.00%
80+	2,501,274	522,504	20.89%	131,240	5.25%	961,684	38.45%	859,346	34.36%	26,498	1.06%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	3,535,216	252,254	7.14%	394,749	11.17%	1,424,505	40.29%	1,463,708	41.40%	0	0.00%	0	0.00%
Other Hispanic	1,835,340	188,640	10.28%	304,836	16.61%	525,138	28.61%	816,726	44.50%	0	0.00%	0	0.00%
Non-Hispanic White	32,645,762	2,395,535	7.34%	3,241,157	9.93%	9,658,181	29.58%	17,213,429	52.73%	137,460	0.42%	0	0.00%
Non-Hispanic Black	3,863,130	194,317	5.03%	328,607	8.51%	1,318,781	34.14%	1,994,926	51.64%	26,498	0.69%	0	0.00%
Other Race - Including Multi-Racial	1,694,691	63,289	3.73%	372,613	21.99%	458,373	27.05%	800,416	47.23%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	950,237	54,347	5.72%	130,352	13.72%	213,296	22.45%	552,242	58.12%	0	0.00%	0	0.00%
Less Than 9th Grade	3,263,040	389,389	11.93%	569,632	17.46%	985,660	30.21%	1,318,359	40.40%	0	0.00%	0	0.00%
9-11th Grade	5,919,271	643,987	10.88%	847,339	14.31%	1,899,174	32.08%	2,478,698	41.88%	50,073	0.85%	0	0.00%
High School Grad/GED or Equivalent	33,429,975	2,006,312	6.00%	3,094,640	9.26%	10,275,233	30.74%	17,939,905	53.66%	113,886	0.34%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	314,106	87,623	27.90%	25,623	8.16%	47,454	15.11%	153,407	48.84%	0	0.00%	0	0.00%
Married	26,289,858	1,430,508	5.44%	2,896,062	11.02%	7,944,198	30.22%	13,947,859	53.05%	71,231	0.27%	0	0.00%
Widowed	4,751,627	607,393	12.78%	358,508	7.54%	1,424,855	29.99%	2,348,692	49.43%	12,178	0.26%	0	0.00%
Divorced	5,862,932	667,088	11.38%	618,716	10.55%	1,736,433	29.62%	2,790,622	47.60%	50,073	0.85%	0	0.00%
Separated	1,146,885	63,410	5.53%	310,057	27.03%	378,488	33.00%	394,930	34.44%	0	0.00%	0	0.00%
Never Married	3,193,357	173,279	5.43%	321,176	10.06%	1,201,704	37.63%	1,466,722	45.93%	30,476	0.95%	0	0.00%
Living with Partner	1,988,990	64,733	3.25%	111,822	5.62%	651,847	32.77%	1,160,588	58.35%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,666,791	397,280	14.90%	523,757	19.64%	610,188	22.88%	1,135,566	42.58%	0	0.00%	0	0.00%
Poverty income ratio less than 1	5,577,123	574,951	10.31%	877,418	15.73%	1,753,553	31.44%	2,359,022	42.30%	12,178	0.22%	0	0.00%
Poverty income ratio 1 or greater	35,330,225	2,121,803	6.01%	3,240,788	9.17%	11,021,237	31.19%	18,794,618	53.20%	151,780	0.43%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	537,545	127,231	23.67%	23,857	4.44%	201,864	37.55%	184,593	34.34%	0	0.00%	0	0.00%
Normal: less than 25	12,125,743	790,732	6.52%	1,262,750	10.41%	3,313,311	27.32%	6,711,018	55.35%	47,931	0.40%	0	0.00%
Overweight: 25.0 - 29.9	12,310,041	857,316	6.96%	1,195,520	9.71%	4,145,785	33.68%	6,111,420	49.65%	0	0.00%	0	0.00%
Obese: 30 or higher	18,600,809	1,318,754	7.09%	2,159,836	11.61%	5,724,018	30.77%	9,282,174	49.90%	116,027	0.62%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,150,933	13,028	1.13%	121,924	10.59%	387,989	33.71%	613,674	53.32%	14,320	1.24%	0	0.00%
Glycohemoglobin < 5.7%	27,395,711	1,405,254	5.13%	2,613,135	9.54%	8,567,986	31.27%	14,809,335	54.06%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	11,037,175	1,142,267	10.35%	1,081,933	9.80%	3,641,364	32.99%	5,123,680	46.42%	47,931	0.43%	0	0.00%
Glycohemoglobin 6.5% or higher	3,990,320	533,485	13.37%	824,972	20.67%	787,639	19.74%	1,742,516	43.67%	101,707	2.55%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,550,032	859,160	15.48%	783,258	14.11%	1,241,408	22.37%	2,564,500	46.21%	101,707	1.83%	0	0.00%
No	38,024,106	2,234,874	5.88%	3,858,705	10.15%	12,143,571	31.94%	19,724,705	51.87%	62,251	0.16%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	19,612,973	1,600,517	8.16%	2,008,808	10.24%	5,940,036	30.29%	9,913,974	50.55%	149,638	0.76%	0	0.00%
No	23,961,165	1,493,517	6.23%	2,633,155	10.99%	7,444,943	31.07%	12,375,231	51.65%	14,320	0.06%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	17,323,777	1,801,096	10.40%	2,028,254	11.71%	5,308,945	30.65%	8,057,276	46.51%	128,205	0.74%	0	0.00%
No	26,250,362	1,292,938	4.93%	2,613,709	9.96%	8,076,033	30.77%	14,231,928	54.22%	35,753	0.14%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,171,749	788,285	12.77%	600,824	9.74%	1,440,738	23.34%	3,311,427	53.65%	30,476	0.49%	0	0.00%
Mobility limitations	5,628,671	484,580	8.61%	825,255	14.66%	1,929,076	34.27%	2,292,032	40.72%	97,729	1.74%	0	0.00%
No mobility limitations	9,028,727	757,738	8.39%	1,050,678	11.64%	2,954,107	32.72%	4,230,451	46.86%	35,753	0.40%	0	0.00%
N/A - Respondent not asked mobility limitation questions	22,744,991	1,063,431	4.68%	2,165,206	9.52%	7,061,058	31.04%	12,455,295	54.76%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,720,848	422,767	11.36%	800,800	21.52%	779,105	20.94%	1,718,176	46.18%	0	0.00%	0	0.00%
No	39,763,911	2,671,267	6.72%	3,841,163	9.66%	12,532,373	31.52%	20,555,150	51.69%	163,958	0.41%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	324,720	58,399	17.98%	100,658	31.00%	47,666	14.68%	87,522	26.95%	30,476	9.39%	0	0.00%
2 or more times a day	5,231,083	585,545	11.19%	764,482	14.61%	1,782,365	34.07%	2,098,690	40.12%	0	0.00%	0	0.00%
Once a day	3,913,332	295,578	7.55%	540,370	13.81%	1,143,144	29.21%	1,934,241	49.43%	0	0.00%	0	0.00%
2 or more time a week	3,844,144	188,309	4.90%	807,580	21.01%	1,391,847	36.21%	1,456,407	37.89%	0	0.00%	0	0.00%
Once a week	3,608,143	122,562	3.40%	249,222	6.91%	958,466	26.56%	2,277,893	63.13%	0	0.00%	0	0.00%
1-3 times a month	7,135,871	764,410	10.71%	639,544	8.96%	2,118,993	29.69%	3,541,695	49.63%	71,231	1.00%	0	0.00%
Never	19,466,051	1,054,935	5.42%	1,540,107	7.91%	5,942,498	30.53%	10,892,758	55.96%	35,753	0.18%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	324,720	58,399	17.98%	100,658	31.00%	47,666	14.68%	87,522	26.95%	30,476	9.39%	0	0.00%
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	1,075,204	252,949	23.53%	47,983	4.46%	220,551	20.51%	482,491	44.87%	71,231	6.62%	0	0.00%
Never	41,498,405	2,570,364	6.19%	4,344,094	10.47%	12,942,880	31.19%	21,628,888	52.12%	12,178	0.03%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,125,672	193,715	17.21%	126,305	11.22%	282,808	25.12%	451,613	40.12%	71,231	6.33%	0	0.00%
30 to 34	1,246,661	37,183	2.98%	201,899	16.20%	431,173	34.59%	576,406	46.24%	0	0.00%	0	0.00%
35 to 39	2,595,857	99,738	3.84%	256,763	9.89%	882,255	33.99%	1,357,101	52.28%	0	0.00%	0	0.00%
40 to 44	4,075,874	394,993	9.69%	565,613	13.88%	1,537,653	37.73%	1,527,543	37.48%	50,073	1.23%	0	0.00%
45 to 49	5,902,598	663,974	11.25%	584,894	9.91%	2,134,517	36.16%	2,519,212	42.68%	0	0.00%	0	0.00%
50 to 54	5,183,409	542,507	10.47%	637,384	12.30%	1,346,924	25.99%	2,644,416	51.02%	12,178	0.23%	0	0.00%
55 or older	2,615,254	167,202	6.39%	232,228	8.88%	800,306	30.60%	1,385,042	52.96%	30,476	1.17%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	20,626,000	980,927	4.76%	1,963,772	9.52%	5,961,907	28.90%	11,719,395	56.82%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
None	4,300,728	283,651	6.60%	469,913	10.93%	996,208	23.16%	2,550,956	59.31%	0	0.00%	0	0.00%
1	4,324,010	173,569	4.01%	492,034	11.38%	1,376,372	31.83%	2,246,282	51.95%	35,753	0.83%	0	0.00%
2	10,327,253	452,700	4.38%	831,947	8.06%	3,441,665	33.33%	5,570,465	53.94%	30,476	0.30%	0	0.00%
More than 2	24,622,147	2,184,113	8.87%	2,848,069	11.57%	7,570,734	30.75%	11,921,502	48.42%	97,729	0.40%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	9,434,212	571,062	6.05%	810,274	8.59%	2,639,278	27.98%	5,276,138	55.93%	137,460	1.46%	0	0.00%
1	7,482,869	439,063	5.87%	894,238	11.95%	2,401,697	32.10%	3,747,872	50.09%	0	0.00%	0	0.00%
2	11,944,848	528,438	4.42%	1,259,354	10.54%	4,080,984	34.17%	6,076,071	50.87%	0	0.00%	0	0.00%
More than 2	14,466,230	1,555,471	10.75%	1,629,627	11.27%	4,263,020	29.47%	7,005,933	48.43%	12,178	0.08%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	12,459,932	1,028,177	8.25%	1,434,325	11.51%	4,483,585	35.98%	5,513,846	44.25%	0	0.00%	0	0.00%
1	4,166,993	328,230	7.88%	368,482	8.84%	1,237,708	29.70%	2,196,820	52.72%	35,753	0.86%	0	0.00%
2	2,055,989	80,866	3.93%	117,127	5.70%	729,910	35.50%	1,026,377	49.92%	101,707	4.95%	0	0.00%
More than 2	1,399,863	223,129	15.94%	248,363	17.74%	230,432	16.46%	697,940	49.86%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	23,174,357	1,433,632	6.19%	2,425,196	10.46%	6,703,343	28.93%	12,600,007	54.37%	12,178	0.05%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,902,879	292,762	4.96%	571,537	9.68%	1,719,898	29.14%	3,318,682	56.22%	0	0.00%	0	0.00%
Yes	6,751,371	454,861	6.74%	869,849	12.88%	1,605,049	23.77%	3,821,612	56.60%	0	0.00%	0	0.00%
No	30,875,242	2,346,411	7.60%	3,200,577	10.37%	10,037,855	32.51%	15,126,441	48.99%	163,958	0.53%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,947,525	292,762	4.92%	571,537	9.61%	1,742,074	29.29%	3,341,152	56.18%	0	0.00%	0	0.00%
35 years old or younger	6,233,281	447,874	7.19%	862,989	13.84%	1,528,757	24.53%	3,393,660	54.44%	0	0.00%	0	0.00%
36 years or older	518,090	6,986	1.35%	6,860	1.32%	76,292	14.73%	427,952	82.60%	0	0.00%	0	0.00%
N/A - Did not have baby weighing 9 lbs or more	30,875,242	2,346,411	7.60%	3,200,577	10.37%	10,037,855	32.51%	15,126,441	48.99%	163,958	0.53%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,602,152	9,111	0.57%	101,624	6.34%	723,691	45.17%	767,726	47.92%	0	0.00%	0	0.00%
29 years old or younger	27,873,639	2,269,621	8.14%	2,820,481	10.12%	8,734,488	31.34%	13,920,844	49.94%	128,205	0.46%	0	0.00%
30 - 40 years old	2,875,571	41,266	1.44%	450,972	15.68%	764,273	26.58%	1,619,060	56.30%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	4,300,728	283,651	6.60%	469,913	10.93%	996,208	23.16%	2,550,956	59.31%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	6,822,202	476,590	6.99%	798,973	11.71%	2,142,567	31.41%	3,368,320	49.37%	35,753	0.52%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,121,717	9,111	0.43%	108,484	5.11%	908,498	42.82%	1,095,624	51.64%	0	0.00%	0	0.00%
29 years old or younger	21,522,115	1,557,617	7.24%	1,853,091	8.61%	6,991,311	32.48%	11,018,389	51.20%	101,707	0.47%	0	0.00%
30 - 40 years old	14,499,324	1,129,071	7.79%	2,107,136	14.53%	4,185,512	28.87%	7,029,674	48.48%	47,931	0.33%	0	0.00%
41 years or older	885,016	93,643	10.58%	75,790	8.56%	234,067	26.45%	481,516	54.41%	0	0.00%	0	0.00%
Refused/Don't know	191,646	20,940	10.93%	27,549	14.37%	45,631	23.81%	83,206	43.42%	14,320	7.47%	0	0.00%
N/A - Never pregnant	4,300,728	283,651	6.60%	469,913	10.93%	996,208	23.16%	2,550,956	59.31%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,219,082	1,282,122	11.43%	1,131,031	10.08%	4,077,738	36.35%	4,642,641	41.38%	85,551	0.76%	0	0.00%
No	32,317,563	1,811,912	5.61%	3,510,932	10.86%	9,307,241	28.80%	17,609,070	54.49%	78,407	0.24%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
35 years old or younger	2,525,405	277,975	11.01%	279,032	11.05%	954,519	37.80%	942,648	37.33%	71,231	2.82%	0	0.00%
36 - 50 years old	7,127,515	744,075	10.44%	703,254	9.87%	2,276,579	31.94%	3,403,607	47.75%	0	0.00%	0	0.00%
51 years or older	1,364,112	222,542	16.31%	148,746	10.90%	714,675	52.39%	278,150	20.39%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	32,317,563	1,811,912	5.61%	3,510,932	10.86%	9,307,241	28.80%	17,609,070	54.49%	78,407	0.24%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	772,445	0	0.00%	72,729	9.42%	226,988	29.39%	472,727	61.20%	0	0.00%	0	0.00%
Yes	6,713,473	919,312	13.69%	666,467	9.93%	2,388,262	35.57%	2,653,881	39.53%	85,551	1.27%	0	0.00%
No	35,893,923	2,174,722	6.06%	3,882,537	10.82%	10,681,840	29.76%	19,076,417	53.15%	78,407	0.22%	0	0.00%
Don't know	194,297	0	0.00%	20,230	10.41%	87,887	45.23%	86,179	44.35%	0	0.00%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	966,742	0	0.00%	92,959	9.62%	314,876	32.57%	558,907	57.81%	0	0.00%	0	0.00%
35 years old or younger	864,424	144,134	16.67%	69,283	8.01%	268,240	31.03%	311,536	36.04%	71,231	8.24%	0	0.00%
36 - 50 years old	4,375,715	491,100	11.22%	461,957	10.56%	1,306,448	29.86%	2,116,211	48.36%	0	0.00%	0	0.00%
51 years or older	1,370,647	284,078	20.73%	135,227	9.87%	725,208	52.91%	226,134	16.50%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	35,893,923	2,174,722	6.06%	3,882,537	10.82%	10,681,840	29.76%	19,076,417	53.15%	78,407	0.22%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	43,574,138	3,094,034	7.10%	4,641,963	10.65%	13,384,978	30.72%	22,289,205	51.15%	163,958	0.38%	0	0.00%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,058,207	1,247,485	11.28%	1,114,673	10.08%	4,060,756	36.72%	4,549,743	41.14%	85,551	0.77%	0	0.00%
No	32,478,438	1,846,549	5.69%	3,527,290	10.86%	9,324,223	28.71%	17,701,969	54.50%	78,407	0.24%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	3,150,713	149,777	4.75%	141,435	4.49%	465,523	14.78%	2,393,978	75.98%	0	0.00%	0	0.00%
30-39	5,460,004	166,973	3.06%	714,152	13.08%	1,637,715	29.99%	2,941,165	53.87%	0	0.00%	0	0.00%
40-49	8,461,428	848,553	10.03%	1,163,774	13.75%	2,643,585	31.24%	3,805,516	44.97%	0	0.00%	0	0.00%
50-59	10,595,004	673,643	6.36%	1,426,700	13.47%	2,732,932	25.79%	5,736,303	54.14%	0	0.00%	25,427	0.24%
60-64	3,593,050	148,356	4.13%	744,807	20.73%	760,829	21.18%	1,939,057	53.97%	0	0.00%	0	0.00%
65-69	2,846,674	370,253	13.01%	206,180	7.24%	1,017,368	35.74%	1,252,873	44.01%	0	0.00%	0	0.00%
70-74	2,783,570	337,224	12.11%	382,243	13.73%	887,222	31.87%	1,176,881	42.28%	0	0.00%	0	0.00%
75-79	2,155,087	414,617	19.24%	156,206	7.25%	926,091	42.97%	658,173	30.54%	0	0.00%	0	0.00%
80+	2,320,241	453,299	19.54%	233,679	10.07%	845,298	36.43%	787,965	33.96%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	3,258,435	312,385	9.59%	422,447	12.96%	1,058,111	32.47%	1,465,492	44.98%	0	0.00%	0	0.00%
Other Hispanic	2,286,436	119,575	5.23%	298,881	13.07%	636,847	27.85%	1,231,133	53.85%	0	0.00%	0	0.00%
Non-Hispanic White	30,321,787	2,789,670	9.20%	4,082,795	13.46%	8,847,720	29.18%	14,576,175	48.07%	0	0.00%	25,427	0.08%
Non-Hispanic Black	3,376,104	204,149	6.05%	204,566	6.06%	964,244	28.56%	2,003,144	59.33%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	2,322,409	156,734	6.75%	160,487	6.91%	485,824	20.92%	1,519,364	65.42%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,124,284	251,411	22.36%	260,754	23.19%	24,088	2.14%	588,032	52.30%	0	0.00%	0	0.00%
Less Than 9th Grade	2,766,828	328,567	11.88%	289,304	10.46%	1,117,409	40.39%	1,031,548	37.28%	0	0.00%	0	0.00%
9-11th Grade	5,087,699	603,947	11.87%	817,396	16.07%	1,724,552	33.90%	1,941,804	38.17%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	32,476,919	2,392,565	7.37%	3,773,418	11.62%	9,126,697	28.10%	17,158,811	52.83%	0	0.00%	25,427	0.08%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	24,874,488	2,003,673	8.06%	3,065,136	12.32%	6,832,698	27.47%	12,947,554	52.05%	0	0.00%	25,427	0.10%
Widowed	4,090,678	771,050	18.85%	720,442	17.61%	1,472,500	36.00%	1,126,686	27.54%	0	0.00%	0	0.00%
Divorced	4,648,101	391,408	8.42%	404,739	8.71%	1,295,888	27.88%	2,556,066	54.99%	0	0.00%	0	0.00%
Separated	812,753	34,958	4.30%	141,458	17.40%	311,163	38.29%	325,175	40.01%	0	0.00%	0	0.00%
Never Married	4,118,534	293,733	7.13%	400,610	9.73%	1,065,009	25.86%	2,359,182	57.28%	0	0.00%	0	0.00%
Living with Partner	2,657,020	67,873	2.55%	311,474	11.72%	939,306	35.35%	1,338,367	50.37%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,647,636	238,350	9.00%	369,639	13.96%	800,569	30.24%	1,239,078	46.80%	0	0.00%	0	0.00%
Poverty income ratio less than 1	5,923,419	570,237	9.63%	852,178	14.39%	1,942,859	32.80%	2,558,146	43.19%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	32,994,116	2,773,928	8.41%	3,947,358	11.96%	9,249,318	28.03%	16,998,084	51.52%	0	0.00%	25,427	0.08%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	11,270,152	775,811	6.88%	898,112	7.97%	2,901,330	25.74%	6,694,899	59.40%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	11,673,193	1,181,504	10.12%	1,313,270	11.25%	3,994,511	34.22%	5,183,908	44.41%	0	0.00%	0	0.00%
Obese: 30 or higher	18,517,821	1,625,199	8.78%	2,921,057	15.77%	5,029,637	27.16%	8,916,501	48.15%	0	0.00%	25,427	0.14%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,376,114	41,882	3.04%	120,038	8.72%	442,818	32.18%	771,375	56.05%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	23,931,482	2,046,823	8.55%	2,638,715	11.03%	7,059,327	29.50%	12,161,189	50.82%	0	0.00%	25,427	0.11%
Glycohemoglobin 5.7% - 6.4%	12,584,363	1,065,854	8.47%	1,644,962	13.07%	3,418,427	27.16%	6,455,119	51.29%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	3,673,212	427,954	11.65%	765,460	20.84%	1,072,173	29.19%	1,407,624	38.32%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,941,122	652,454	13.20%	810,405	16.40%	1,470,176	29.75%	2,008,088	40.64%	0	0.00%	0	0.00%
No	36,624,049	2,930,060	8.00%	4,358,771	11.90%	10,522,570	28.73%	18,787,220	51.30%	0	0.00%	25,427	0.07%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	17,189,244	1,718,072	10.00%	2,841,512	16.53%	4,875,498	28.36%	7,728,734	44.96%	0	0.00%	25,427	0.15%
No	24,375,928	1,864,442	7.65%	2,327,664	9.55%	7,117,248	29.20%	13,066,574	53.60%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	15,658,321	1,788,266	11.42%	2,328,121	14.87%	4,664,726	29.79%	6,851,780	43.76%	0	0.00%	25,427	0.16%
No	25,889,369	1,794,248	6.93%	2,841,055	10.97%	7,310,538	28.24%	13,943,528	53.86%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,512,998	871,699	15.81%	836,764	15.18%	1,623,778	29.45%	2,180,756	39.56%	0	0.00%	0	0.00%
Mobility limitations	5,546,085	747,245	13.47%	722,231	13.02%	1,940,255	34.98%	2,110,927	38.06%	0	0.00%	25,427	0.46%
No mobility limitations	9,189,061	873,813	9.51%	1,083,807	11.79%	2,782,704	30.28%	4,448,737	48.41%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	21,317,028	1,089,756	5.11%	2,526,374	11.85%	5,646,009	26.49%	12,054,888	56.55%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,034,403	606,266	15.03%	588,523	14.59%	1,338,743	33.18%	1,500,870	37.20%	0	0.00%	0	0.00%
No	37,461,743	2,976,248	7.94%	4,511,626	12.04%	10,654,004	28.44%	19,294,438	51.50%	0	0.00%	25,427	0.07%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	4,953,414	511,268	10.32%	777,503	15.70%	1,685,492	34.03%	1,979,151	39.96%	0	0.00%	0	0.00%
Once a day	3,644,732	382,045	10.48%	481,700	13.22%	734,410	20.15%	2,046,576	56.15%	0	0.00%	0	0.00%
2 or more time a week	4,467,470	569,206	12.74%	742,168	16.61%	1,302,046	29.15%	1,854,050	41.50%	0	0.00%	0	0.00%
Once a week	2,649,682	134,927	5.09%	383,883	14.49%	936,551	35.35%	1,194,322	45.07%	0	0.00%	0	0.00%
1-3 times a month	6,040,827	435,385	7.21%	549,064	9.09%	1,891,031	31.30%	3,165,346	52.40%	0	0.00%	0	0.00%
Never	19,557,241	1,472,765	7.53%	2,234,858	11.43%	5,408,581	27.66%	10,441,037	53.39%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	267,556	76,918	28.75%	0	0.00%	12,000	4.49%	153,210	57.26%	0	0.00%	25,427	9.50%
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	884,074	15,052	1.70%	58,596	6.63%	336,257	38.03%	474,170	53.63%	0	0.00%	0	0.00%
Never	39,720,067	3,335,290	8.40%	4,889,328	12.31%	11,392,082	28.68%	20,103,367	50.61%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,390,166	410,182	29.51%	168,921	12.15%	368,455	26.50%	442,608	31.84%	0	0.00%	0	0.00%
30 to 34	1,667,940	119,147	7.14%	309,324	18.55%	466,924	27.99%	772,545	46.32%	0	0.00%	0	0.00%
35 to 39	2,132,891	329,667	15.46%	229,358	10.75%	747,085	35.03%	826,781	38.76%	0	0.00%	0	0.00%
40 to 44	3,834,630	528,194	13.77%	578,596	15.09%	1,095,485	28.57%	1,632,354	42.57%	0	0.00%	0	0.00%
45 to 49	5,016,951	640,875	12.77%	645,092	12.86%	1,211,090	24.14%	2,519,894	50.23%	0	0.00%	0	0.00%
50 to 54	7,197,564	541,514	7.52%	911,385	12.66%	2,291,277	31.83%	3,453,388	47.98%	0	0.00%	0	0.00%
55 or older	2,247,737	220,110	9.79%	374,534	16.66%	703,421	31.29%	949,672	42.25%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	17,811,880	746,897	4.19%	1,908,540	10.71%	5,023,967	28.21%	10,107,048	56.74%	0	0.00%	25,427	0.14%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	4,012,962	322,878	8.05%	258,274	6.44%	1,043,461	26.00%	2,388,349	59.52%	0	0.00%	0	0.00%
1	4,743,482	325,671	6.87%	776,977	16.38%	1,014,718	21.39%	2,626,116	55.36%	0	0.00%	0	0.00%
2	10,672,802	1,032,345	9.67%	1,170,399	10.97%	2,850,324	26.71%	5,594,307	52.42%	0	0.00%	25,427	0.24%
More than 2	22,062,090	1,901,621	8.62%	2,963,526	13.43%	7,061,485	32.01%	10,135,458	45.94%	0	0.00%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	9,409,713	517,125	5.50%	1,242,362	13.20%	2,375,768	25.25%	5,249,031	55.78%	0	0.00%	25,427	0.27%
1	6,701,306	512,439	7.65%	744,057	11.10%	2,069,541	30.88%	3,375,269	50.37%	0	0.00%	0	0.00%
2	12,237,832	1,056,505	8.63%	1,619,051	13.23%	3,383,624	27.65%	6,178,653	50.49%	0	0.00%	0	0.00%
More than 2	13,068,928	1,496,445	11.45%	1,511,251	11.56%	4,163,814	31.86%	5,897,418	45.13%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	477,152	64,696	13.56%	146,758	30.76%	24,980	5.24%	240,718	50.45%	0	0.00%	0	0.00%
None	12,464,366	993,680	7.97%	1,826,759	14.66%	3,559,518	28.56%	6,058,981	48.61%	0	0.00%	25,427	0.20%
1	4,283,563	238,801	5.57%	413,477	9.65%	1,171,853	27.36%	2,459,432	57.42%	0	0.00%	0	0.00%
2	2,177,127	130,607	6.00%	366,876	16.85%	762,515	35.02%	917,129	42.13%	0	0.00%	0	0.00%
More than 2	878,903	38,750	4.41%	226,878	25.81%	268,095	30.50%	345,180	39.27%	0	0.00%	0	0.00%
N/A - Vaginal delivery/deliveries only	21,284,061	2,115,980	9.94%	2,188,428	10.28%	6,205,785	29.16%	10,773,868	50.62%	0	0.00%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,961,702	422,504	7.09%	715,920	12.01%	1,409,533	23.64%	3,388,318	56.83%	0	0.00%	25,427	0.43%
Yes	6,966,434	747,806	10.73%	715,913	10.28%	2,020,318	29.00%	3,482,397	49.99%	0	0.00%	0	0.00%
No	28,431,631	2,387,878	8.40%	3,711,899	13.06%	8,429,550	29.65%	13,902,304	48.90%	0	0.00%	0	0.00%
Don't know	205,405	24,325	11.84%	25,444	12.39%	133,346	64.92%	22,290	10.85%	0	0.00%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,167,107	446,829	7.25%	741,364	12.02%	1,542,879	25.02%	3,410,608	55.30%	0	0.00%	25,427	0.41%
35 years old or younger	6,388,489	634,909	9.94%	667,201	10.44%	1,731,679	27.11%	3,354,700	52.51%	0	0.00%	0	0.00%
36 years or older	577,944	112,898	19.53%	48,712	8.43%	288,639	49.94%	127,696	22.09%	0	0.00%	0	0.00%
N/A - Did not have baby weighing 9 lbs or more	28,431,631	2,387,878	8.40%	3,711,899	13.06%	8,429,550	29.65%	13,902,304	48.90%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,948,740	99,627	5.11%	457,646	23.48%	366,072	18.79%	999,968	51.31%	0	0.00%	25,427	1.30%
29 years old or younger	26,595,475	2,555,698	9.61%	3,245,434	12.20%	8,218,416	30.90%	12,575,927	47.29%	0	0.00%	0	0.00%
30 - 40 years old	2,256,437	139,494	6.18%	347,769	15.41%	551,246	24.43%	1,217,928	53.98%	0	0.00%	0	0.00%
N/A - Never pregnant	4,012,962	322,878	8.05%	258,274	6.44%	1,043,461	26.00%	2,388,349	59.52%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	6,481,179	420,455	6.49%	769,496	11.87%	1,813,552	27.98%	3,477,676	53.66%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,512,359	119,640	4.76%	457,646	18.22%	468,674	18.65%	1,440,972	57.36%	0	0.00%	25,427	1.01%
29 years old or younger	18,932,870	1,935,233	10.22%	2,452,647	12.95%	5,795,800	30.61%	8,749,191	46.21%	0	0.00%	0	0.00%
30 - 40 years old	14,850,293	1,036,006	6.98%	1,834,641	12.35%	4,352,237	29.31%	7,627,409	51.36%	0	0.00%	0	0.00%
41 years or older	862,532	40,030	4.64%	62,320	7.23%	306,254	35.51%	453,927	52.63%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	4,012,962	322,878	8.05%	258,274	6.44%	1,043,461	26.00%	2,388,349	59.52%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,106,953	1,672,297	15.06%	1,630,536	14.68%	3,130,207	28.18%	4,673,912	42.08%	0	0.00%	0	0.00%
No	30,458,218	1,910,217	6.27%	3,538,640	11.62%	8,862,539	29.10%	16,121,396	52.93%	0	0.00%	25,427	0.08%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
35 years old or younger	2,928,483	622,743	21.27%	398,526	13.61%	711,046	24.28%	1,196,168	40.85%	0	0.00%	0	0.00%
36 - 50 years old	6,136,538	853,003	13.90%	855,140	13.94%	1,823,503	29.72%	2,604,893	42.45%	0	0.00%	0	0.00%
51 years or older	1,791,666	116,794	6.52%	376,870	21.03%	501,486	27.99%	796,516	44.46%	0	0.00%	0	0.00%
Refused/Don't know	250,266	79,757	31.87%	0	0.00%	94,173	37.63%	76,335	30.50%	0	0.00%	0	0.00%
N/A - Did not have procedure	30,458,218	1,910,217	6.27%	3,538,640	11.62%	8,862,539	29.10%	16,121,396	52.93%	0	0.00%	25,427	0.08%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,167,397	64,696	5.54%	306,483	26.25%	297,309	25.47%	498,908	42.74%	0	0.00%	0	0.00%
Yes	5,650,720	626,485	11.09%	623,051	11.03%	1,935,044	34.24%	2,466,139	43.64%	0	0.00%	0	0.00%
No	34,370,000	2,869,751	8.35%	4,135,454	12.03%	9,715,685	28.27%	17,623,682	51.28%	0	0.00%	25,427	0.07%
Don't know	377,055	21,581	5.72%	104,187	27.63%	44,709	11.86%	206,578	54.79%	0	0.00%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,544,452	86,277	5.59%	410,670	26.59%	342,018	22.14%	705,486	45.68%	0	0.00%	0	0.00%
35 years old or younger	1,140,620	232,956	20.42%	125,408	10.99%	322,785	28.30%	459,471	40.28%	0	0.00%	0	0.00%
36 - 50 years old	3,156,611	253,919	8.04%	300,977	9.53%	1,137,097	36.02%	1,464,617	46.40%	0	0.00%	0	0.00%
51 years or older	1,210,967	126,748	10.47%	196,666	16.24%	421,837	34.83%	465,716	38.46%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	34,370,000	2,869,751	8.35%	4,135,454	12.03%	9,715,685	28.27%	17,623,682	51.28%	0	0.00%	25,427	0.07%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	41,565,171	3,582,514	8.62%	5,169,176	12.44%	11,992,746	28.85%	20,795,308	50.03%	0	0.00%	25,427	0.06%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,070,262	1,672,297	15.11%	1,593,845	14.40%	3,130,207	28.28%	4,673,912	42.22%	0	0.00%	0	0.00%
No	30,494,910	1,910,217	6.26%	3,575,331	11.72%	8,862,539	29.06%	16,121,396	52.87%	0	0.00%	25,427	0.08%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.043: How frequently does this (referring to KI.Q.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.