

Table F03 - Prevalence of urge urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*
20-29	12,825,480	1,845,784	14.39%	9.49%	19.29%	10,979,696	85.61%
30-39	18,943,946	3,089,960	16.31%	11.51%	21.11%	15,853,987	83.69%
40-49	22,500,594	4,893,619	21.75%	17.71%	25.78%	17,606,976	78.25%
50-59	17,533,327	4,762,544	27.16%	22.40%	31.92%	12,770,783	72.84%
60-64	5,474,602	1,770,182	32.33%	23.61%	41.06%	3,704,420	67.67%
65-69	5,240,754	1,696,659	32.37%	23.81%	40.94%	3,544,095	67.63%
70-74	4,310,387	1,577,521	36.60%	23.35%	49.85%	2,732,866	63.40%
75-79	4,281,324	1,913,177	44.69%	36.06%	53.31%	2,368,146	55.31%
80-84	2,864,438	1,335,229	46.61%	37.78%	55.45%	1,529,209	53.39%
85+	2,142,995	1,019,446	47.57%	27.92%	67.22%	1,123,549	52.43%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,044,522	1,221,405	20.21%	16.93%	23.48%	4,823,117	79.79%
Other Hispanic	5,721,649	1,371,547	23.97%	15.64%	32.30%	4,350,103	76.03%
Non-Hispanic White	69,089,229	17,764,803	25.71%	22.58%	28.85%	51,324,426	74.29%
Non-Hispanic Black	10,606,305	2,277,418	21.47%	17.97%	24.97%	8,328,887	78.53%
Other Race - Including Multi-Racial	4,925,900	1,290,578	26.20%	14.69%	37.71%	3,635,321	73.80%
Education	-	-	-	-	-	-	-
Missing	2,566,250	533,433	20.79%	10.63%	30.94%	2,032,817	79.21%
Less Than 9th Grade	6,983,938	2,352,307	33.68%	26.12%	41.24%	4,631,631	66.32%
9-11th Grade	11,493,555	2,903,325	25.26%	20.03%	30.49%	8,590,230	74.74%
High School Grad/GED or Equivalent	75,136,791	18,121,021	24.12%	21.92%	26.31%	57,015,770	75.88%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	606,658	105,325	17.36%	(2.21%)	36.93%	501,332	82.64%
Married	56,394,435	12,998,151	23.05%	20.26%	25.84%	43,396,284	76.95%
Widowed	9,939,760	4,277,040	43.03%	34.59%	51.47%	5,662,720	56.97%
Divorced	9,569,819	2,507,867	26.21%	21.72%	30.70%	7,061,952	73.79%
Separated	3,664,817	830,030	22.65%	13.68%	31.62%	2,834,787	77.35%
Never Married	10,695,115	2,685,551	25.11%	18.28%	31.94%	8,009,564	74.89%
Living with Partner	5,458,082	497,652	9.12%	1.75%	16.48%	4,960,430	90.88%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,220,599	1,736,974	27.92%	21.79%	34.05%	4,483,625	72.08%
Poverty income ratio less than 1	14,133,124	3,835,148	27.14%	22.20%	32.07%	10,297,976	72.86%
Poverty income ratio 1 or greater	76,033,883	18,353,629	24.14%	21.91%	26.36%	57,680,253	75.86%
Body mass index	-	-	-	-	-	-	-
Missing	3,885,282	1,761,921	45.35%	35.44%	55.26%	2,123,361	54.65%
Normal: less than 25	36,600,681	6,524,469	17.83%	14.66%	20.99%	30,076,212	82.17%
Overweight: 25.0 - 29.9	25,489,522	6,538,274	25.65%	23.35%	27.95%	18,951,248	74.35%
Obese: 30 or higher	30,412,122	9,101,087	29.93%	27.22%	32.63%	21,311,035	70.07%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,329,836	709,752	21.31%	10.20%	32.43%	2,620,083	78.69%
Glycohemoglobin < 5.7%	75,054,536	16,548,105	22.05%	19.69%	24.41%	58,506,431	77.95%
Glycohemoglobin 5.7% - 6.4%	13,338,158	5,024,721	37.67%	30.09%	45.25%	8,313,438	62.33%
Glycohemoglobin 6.5% or higher	4,665,076	1,643,173	35.22%	24.88%	45.56%	3,021,903	64.78%
Diabetes	-	-	-	-	-	-	-
Yes	6,392,249	2,420,010	37.86%	29.85%	45.87%	3,972,239	62.14%
No	89,995,357	21,505,742	23.90%	21.92%	25.88%	68,489,615	76.10%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,779,377	10,838,738	25.94%	23.39%	28.50%	30,940,639	74.06%
No	54,515,464	13,087,014	24.01%	20.95%	27.06%	41,428,451	75.99%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	27,312,973	8,693,824	31.83%	27.97%	35.69%	18,619,149	68.17%
No	68,684,683	15,114,221	22.01%	20.04%	23.97%	53,570,462	77.99%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,316,505	2,699,779	42.74%	33.61%	51.88%	3,616,726	57.26%
Mobility limitations	13,174,109	6,100,788	46.31%	39.61%	53.01%	7,073,321	53.69%
No mobility limitations	17,642,704	5,022,436	28.47%	22.50%	34.43%	12,620,268	71.53%
N/A - Respondent not asked mobility limitation questions	59,254,288	10,102,749	17.05%	14.43%	19.67%	49,151,540	82.95%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	3,837,297	1,149,701	29.96%	15.04%	44.88%	2,687,596	70.04%
30 to 34	3,723,416	1,546,562	41.54%	25.88%	57.19%	2,176,854	58.46%
35 to 39	4,731,533	1,921,406	40.61%	31.54%	49.68%	2,810,127	59.39%
40 to 44	5,719,731	1,615,309	28.24%	19.99%	36.49%	4,104,422	71.76%
45 to 49	10,150,431	3,088,647	30.43%	21.95%	38.91%	7,061,784	69.57%
50 to 54	9,419,020	3,764,397	39.97%	32.13%	47.80%	5,654,623	60.03%
55 or older	3,441,979	1,387,354	40.31%	30.25%	50.37%	2,054,625	59.69%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	55,153,584	9,377,790	17.00%	15.16%	18.84%	45,775,794	83.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,404,672	482,168	34.33%	20.81%	47.85%	922,505	65.67%
None	15,771,213	2,498,228	15.84%	8.62%	23.06%	13,272,985	84.16%
1	10,550,872	2,331,034	22.09%	14.24%	29.94%	8,219,839	77.91%
2	18,612,397	3,932,623	21.13%	15.71%	26.55%	14,679,774	78.87%
More than 2	50,048,451	14,681,699	29.33%	26.45%	32.22%	35,366,752	70.67%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,107,547	601,966	28.56%	14.35%	42.78%	1,505,581	71.44%
29 years old or younger	56,123,966	15,812,185	28.17%	25.61%	30.73%	40,311,781	71.83%
30 - 40 years old	3,432,833	800,094	23.31%	16.15%	30.46%	2,632,739	76.69%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	2,498,228	15.84%	8.62%	23.06%	13,272,985	84.16%
Only 1 live birth - see age at last live birth	15,349,454	3,293,674	21.46%	15.24%	27.68%	12,055,780	78.54%
No pregnancies resulting in live births	3,535,523	903,674	25.56%	13.42%	37.70%	2,631,849	74.44%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	7,125,329	2,331,696	32.72%	26.34%	39.11%	4,793,633	67.28%
29 years old or younger	41,926,311	9,114,520	21.74%	19.63%	23.85%	32,811,791	78.26%
30 - 40 years old	25,810,467	7,943,374	30.78%	25.64%	35.91%	17,867,093	69.22%
41 years or older	2,115,817	1,134,260	53.61%	26.81%	80.41%	981,557	46.39%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	2,498,228	15.84%	8.62%	23.06%	13,272,985	84.16%
No pregnancies resulting in live births	3,535,523	903,674	25.56%	13.42%	37.70%	2,631,849	74.44%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	42,912,018	7,531,384	17.55%	14.97%	20.13%	35,380,634	82.45%
Yes	20,960,048	8,240,673	39.32%	34.96%	43.67%	12,719,375	60.68%
No	32,486,648	8,153,694	25.10%	21.18%	29.01%	24,332,954	74.90%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	42,973,654	7,564,128	17.60%	15.07%	20.14%	35,409,527	82.40%
35 years old or younger	7,522,654	3,024,492	40.21%	29.79%	50.62%	4,498,162	59.79%
36 - 50 years old	10,843,595	3,911,183	36.07%	29.82%	42.32%	6,932,412	63.93%
51 years or older	2,336,250	1,144,392	48.98%	34.66%	63.31%	1,191,858	51.02%
Refused/Don't know	224,806	127,863	56.88%	22.23%	91.52%	96,943	43.12%
N/A - Did not have procedure	32,486,648	8,153,694	25.10%	21.18%	29.01%	24,332,954	74.90%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2001-2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	96,387,606	23,925,751	24.82%	22.90%	26.74%	72,461,855	75.18%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	8,390,140	1,493,988	17.81%	11.42%	24.19%	6,896,152	82.19%
Yes	20,960,048	8,240,673	39.32%	34.96%	43.67%	12,719,375	60.68%
No	67,008,526	14,191,090	21.18%	19.18%	23.17%	52,817,436	78.82%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Age at screening	-	-	-	-	-	-	-
10-19	696,430	127,044	18.24%	(15.99%)	52.48%	569,386	81.76%
20-29	12,963,783	1,456,241	11.23%	6.69%	15.78%	11,507,542	88.77%
30-39	17,035,030	3,216,701	18.88%	13.03%	24.74%	13,818,329	81.12%
40-49	20,812,873	4,170,880	20.04%	15.86%	24.22%	16,641,993	79.96%
50-59	18,372,452	5,527,419	30.09%	22.81%	37.36%	12,845,033	69.91%
60-64	5,613,252	1,283,925	22.87%	14.89%	30.85%	4,329,327	77.13%
65-69	6,373,059	2,590,075	40.64%	30.41%	50.87%	3,782,984	59.36%
70-74	5,291,067	2,088,104	39.46%	23.43%	55.50%	3,202,963	60.54%
75-79	5,602,444	2,415,697	43.12%	32.57%	53.67%	3,186,747	56.88%
80-84	3,339,588	1,087,353	32.56%	23.43%	41.69%	2,252,235	67.44%
85+	2,149,983	1,051,109	48.89%	37.59%	60.19%	1,098,874	51.11%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,686,810	1,422,888	21.28%	16.23%	26.33%	5,263,922	78.72%
Other Hispanic	3,490,447	1,020,738	29.24%	14.28%	44.21%	2,469,709	70.76%
Non-Hispanic White	71,803,772	18,440,395	25.68%	22.09%	29.27%	53,363,377	74.32%
Non-Hispanic Black	11,514,225	3,321,231	28.84%	24.38%	33.31%	8,192,995	71.16%
Other Race - Including Multi-Racial	4,754,706	809,297	17.02%	8.95%	25.09%	3,945,409	82.98%
Education	-	-	-	-	-	-	-
Missing	3,272,227	912,462	27.89%	8.79%	46.98%	2,359,765	72.11%
Less Than 9th Grade	7,079,482	2,200,338	31.08%	22.41%	39.75%	4,879,144	68.92%
9-11th Grade	10,222,174	3,463,925	33.89%	25.22%	42.55%	6,758,250	66.11%
High School Grad/GED or Equivalent	77,555,073	18,399,215	23.72%	20.52%	26.93%	59,155,858	76.28%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	538,429	85,526	15.88%	(4.01%)	35.78%	452,903	84.12%
Married	55,190,860	12,323,620	22.33%	18.84%	25.82%	42,867,240	77.67%
Widowed	11,473,706	4,599,609	40.09%	33.40%	46.77%	6,874,097	59.91%
Divorced	11,142,010	3,746,586	33.63%	22.62%	44.63%	7,395,423	66.37%
Separated	2,410,096	419,989	17.43%	5.49%	29.36%	1,990,107	82.57%
Never Married	12,255,856	2,647,160	21.60%	15.73%	27.46%	9,608,696	78.40%
Living with Partner	5,186,193	1,192,059	22.99%	12.24%	33.73%	3,994,134	77.01%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	5,148,576	1,169,561	22.72%	13.35%	32.08%	3,979,015	77.28%
Poverty income ratio less than 1	13,079,770	4,158,086	31.79%	25.08%	38.50%	8,921,684	68.21%
Poverty income ratio 1 or greater	80,021,615	19,686,902	24.60%	21.78%	27.43%	60,334,714	75.40%
Body mass index	-	-	-	-	-	-	-
Missing	1,417,889	431,430	30.43%	15.01%	45.85%	986,459	69.57%
Normal: less than 25	36,280,183	6,694,721	18.45%	14.78%	22.12%	29,585,462	81.55%
Overweight: 25.0 - 29.9	27,727,409	7,688,386	27.73%	22.29%	33.17%	20,039,023	72.27%
Obese: 30 or higher	32,824,479	10,200,012	31.07%	27.34%	34.81%	22,624,467	68.93%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,302,444	628,980	19.05%	9.93%	28.17%	2,673,465	80.95%
Glycohemoglobin < 5.7%	75,269,475	17,412,020	23.13%	20.49%	25.77%	57,857,456	76.87%
Glycohemoglobin 5.7% - 6.4%	13,399,092	4,725,769	35.27%	30.36%	40.18%	8,673,323	64.73%
Glycohemoglobin 6.5% or higher	6,278,949	2,247,780	35.80%	24.97%	46.63%	4,031,169	64.20%
Diabetes	-	-	-	-	-	-	-
Yes	8,142,154	3,455,291	42.44%	34.03%	50.85%	4,686,863	57.56%
No	90,107,806	21,559,257	23.93%	20.98%	26.87%	68,548,549	76.07%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	43,300,708	12,859,201	29.70%	26.79%	32.60%	30,441,506	70.30%
No	54,918,150	12,155,348	22.13%	18.41%	25.85%	42,762,803	77.87%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,590,425	10,715,400	35.03%	31.31%	38.74%	19,875,025	64.97%
No	67,086,850	14,082,853	20.99%	17.96%	24.02%	53,003,997	79.01%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,798,836	5,460,263	46.28%	37.47%	55.09%	6,338,573	53.72%
Mobility limitations	13,102,646	4,852,008	37.03%	31.37%	42.70%	8,250,638	62.97%
No mobility limitations	20,850,571	5,878,766	28.19%	24.30%	32.09%	14,971,804	71.81%
N/A - Respondent not asked mobility limitation questions	52,497,907	8,823,512	16.81%	13.65%	19.97%	43,674,395	83.19%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	2,535,421	633,447	24.98%	16.21%	33.76%	1,901,974	75.02%
Younger than 30	2,891,152	1,102,451	38.13%	20.92%	55.34%	1,788,701	61.87%
30 to 34	4,500,186	2,166,179	48.14%	28.97%	67.30%	2,334,008	51.86%
35 to 39	5,581,951	2,300,079	41.21%	33.65%	48.76%	3,281,871	58.79%
40 to 44	6,448,152	2,136,702	33.14%	27.31%	38.96%	4,311,450	66.86%
45 to 49	10,508,053	3,579,247	34.06%	24.16%	43.97%	6,928,807	65.94%
50 to 54	9,282,520	2,980,449	32.11%	25.89%	38.33%	6,302,070	67.89%
55 or older	4,537,319	2,030,084	44.74%	33.98%	55.50%	2,507,234	55.26%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	51,728,762	8,031,301	15.53%	12.64%	18.41%	43,697,460	84.47%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,867,248	626,612	33.56%	17.57%	49.55%	1,240,636	66.44%
None	17,857,228	2,939,167	16.46%	11.52%	21.39%	14,918,061	83.54%
1	13,698,789	2,664,303	19.45%	12.97%	25.93%	11,034,486	80.55%
2	22,340,303	4,860,387	21.76%	15.52%	27.99%	17,479,916	78.24%
More than 2	42,486,392	13,924,079	32.77%	29.18%	36.36%	28,562,313	67.23%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,499,757	550,922	22.04%	11.99%	32.09%	1,948,835	77.96%
29 years old or younger	52,992,653	16,012,220	30.22%	26.18%	34.25%	36,980,433	69.78%
30 - 40 years old	4,126,416	839,379	20.34%	11.00%	29.68%	3,287,038	79.66%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	2,939,167	16.46%	11.52%	21.39%	14,918,061	83.54%
Only 1 live birth - see age at last live birth	16,014,683	3,876,784	24.21%	17.49%	30.93%	12,137,899	75.79%
No pregnancies resulting in live births	4,581,496	733,146	16.00%	4.37%	27.63%	3,848,350	84.00%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,852,334	2,055,229	29.99%	21.88%	38.10%	4,797,105	70.01%
29 years old or younger	42,675,691	12,007,836	28.14%	22.95%	33.33%	30,667,855	71.86%
30 - 40 years old	24,321,854	6,573,739	27.03%	20.90%	33.16%	17,748,115	72.97%
41 years or older	1,773,198	675,009	38.07%	23.83%	52.30%	1,098,189	61.93%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	2,939,167	16.46%	11.52%	21.39%	14,918,061	83.54%
No pregnancies resulting in live births	4,581,496	733,146	16.00%	4.37%	27.63%	3,848,350	84.00%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	41,295,719	6,629,010	16.05%	12.42%	19.69%	34,666,709	83.95%
Yes	21,148,808	8,638,932	40.85%	33.54%	48.16%	12,509,876	59.15%
No	35,742,140	9,683,313	27.09%	23.05%	31.13%	26,058,827	72.91%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	41,359,012	6,692,303	16.18%	12.54%	19.82%	34,666,709	83.82%
35 years old or younger	7,565,019	3,606,573	47.67%	34.55%	60.80%	3,958,446	52.33%
36 - 50 years old	10,954,863	3,491,696	31.87%	25.96%	37.79%	7,463,167	68.13%
51 years or older	2,435,004	1,455,269	59.76%	45.68%	73.85%	979,735	40.24%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	35,742,140	9,683,313	27.09%	23.05%	31.13%	26,058,827	72.91%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2003-2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	98,249,961	25,014,549	25.46%	22.49%	28.43%	73,235,412	74.54%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	1,936,435	536,508	27.71%	16.83%	38.58%	1,399,927	72.29%
Yes	21,148,808	8,638,932	40.85%	33.54%	48.16%	12,509,876	59.15%
No	75,101,424	15,775,816	21.01%	17.96%	24.05%	59,325,608	78.99%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Age at screening	-	-	-	-	-	-	-
10-19	398,858	58,178	14.59%	(3.26%)	32.43%	340,679	85.41%
20-29	12,756,001	1,590,931	12.47%	8.68%	16.26%	11,165,070	87.53%
30-39	17,627,319	3,254,841	18.46%	13.67%	23.26%	14,372,478	81.54%
40-49	23,800,262	4,526,778	19.02%	12.20%	25.84%	19,273,485	80.98%
50-59	19,102,084	5,282,835	27.66%	22.01%	33.31%	13,819,249	72.34%
60-64	6,594,843	2,330,182	35.33%	29.08%	41.59%	4,264,661	64.67%
65-69	5,449,338	2,237,833	41.07%	29.33%	52.81%	3,211,505	58.93%
70-74	5,004,818	2,329,200	46.54%	38.99%	54.09%	2,675,619	53.46%
75-79	4,045,527	1,408,061	34.81%	21.73%	47.88%	2,637,465	65.19%
80-84	2,919,334	1,261,821	43.22%	34.67%	51.78%	1,657,512	56.78%
85+	2,013,172	940,262	46.71%	32.42%	60.99%	1,072,910	53.29%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	6,995,209	1,628,304	23.28%	17.25%	29.30%	5,366,906	76.72%
Other Hispanic	3,345,447	879,742	26.30%	12.33%	40.26%	2,465,705	73.70%
Non-Hispanic White	72,344,228	18,283,765	25.27%	21.12%	29.43%	54,060,463	74.73%
Non-Hispanic Black	11,495,230	3,525,787	30.67%	26.30%	35.04%	7,969,444	69.33%
Other Race - Including Multi-Racial	5,531,440	903,325	16.33%	8.29%	24.37%	4,628,116	83.67%
Education	-	-	-	-	-	-	-
Missing	2,515,028	604,008	24.02%	7.99%	40.04%	1,911,020	75.98%
Less Than 9th Grade	5,421,258	1,661,873	30.65%	22.02%	39.29%	3,759,385	69.35%
9-11th Grade	11,048,856	3,465,061	31.36%	23.79%	38.93%	7,583,795	68.64%
High School Grad/GED or Equivalent	80,515,675	19,457,204	24.17%	20.86%	27.47%	61,058,471	75.83%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	745,740	36,081	4.84%	(3.38%)	13.06%	709,659	95.16%
Married	57,213,865	14,167,387	24.76%	20.65%	28.87%	43,046,478	75.24%
Widowed	9,018,832	3,356,052	37.21%	30.08%	44.34%	5,662,780	62.79%
Divorced	12,389,946	3,357,375	27.10%	19.62%	34.57%	9,032,571	72.90%
Separated	3,088,185	877,219	28.41%	18.16%	38.65%	2,210,966	71.59%
Never Married	10,375,951	2,205,102	21.25%	14.70%	27.81%	8,170,849	78.75%
Living with Partner	6,702,495	1,131,141	16.88%	9.76%	23.99%	5,571,354	83.12%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	3,235,408	1,390,005	42.96%	33.15%	52.78%	1,845,403	57.04%
Poverty income ratio less than 1	10,921,963	3,182,392	29.14%	21.75%	36.53%	7,739,571	70.86%
Poverty income ratio 1 or greater	85,554,185	20,648,525	24.14%	20.27%	28.00%	64,905,659	75.86%
Body mass index	-	-	-	-	-	-	-
Missing	922,023	491,432	53.30%	32.09%	74.50%	430,591	46.70%
Normal: less than 25	38,407,533	6,846,555	17.83%	13.88%	21.77%	31,560,977	82.17%
Overweight: 25.0 - 29.9	25,231,035	6,576,612	26.07%	20.81%	31.32%	18,654,423	73.93%
Obese: 30 or higher	35,150,965	11,306,323	32.17%	27.89%	36.44%	23,844,642	67.83%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,122,733	667,970	21.39%	9.79%	32.99%	2,454,763	78.61%
Glycohemoglobin < 5.7%	75,153,124	17,400,028	23.15%	18.83%	27.47%	57,753,096	76.85%
Glycohemoglobin 5.7% - 6.4%	15,501,614	5,047,775	32.56%	27.16%	37.97%	10,453,838	67.44%
Glycohemoglobin 6.5% or higher	5,934,085	2,105,149	35.48%	26.03%	44.92%	3,828,936	64.52%
Diabetes	-	-	-	-	-	-	-
Yes	8,498,641	3,250,395	38.25%	29.24%	47.25%	5,248,246	61.75%
No	91,212,915	21,970,527	24.09%	20.47%	27.71%	69,242,387	75.91%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,987,511	10,359,802	24.67%	20.09%	29.26%	31,627,709	75.33%
No	57,715,558	14,852,634	25.73%	22.32%	29.15%	42,862,924	74.27%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	31,090,111	10,776,742	34.66%	30.41%	38.92%	20,313,369	65.34%
No	68,353,585	14,356,119	21.00%	17.47%	24.53%	53,997,466	79.00%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	10,862,345	4,973,568	45.79%	38.31%	53.26%	5,888,777	54.21%
Mobility limitations	8,792,109	3,829,182	43.55%	31.54%	55.57%	4,962,928	56.45%
No mobility limitations	20,550,571	6,568,299	31.96%	25.90%	38.02%	13,982,272	68.04%
N/A - Respondent not asked mobility limitation questions	59,506,530	9,849,874	16.55%	13.74%	19.36%	49,656,657	83.45%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,664,983	2,532,603	38.00%	24.61%	51.39%	4,132,379	62.00%
No	92,968,325	22,688,319	24.40%	20.64%	28.17%	70,280,007	75.60%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	761,127	245,830	32.30%	4.65%	59.95%	515,297	67.70%
2 or more times a day	11,136,447	3,512,561	31.54%	26.47%	36.61%	7,623,886	68.46%
Once a day	9,094,735	2,391,390	26.29%	19.44%	33.15%	6,703,345	73.71%
2 or more time a week	10,514,091	2,694,274	25.63%	18.45%	32.80%	7,819,817	74.37%
Once a week	6,191,809	1,733,866	28.00%	20.62%	35.38%	4,457,943	72.00%
1-3 times a month	13,509,041	4,212,110	31.18%	25.51%	36.85%	9,296,931	68.82%
Never	48,418,187	10,350,050	21.38%	17.25%	25.51%	38,068,137	78.62%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	761,127	245,830	32.30%	4.65%	59.95%	515,297	67.70%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	10,514,091	2,694,274	25.63%	7.63%	45.39%	7,819,817	74.37%
Once a week	*	*	*	*	*	*	*
1-3 times a month	13,509,041	4,212,110	31.18%	18.22%	76.00%	9,296,931	68.82%
Never	48,418,187	10,350,050	21.38%	21.63%	28.24%	38,068,137	78.62%
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	480,340	38,237	7.96%	(8.43%)	24.35%	442,103	92.04%
Younger than 30	3,249,527	1,324,106	40.75%	25.60%	55.90%	1,925,421	59.25%
30 to 34	3,113,296	1,348,019	43.30%	28.43%	58.16%	1,765,277	56.70%
35 to 39	5,187,920	2,097,662	40.43%	30.85%	50.01%	3,090,257	59.57%
40 to 44	6,708,828	2,901,226	43.24%	35.49%	51.00%	3,807,602	56.76%
45 to 49	10,909,757	4,183,891	38.35%	30.28%	46.42%	6,725,866	61.65%
50 to 54	11,102,718	3,495,074	31.48%	26.22%	36.74%	7,607,644	68.52%
55 or older	3,656,590	1,010,289	27.63%	14.01%	41.25%	2,646,300	72.37%
Don't know	544,573	117,977	21.66%	(8.88%)	52.21%	426,596	78.34%
N/A - Had period in last 12 months	54,758,007	8,704,440	15.90%	12.22%	19.58%	46,053,566	84.10%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,268,216	1,773,752	10.90%	6.48%	15.32%	14,494,464	89.10%
1	12,829,048	3,109,925	24.24%	18.37%	30.11%	9,719,123	75.76%
2	20,643,676	5,878,129	28.47%	23.49%	33.46%	14,765,547	71.53%
More than 2	49,916,637	14,459,116	28.97%	23.67%	34.26%	35,457,521	71.03%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	993,141	44,723	4.50%	0.27%	8.73%	948,418	95.50%
None	32,053,075	5,294,183	16.52%	12.84%	20.20%	26,758,893	83.48%
1	14,883,888	4,287,844	28.81%	23.53%	34.09%	10,596,044	71.19%
2	21,331,484	5,683,911	26.65%	22.15%	31.14%	15,647,573	73.35%
More than 2	30,426,938	9,910,261	32.57%	26.06%	39.08%	20,516,678	67.43%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	1,016,170	44,723	4.40%	0.39%	8.41%	971,446	95.60%
None	29,973,541	8,584,518	28.64%	24.34%	32.94%	21,389,023	71.36%
1	9,918,573	1,829,322	18.44%	9.69%	27.20%	8,089,252	81.56%
2	5,209,552	1,397,266	26.82%	15.69%	37.95%	3,812,285	73.18%
More than 2	1,827,082	488,819	26.75%	17.58%	35.93%	1,338,263	73.25%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	51,585,154	12,876,274	24.96%	20.86%	29.06%	38,708,880	75.04%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	3,595,632	646,120	17.97%	7.93%	28.01%	2,949,511	82.03%
29 years old or younger	55,693,041	16,623,518	29.85%	25.07%	34.63%	39,069,523	70.15%
30 - 40 years old	3,767,991	780,039	20.70%	8.94%	32.46%	2,987,952	79.30%
Refused/Don't know	288,861	141,956	49.14%	17.41%	80.88%	146,905	50.86%
N/A - Never pregnant	16,268,216	1,773,752	10.90%	6.48%	15.32%	14,494,464	89.10%
Only 1 live birth - see age at last live birth	16,042,594	4,188,366	26.11%	20.90%	31.31%	11,854,228	73.89%
No pregnancies resulting in live births	4,055,220	1,067,171	26.32%	15.30%	37.33%	2,988,049	73.68%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	3,639,681	646,120	17.75%	7.73%	27.77%	2,993,561	82.25%
29 years old or younger	42,813,220	11,919,979	27.84%	22.93%	32.75%	30,893,241	72.16%
30 - 40 years old	31,117,341	9,123,265	29.32%	23.25%	35.39%	21,994,076	70.68%
41 years or older	1,331,628	451,924	33.94%	22.09%	45.78%	879,704	66.06%
Refused/Don't know	486,249	238,711	49.09%	15.45%	82.73%	247,539	50.91%
N/A - Never pregnant	16,268,216	1,773,752	10.90%	6.48%	15.32%	14,494,464	89.10%
No pregnancies resulting in live births	4,055,220	1,067,171	26.32%	15.30%	37.33%	2,988,049	73.68%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	44,487,469	7,570,700	17.02%	12.95%	21.09%	36,916,769	82.98%
Yes	20,961,146	8,113,279	38.71%	31.88%	45.53%	12,847,867	61.29%
No	34,229,572	9,503,575	27.76%	22.68%	32.85%	24,725,997	72.24%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	44,520,837	7,604,068	17.08%	13.05%	21.11%	36,916,769	82.92%
35 years old or younger	6,450,871	2,718,788	42.15%	33.17%	51.12%	3,732,083	57.85%
36 - 50 years old	11,493,877	4,555,075	39.63%	30.30%	48.97%	6,938,802	60.37%
51 years or older	2,278,795	602,808	26.45%	7.00%	45.90%	1,675,987	73.55%
Refused/Don't know	737,604	236,608	32.08%	10.72%	53.44%	500,996	67.92%
N/A - Did not have procedure	34,229,572	9,503,575	27.76%	22.68%	32.85%	24,725,997	72.24%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2005-2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	99,711,555	25,220,922	25.29%	21.98%	28.61%	74,490,633	74.71%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,961,146	8,113,279	38.71%	31.88%	45.53%	12,847,867	61.29%
No	78,717,041	17,074,275	21.69%	18.07%	25.32%	61,642,766	78.31%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,194,670	26,250,782	25.94%	24.34%	27.54%	74,943,888	74.06%
Age at screening	-	-	-	-	-	-	-
10-19	723,602	131,239	18.14%	(10.37%)	46.64%	592,364	81.86%
20-29	12,424,789	1,070,655	8.62%	5.17%	12.07%	11,354,134	91.38%
30-39	17,248,808	2,291,102	13.28%	10.17%	16.39%	14,957,705	86.72%
40-49	20,998,655	5,124,808	24.41%	19.48%	29.33%	15,873,847	75.59%
50-59	20,085,163	6,035,135	30.05%	24.97%	35.13%	14,050,028	69.95%
60-64	7,675,019	2,365,499	30.82%	22.64%	39.00%	5,309,520	69.18%
65-69	6,578,135	2,249,997	34.20%	26.07%	42.34%	4,328,138	65.80%
70-74	5,328,792	2,543,230	47.73%	36.16%	59.29%	2,785,562	52.27%
75-79	4,498,310	1,844,535	41.01%	33.36%	48.65%	2,653,776	58.99%
80+	5,633,398	2,594,583	46.06%	37.20%	54.92%	3,038,815	53.94%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,563,856	1,553,678	20.54%	16.70%	24.38%	6,010,178	79.46%
Other Hispanic	4,965,594	1,301,346	26.21%	22.24%	30.17%	3,664,248	73.79%
Non-Hispanic White	71,403,732	18,979,877	26.58%	24.39%	28.78%	52,423,855	73.42%
Non-Hispanic Black	12,035,915	3,610,806	30.00%	24.76%	35.24%	8,425,109	70.00%
Other Race - Including Multi-Racial	5,225,573	805,075	15.41%	9.27%	21.54%	4,420,499	84.59%
Education	-	-	-	-	-	-	-
Missing	2,863,940	780,652	27.26%	14.52%	40.00%	2,083,288	72.74%
Less Than 9th Grade	6,416,781	2,242,619	34.95%	27.79%	42.11%	4,174,162	65.05%
9-11th Grade	13,269,620	3,881,059	29.25%	25.34%	33.16%	9,388,561	70.75%
High School Grad/GED or Equivalent	78,632,714	19,334,836	24.59%	22.54%	26.64%	59,297,878	75.41%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,342,297	260,286	19.39%	3.31%	35.47%	1,082,010	80.61%
Married	57,582,615	14,074,127	24.44%	21.79%	27.09%	43,508,488	75.56%
Widowed	9,356,585	3,526,660	37.69%	33.20%	42.19%	5,829,925	62.31%
Divorced	12,570,976	4,100,623	32.62%	25.59%	39.65%	8,470,353	67.38%
Separated	2,378,944	606,331	25.49%	16.03%	34.94%	1,772,613	74.51%
Never Married	11,774,159	2,474,011	21.01%	16.56%	25.46%	9,300,148	78.99%
Living with Partner	6,162,712	1,208,744	19.61%	10.80%	28.43%	4,953,968	80.39%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	7,061,696	1,785,020	25.28%	18.08%	32.48%	5,276,676	74.72%
Poverty income ratio less than 1	13,770,564	3,819,925	27.74%	22.71%	32.77%	9,950,639	72.26%
Poverty income ratio 1 or greater	80,362,410	20,645,837	25.69%	23.67%	27.71%	59,716,573	74.31%
Body mass index	-	-	-	-	-	-	-
Missing	950,166	426,688	44.91%	24.78%	65.04%	523,478	55.09%
Normal: less than 25	35,587,048	7,329,434	20.60%	17.12%	24.07%	28,257,614	79.40%
Overweight: 25.0 - 29.9	29,281,781	7,888,995	26.94%	24.08%	29.80%	21,392,786	73.06%
Obese: 30 or higher	35,375,675	10,605,665	29.98%	25.99%	33.97%	24,770,010	70.02%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,619,352	781,126	21.58%	14.98%	28.18%	2,838,226	78.42%
Glycohemoglobin < 5.7%	68,246,278	13,407,599	19.65%	17.83%	21.46%	54,838,678	80.35%
Glycohemoglobin 5.7% - 6.4%	22,358,589	9,014,889	40.32%	36.69%	43.95%	13,343,699	59.68%
Glycohemoglobin 6.5% or higher	6,970,452	3,047,167	43.72%	35.16%	52.28%	3,923,284	56.28%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,194,670	26,250,782	25.94%	24.34%	27.54%	74,943,888	74.06%
Diabetes	-	-	-	-	-	-	-
Yes	9,392,511	4,408,790	46.94%	38.67%	55.21%	4,983,721	53.06%
No	91,802,159	21,841,991	23.79%	22.13%	25.46%	69,960,167	76.21%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,148,068	12,368,068	30.06%	27.78%	32.34%	28,780,000	69.94%
No	59,982,420	13,882,714	23.14%	20.85%	25.44%	46,099,706	76.86%
Refused	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	32,797,113	12,649,611	38.57%	34.93%	42.21%	20,147,502	61.43%
No	68,313,973	13,601,171	19.91%	18.48%	21.34%	54,712,802	80.09%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,630,095	3,900,280	33.54%	28.94%	38.13%	7,729,814	66.46%
Mobility limitations	10,714,977	4,407,431	41.13%	33.01%	49.26%	6,307,546	58.87%
No mobility limitations	20,400,274	7,499,869	36.76%	32.03%	41.50%	12,900,404	63.24%
N/A - Respondent not asked mobility limitation questions	58,449,325	10,443,202	17.87%	15.47%	20.27%	48,006,123	82.13%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,361,300	2,406,317	32.69%	23.16%	42.21%	4,954,983	67.31%
No	93,743,992	23,801,450	25.39%	23.86%	26.92%	69,942,542	74.61%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	583,024	168,164	28.84%	3.90%	53.79%	414,860	71.16%
2 or more times a day	9,905,116	2,692,415	27.18%	18.33%	36.03%	7,212,701	72.82%
Once a day	9,682,887	2,607,295	26.93%	22.33%	31.53%	7,075,592	73.07%
2 or more time a week	7,646,398	2,294,811	30.01%	25.74%	34.28%	5,351,587	69.99%
Once a week	7,022,920	2,067,749	29.44%	19.91%	38.97%	4,955,171	70.56%
1-3 times a month	14,326,336	5,053,236	35.27%	29.77%	40.77%	9,273,100	64.73%
Never	51,785,809	11,298,450	21.82%	20.37%	23.26%	40,487,359	78.18%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	583,024	168,164	28.84%	3.90%	53.79%	414,860	71.16%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	7,646,398	2,294,811	30.01%	8.26%	108.70%	5,351,587	69.99%
Once a week	*	*	*	*	*	*	*
1-3 times a month	14,326,336	5,053,236	35.27%	50.38%	82.47%	9,273,100	64.73%
Never	51,785,809	11,298,450	21.82%	23.62%	26.49%	40,487,359	78.18%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,194,670	26,250,782	25.94%	24.34%	27.54%	74,943,888	74.06%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	432,110	17,263	4.00%	(4.50%)	12.49%	414,848	96.01%
Younger than 30	3,069,584	1,092,036	35.58%	22.90%	48.25%	1,977,548	64.42%
30 to 34	3,711,338	1,445,999	38.96%	25.76%	52.16%	2,265,339	61.04%
35 to 39	5,113,159	1,852,481	36.23%	26.21%	46.25%	3,260,678	63.77%
40 to 44	7,502,270	3,215,975	42.87%	31.85%	53.88%	4,286,295	57.13%
45 to 49	11,650,306	4,624,738	39.70%	32.93%	46.46%	7,025,568	60.30%
50 to 54	10,341,790	3,240,354	31.33%	22.41%	40.26%	7,101,436	68.67%
55 or older	5,101,078	2,238,244	43.88%	34.83%	52.92%	2,862,834	56.12%
Don't know	224,950	54,362	24.17%	(0.88%)	49.21%	170,589	75.83%
N/A - Had period in last 12 months	54,048,084	8,469,331	15.67%	13.19%	18.15%	45,578,753	84.33%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,962,970	2,580,354	15.21%	10.42%	20.00%	14,382,616	84.79%
1	13,365,714	2,621,413	19.61%	13.20%	26.03%	10,744,302	80.39%
2	22,668,126	6,031,539	26.61%	22.26%	30.96%	16,636,587	73.39%
More than 2	47,917,311	14,959,853	31.22%	27.62%	34.82%	32,957,458	68.78%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	674,220	101,788	15.10%	(1.68%)	31.87%	572,432	84.90%
None	32,052,425	5,804,859	18.11%	15.63%	20.59%	26,247,566	81.89%
1	17,728,714	3,566,141	20.12%	15.86%	24.37%	14,162,573	79.88%
2	23,024,729	6,946,976	30.17%	25.84%	34.51%	16,077,753	69.83%
More than 2	27,700,262	9,816,697	35.44%	32.13%	38.75%	17,883,564	64.56%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	964,977	124,388	12.89%	(3.89%)	29.67%	840,590	87.11%
None	25,242,835	7,177,469	28.43%	25.12%	31.74%	18,065,365	71.57%
1	9,899,641	2,221,278	22.44%	15.27%	29.61%	7,678,364	77.56%
2	5,518,794	1,299,049	23.54%	15.85%	31.22%	4,219,745	76.46%
More than 2	2,981,345	732,226	24.56%	14.46%	34.66%	2,249,119	75.44%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	56,572,758	14,682,052	25.95%	23.54%	28.37%	41,890,705	74.05%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	21,543,739	3,606,796	16.74%	13.20%	20.29%	17,936,943	83.26%
Yes	13,100,217	4,326,233	33.02%	25.36%	40.69%	8,773,983	66.98%
No	66,370,995	18,280,301	27.54%	26.28%	28.81%	48,090,694	72.46%
Don't know	179,719	37,451	20.84%	(7.65%)	49.33%	142,268	79.16%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	21,723,458	3,644,248	16.78%	13.38%	20.17%	18,079,211	83.22%
35 years old or younger	12,074,632	3,922,334	32.48%	24.58%	40.39%	8,152,299	67.52%
36 years or older	932,800	395,986	42.45%	12.03%	72.88%	536,814	57.55%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	66,370,995	18,280,301	27.54%	26.28%	28.81%	48,090,694	72.46%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,194,670	26,250,782	25.94%	24.34%	27.54%	74,943,888	74.06%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	4,580,770	1,026,443	22.41%	13.37%	31.45%	3,554,327	77.59%
29 years old or younger	55,850,505	17,896,821	32.04%	30.20%	33.89%	37,953,684	67.96%
30 - 40 years old	5,156,343	1,154,942	22.40%	12.47%	32.33%	4,001,401	77.60%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	16,962,970	2,580,354	15.21%	10.42%	20.00%	14,382,616	84.79%
Only 1 live birth - see age at last live birth	18,089,584	3,505,690	19.38%	14.95%	23.81%	14,583,894	80.62%
No pregnancies resulting in live births	491,126	73,133	14.89%	(0.17%)	29.95%	417,994	85.11%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	5,813,479	1,433,545	24.66%	16.45%	32.86%	4,379,934	75.34%
29 years old or younger	46,337,613	12,999,816	28.05%	26.18%	29.93%	33,337,797	71.95%
30 - 40 years old	29,435,374	8,310,231	28.23%	25.16%	31.31%	21,125,143	71.77%
41 years or older	1,806,507	717,691	39.73%	16.18%	63.28%	1,088,816	60.27%
Refused/Don't know	347,601	136,012	39.13%	16.97%	61.29%	211,589	60.87%
N/A - Never pregnant	16,962,970	2,580,354	15.21%	10.42%	20.00%	14,382,616	84.79%
No pregnancies resulting in live births	491,126	73,133	14.89%	(0.17%)	29.95%	417,994	85.11%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,875,967	9,363,305	42.80%	38.61%	47.00%	12,512,662	57.20%
No	79,160,905	16,849,984	21.29%	19.58%	22.99%	62,310,921	78.71%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	6,125,811	2,492,721	40.69%	33.81%	47.58%	3,633,090	59.31%
36 - 50 years old	12,707,817	5,476,908	43.10%	35.59%	50.61%	7,230,910	56.90%
51 years or older	2,470,853	1,199,334	48.54%	38.36%	58.72%	1,271,519	51.46%
Refused/Don't know	571,486	194,342	34.01%	6.77%	61.24%	377,144	65.99%
N/A - Did not have procedure	79,160,905	16,849,984	21.29%	19.58%	22.99%	62,310,921	78.71%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,803,240	143,145	7.94%	0.34%	15.53%	1,660,095	92.06%
Yes	11,782,493	5,305,756	45.03%	37.04%	53.02%	6,476,737	54.97%
No	86,948,198	20,579,534	23.67%	21.77%	25.57%	66,368,664	76.33%
Don't know	660,739	222,347	33.65%	7.80%	59.50%	438,392	66.35%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,463,979	365,492	14.83%	4.15%	25.52%	2,098,487	85.17%
35 years old or younger	2,315,265	843,180	36.42%	22.11%	50.72%	1,472,085	63.58%
36 - 50 years old	6,618,268	3,322,455	50.20%	38.28%	62.12%	3,295,814	49.80%
51 years or older	2,584,749	1,024,360	39.63%	30.20%	49.06%	1,560,389	60.37%
Refused/Don't know	264,211	115,761	43.81%	19.01%	68.62%	148,450	56.19%
N/A - Did not have procedure	86,948,198	20,579,534	23.67%	21.77%	25.57%	66,368,664	76.33%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2007-2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	101,194,670	26,250,782	25.94%	24.34%	27.54%	74,943,888	74.06%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,299,155	9,166,381	43.04%	38.70%	47.37%	12,132,775	56.96%
No	79,737,716	17,046,908	21.38%	19.73%	23.03%	62,690,808	78.62%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,834,970	25,417,806	25.98%	23.15%	28.81%	72,417,164	74.02%
Age at screening	-	-	-	-	-	-	-
10-19	353,109	94,927	26.88%	4.53%	49.24%	258,182	73.12%
20-29	12,072,924	1,840,192	15.24%	11.56%	18.93%	10,232,732	84.76%
30-39	15,950,025	2,687,312	16.85%	12.96%	20.74%	13,262,713	83.15%
40-49	19,772,641	4,635,591	23.44%	16.79%	30.10%	15,137,050	76.56%
50-59	20,704,188	5,211,647	25.17%	18.75%	31.59%	15,492,541	74.83%
60-64	7,767,115	2,437,163	31.38%	19.28%	43.47%	5,329,952	68.62%
65-69	6,636,364	2,072,827	31.23%	22.70%	39.77%	4,563,537	68.77%
70-74	5,133,707	2,365,064	46.07%	40.54%	51.59%	2,768,643	53.93%
75-79	4,161,433	1,777,635	42.72%	34.01%	51.42%	2,383,798	57.28%
80+	5,283,463	2,295,448	43.45%	34.13%	52.76%	2,988,015	56.55%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,365,117	2,092,224	28.41%	24.03%	32.78%	5,272,893	71.59%
Other Hispanic	4,864,518	1,184,251	24.34%	19.81%	28.88%	3,680,267	75.66%
Non-Hispanic White	67,979,452	17,480,786	25.71%	22.25%	29.18%	50,498,666	74.29%
Non-Hispanic Black	11,442,102	3,812,462	33.32%	26.36%	40.27%	7,629,641	66.68%
Other Race - Including Multi-Racial	6,183,781	848,083	13.71%	7.02%	20.41%	5,335,697	86.29%
Education	-	-	-	-	-	-	-
Missing	2,266,489	635,598	28.04%	11.43%	44.65%	1,630,891	71.96%
Less Than 9th Grade	5,812,537	1,844,541	31.73%	25.38%	38.09%	3,967,996	68.27%
9-11th Grade	11,569,166	3,504,140	30.29%	23.21%	37.37%	8,065,026	69.71%
High School Grad/GED or Equivalent	77,985,423	19,358,370	24.82%	21.91%	27.74%	58,627,053	75.18%
Refused	*	*	*	*	*	*	*
Don't Know	156,713	75,158	47.96%	15.40%	80.52%	81,555	52.04%
Marital status	-	-	-	-	-	-	-
Missing	353,109	94,927	26.88%	4.53%	49.24%	258,182	73.12%
Married	56,002,320	13,148,028	23.48%	19.60%	27.36%	42,854,292	76.52%
Widowed	8,890,620	3,965,487	44.60%	35.70%	53.50%	4,925,133	55.40%
Divorced	11,116,682	2,951,626	26.55%	19.91%	33.19%	8,165,057	73.45%
Separated	2,211,003	849,896	38.44%	22.06%	54.82%	1,361,106	61.56%
Never Married	11,937,752	2,776,325	23.26%	19.31%	27.20%	9,161,427	76.74%
Living with Partner	7,120,836	1,455,784	20.44%	14.35%	26.53%	5,665,053	79.56%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,365,743	1,897,549	29.81%	22.30%	37.32%	4,468,194	70.19%
Poverty income ratio less than 1	13,921,006	4,290,978	30.82%	26.17%	35.47%	9,630,027	69.18%
Poverty income ratio 1 or greater	77,548,222	19,229,279	24.80%	21.51%	28.08%	58,318,943	75.20%
Body mass index	-	-	-	-	-	-	-
Missing	432,161	261,788	60.58%	27.19%	93.96%	170,373	39.42%
Normal: less than 25	34,188,422	5,714,127	16.71%	13.82%	19.60%	28,474,295	83.29%
Overweight: 25.0 - 29.9	27,289,741	7,234,091	26.51%	20.64%	32.37%	20,055,650	73.49%
Obese: 30 or higher	35,924,646	12,207,800	33.98%	30.60%	37.37%	23,716,846	66.02%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,219,727	853,886	26.52%	19.84%	33.20%	2,365,841	73.48%
Glycohemoglobin < 5.7%	62,637,820	13,935,553	22.25%	19.94%	24.56%	48,702,268	77.75%
Glycohemoglobin 5.7% - 6.4%	25,204,944	7,795,151	30.93%	25.73%	36.12%	17,409,793	69.07%
Glycohemoglobin 6.5% or higher	6,772,479	2,833,217	41.83%	31.08%	52.59%	3,939,262	58.17%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,834,970	25,417,806	25.98%	23.15%	28.81%	72,417,164	74.02%
Diabetes	-	-	-	-	-	-	-
Yes	8,795,160	4,128,037	46.94%	38.31%	55.56%	4,667,123	53.06%
No	89,039,811	21,289,769	23.91%	20.99%	26.83%	67,750,041	76.09%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	39,138,315	11,091,913	28.34%	23.63%	33.05%	28,046,402	71.66%
No	58,696,655	14,325,894	24.41%	21.20%	27.62%	44,370,762	75.59%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,250,076	11,606,108	38.37%	33.97%	42.77%	18,643,968	61.63%
No	67,523,526	13,796,321	20.43%	17.76%	23.10%	53,727,205	79.57%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	11,418,591	5,124,229	44.88%	36.96%	52.79%	6,294,362	55.12%
Mobility limitations	9,634,811	3,894,488	40.42%	33.22%	47.62%	5,740,323	59.58%
No mobility limitations	20,709,889	6,222,816	30.05%	23.04%	37.06%	14,487,074	69.95%
N/A - Respondent not asked mobility limitation questions	56,071,679	10,176,273	18.15%	15.48%	20.82%	45,895,406	81.85%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,026,447	2,780,568	39.57%	30.04%	49.11%	4,245,879	60.43%
No	90,732,584	22,630,326	24.94%	22.03%	27.86%	68,102,259	75.06%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	514,556	156,776	30.47%	2.39%	58.54%	357,780	69.53%
2 or more times a day	9,005,246	2,801,521	31.11%	23.70%	38.52%	6,203,725	68.89%
Once a day	8,275,588	2,380,873	28.77%	21.59%	35.95%	5,894,715	71.23%
2 or more time a week	8,149,977	2,678,116	32.86%	24.89%	40.83%	5,471,861	67.14%
Once a week	6,154,232	1,874,622	30.46%	24.20%	36.72%	4,279,611	69.54%
1-3 times a month	13,260,263	3,679,787	27.75%	21.21%	34.30%	9,580,476	72.25%
Never	52,378,637	11,808,270	22.54%	19.22%	25.87%	40,570,368	77.46%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	514,556	156,776	30.47%	1.13%	53.73%	357,780	69.53%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	8,149,977	2,678,116	32.86%	10.11%	51.42%	5,471,861	67.14%
Once a week	6,154,232	1,874,622	30.46%	36.20%	100.20%	4,279,611	69.54%
1-3 times a month	13,260,263	3,679,787	27.75%	25.87%	67.93%	9,580,476	72.25%
Never	52,378,637	11,808,270	22.54%	22.52%	28.23%	40,570,368	77.46%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,834,970	25,417,806	25.98%	23.15%	28.81%	72,417,164	74.02%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	624,425	84,593	13.55%	(1.44%)	28.54%	539,833	86.45%
Younger than 30	2,882,766	1,039,094	36.05%	20.14%	51.95%	1,843,673	63.96%
30 to 34	3,254,943	1,006,929	30.94%	20.13%	41.74%	2,248,014	69.06%
35 to 39	4,541,323	1,347,858	29.68%	18.01%	41.35%	3,193,465	70.32%
40 to 44	7,479,695	2,873,502	38.42%	33.30%	43.54%	4,606,193	61.58%
45 to 49	9,847,084	3,356,466	34.09%	27.24%	40.93%	6,490,618	65.91%
50 to 54	13,757,434	4,611,844	33.52%	26.00%	41.05%	9,145,590	66.48%
55 or older	5,142,489	1,942,557	37.77%	26.97%	48.58%	3,199,932	62.23%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	50,018,311	9,032,022	18.06%	14.93%	21.19%	40,986,290	81.94%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	17,565,210	2,652,878	15.10%	11.57%	18.63%	14,912,332	84.90%
1	12,038,159	2,655,546	22.06%	16.02%	28.10%	9,382,614	77.94%
2	23,689,401	5,674,202	23.95%	17.46%	30.44%	18,015,199	76.05%
More than 2	44,437,847	14,435,181	32.48%	29.50%	35.47%	30,002,666	67.52%
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	310,780	135,580	43.63%	16.00%	71.25%	175,200	56.37%
None	32,886,315	5,869,186	17.85%	14.22%	21.47%	27,017,130	82.15%
1	14,225,353	3,586,898	25.21%	19.34%	31.09%	10,638,454	74.79%
2	23,367,578	6,391,261	27.35%	19.51%	35.20%	16,976,316	72.65%
More than 2	27,044,944	9,434,881	34.89%	30.56%	39.21%	17,610,064	65.11%
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	779,924	233,202	29.90%	11.71%	48.09%	546,722	70.10%
None	25,110,253	7,632,579	30.40%	26.06%	34.73%	17,477,675	69.60%
1	9,753,274	2,299,292	23.57%	15.94%	31.21%	7,453,982	76.43%
2	5,288,336	1,312,038	24.81%	17.61%	32.01%	3,976,297	75.19%
More than 2	2,314,074	623,369	26.94%	13.98%	39.90%	1,690,705	73.06%
N/A - Vaginal delivery/deliveries only	54,589,109	13,317,326	24.40%	21.21%	27.59%	41,271,783	75.60%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	22,967,058	3,861,897	16.81%	12.51%	21.12%	19,105,161	83.19%
Yes	14,458,970	4,855,510	33.58%	27.07%	40.09%	9,603,461	66.42%
No	60,016,411	16,538,839	27.56%	24.10%	31.02%	43,477,572	72.44%
Don't know	392,531	161,561	41.16%	11.23%	71.08%	230,970	58.84%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	23,359,589	4,023,458	17.22%	13.17%	21.28%	19,336,131	82.78%
35 years old or younger	13,118,409	4,345,629	33.13%	25.58%	40.67%	8,772,780	66.87%
36 years or older	1,284,225	491,599	38.28%	24.89%	51.67%	792,626	61.72%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	60,016,411	16,538,839	27.56%	24.10%	31.02%	43,477,572	72.44%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,834,970	25,417,806	25.98%	23.15%	28.81%	72,417,164	74.02%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	5,401,849	1,209,019	22.38%	8.87%	35.89%	4,192,830	77.62%
29 years old or younger	54,084,899	16,270,236	30.08%	27.39%	32.78%	37,814,663	69.92%
30 - 40 years old	5,231,391	1,542,350	29.48%	17.70%	41.26%	3,689,041	70.52%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,565,210	2,652,878	15.10%	11.57%	18.63%	14,912,332	84.90%
Only 1 live birth - see age at last live birth	15,122,928	3,527,466	23.33%	18.77%	27.88%	11,595,462	76.67%
No pregnancies resulting in live births	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	6,887,500	1,344,681	19.52%	8.26%	30.79%	5,542,819	80.48%
29 years old or younger	41,075,647	12,042,012	29.32%	25.88%	32.75%	29,033,635	70.68%
30 - 40 years old	29,556,615	8,339,256	28.21%	23.12%	33.31%	21,217,359	71.79%
41 years or older	2,166,264	727,954	33.60%	23.24%	43.97%	1,438,310	66.40%
Refused/Don't know	254,181	95,168	37.44%	0.30%	74.58%	159,013	62.56%
N/A - Never pregnant	17,565,210	2,652,878	15.10%	11.57%	18.63%	14,912,332	84.90%
No pregnancies resulting in live births	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,537,624	7,934,861	38.64%	33.56%	43.71%	12,602,763	61.36%
No	77,219,045	17,454,893	22.60%	19.65%	25.56%	59,764,152	77.40%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	5,391,230	2,062,218	38.25%	28.29%	48.21%	3,329,012	61.75%
36 - 50 years old	11,048,011	4,064,475	36.79%	28.41%	45.17%	6,983,537	63.21%
51 years or older	3,681,428	1,532,708	41.63%	27.96%	55.31%	2,148,720	58.37%
Refused/Don't know	416,955	275,460	66.06%	40.89%	91.24%	141,495	33.94%
N/A - Did not have procedure	77,219,045	17,454,893	22.60%	19.65%	25.56%	59,764,152	77.40%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,954,499	428,996	21.95%	10.13%	33.77%	1,525,503	78.05%
Yes	10,117,822	3,927,401	38.82%	31.23%	46.40%	6,190,421	61.18%
No	85,027,169	20,752,776	24.41%	21.66%	27.16%	64,274,393	75.59%
Refused	*	*	*	*	*	*	*
Don't know	723,930	308,633	42.63%	10.17%	75.09%	415,296	57.37%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	2,689,979	737,629	27.42%	13.53%	41.31%	1,952,350	72.58%
35 years old or younger	2,196,357	891,251	40.58%	27.80%	53.36%	1,305,105	59.42%
36 - 50 years old	5,245,669	2,089,328	39.83%	29.81%	49.85%	3,156,341	60.17%
51 years or older	2,452,490	766,122	31.24%	19.22%	43.25%	1,686,368	68.76%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	85,027,169	20,752,776	24.41%	21.66%	27.16%	64,274,393	75.59%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03 - Prevalence of urge urinary incontinence among women, 2009-2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	97,834,970	25,417,806	25.98%	23.15%	28.81%	72,417,164	74.02%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,338,065	7,918,952	38.94%	33.88%	44.00%	12,419,113	61.06%
No	77,418,604	17,470,803	22.57%	19.61%	25.53%	59,947,802	77.43%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.