

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	1,845,784	108,099	5.86%	91,747	4.97%	541,822	29.35%	1,104,116	59.82%	0	0.00%	0	0.00%
30-39	3,089,960	264,958	8.57%	360,809	11.68%	725,447	23.48%	1,738,745	56.27%	0	0.00%	0	0.00%
40-49	4,893,619	524,404	10.72%	906,455	18.52%	1,303,257	26.63%	2,159,503	44.13%	0	0.00%	0	0.00%
50-59	4,762,544	525,686	11.04%	1,108,030	23.27%	1,288,057	27.05%	1,840,772	38.65%	0	0.00%	0	0.00%
60-64	1,770,182	429,919	24.29%	703,744	39.76%	372,716	21.06%	263,803	14.90%	0	0.00%	0	0.00%
65-69	1,696,659	446,356	26.31%	509,522	30.03%	412,346	24.30%	328,435	19.36%	0	0.00%	0	0.00%
70-74	1,577,521	432,496	27.42%	296,576	18.80%	653,549	41.43%	194,900	12.35%	0	0.00%	0	0.00%
75-79	1,913,177	502,269	26.25%	451,625	23.61%	541,439	28.30%	376,834	19.70%	41,011	2.14%	0	0.00%
80-84	1,335,229	414,735	31.06%	324,824	24.33%	397,768	29.79%	168,227	12.60%	29,674	2.22%	0	0.00%
85+	1,019,446	400,524	39.29%	190,182	18.66%	297,050	29.14%	114,646	11.25%	17,043	1.67%	0	0.00%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,221,405	129,488	10.60%	266,657	21.83%	259,051	21.21%	566,208	46.36%	0	0.00%	0	0.00%
Other Hispanic	1,371,547	334,115	24.36%	188,569	13.75%	269,775	19.67%	579,087	42.22%	0	0.00%	0	0.00%
Non-Hispanic White	17,764,803	2,827,019	15.91%	3,814,785	21.47%	5,018,880	28.25%	6,016,390	33.87%	87,728	0.49%	0	0.00%
Non-Hispanic Black	2,277,418	504,492	22.15%	565,093	24.81%	730,241	32.06%	477,592	20.97%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	1,290,578	254,330	19.71%	108,410	8.40%	255,504	19.80%	672,334	52.10%	0	0.00%	0	0.00%
<b>Education</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	533,433	114,001	21.37%	89,456	16.77%	192,086	36.01%	137,890	25.85%	0	0.00%	0	0.00%
Less Than 9th Grade	2,352,307	649,788	27.62%	571,194	24.28%	410,075	17.43%	663,195	28.19%	58,054	2.47%	0	0.00%
9-11th Grade	2,903,325	391,160	13.47%	652,482	22.47%	1,147,917	39.54%	711,766	24.52%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	18,121,021	2,891,070	15.95%	3,625,960	20.01%	4,783,373	26.40%	6,790,944	37.48%	29,674	0.16%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Marital status</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	12,998,151	1,848,190	14.22%	2,753,685	21.19%	3,652,248	28.10%	4,726,985	36.37%	17,043	0.13%	0	0.00%
Widowed	4,277,040	1,340,000	31.33%	1,171,428	27.39%	1,127,909	26.37%	567,017	13.26%	70,685	1.65%	0	0.00%
Divorced	2,507,867	356,496	14.22%	427,672	17.05%	575,215	22.94%	1,148,485	45.80%	0	0.00%	0	0.00%
Separated	830,030	64,182	7.73%	132,911	16.01%	192,953	23.25%	439,985	53.01%	0	0.00%	0	0.00%
Never Married	2,685,551	424,439	15.80%	344,998	12.85%	786,018	29.27%	1,130,096	42.08%	0	0.00%	0	0.00%
Living with Partner	497,652	16,139	3.24%	88,685	17.82%	105,073	21.11%	287,755	57.82%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,736,974	269,352	15.51%	307,830	17.72%	352,337	20.28%	766,444	44.13%	41,011	2.36%	0	0.00%
Poverty income ratio less than 1	3,835,148	850,164	22.17%	698,032	18.20%	1,187,058	30.95%	1,099,894	28.68%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	18,353,629	2,929,929	15.96%	3,937,652	21.45%	4,994,057	27.21%	6,445,274	35.12%	46,717	0.25%	0	0.00%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,761,921	778,417	44.18%	131,711	7.48%	504,781	28.65%	317,338	18.01%	29,674	1.68%	0	0.00%
Normal: less than 25	6,524,469	841,852	12.90%	1,304,570	20.00%	1,461,306	22.40%	2,916,741	44.70%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	6,538,274	849,064	12.99%	1,308,233	20.01%	2,024,121	30.96%	2,298,801	35.16%	58,054	0.89%	0	0.00%
Obese: 30 or higher	9,101,087	1,580,112	17.36%	2,199,001	24.16%	2,543,244	27.94%	2,778,731	30.53%	0	0.00%	0	0.00%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	709,752	267,155	37.64%	65,142	9.18%	219,105	30.87%	158,350	22.31%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	16,548,105	2,021,904	12.22%	3,317,811	20.05%	4,415,552	26.68%	6,746,120	40.77%	46,717	0.28%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	5,024,721	1,154,638	22.98%	1,104,488	21.98%	1,584,871	31.54%	1,180,724	23.50%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,643,173	605,748	36.86%	456,073	27.76%	313,924	19.10%	226,417	13.78%	41,011	2.50%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Diabetes</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,420,010	992,341	41.01%	434,223	17.94%	564,346	23.32%	388,089	16.04%	41,011	1.69%	0	0.00%
No	21,505,742	3,057,104	14.22%	4,509,292	20.97%	5,969,106	27.76%	7,923,523	36.84%	46,717	0.22%	0	0.00%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	10,838,738	2,160,580	19.93%	1,942,712	17.92%	3,032,963	27.98%	3,702,482	34.16%	0	0.00%	0	0.00%
No	13,087,014	1,888,865	14.43%	3,000,803	22.93%	3,500,488	26.75%	4,609,130	35.22%	87,728	0.67%	0	0.00%
<b>High blood pressure</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	8,693,824	2,254,770	25.94%	2,018,080	23.21%	2,458,697	28.28%	1,962,276	22.57%	0	0.00%	0	0.00%
No	15,114,221	1,794,675	11.87%	2,925,434	19.36%	4,074,755	26.96%	6,231,628	41.23%	87,728	0.58%	0	0.00%
<b>Mobility limitation</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,699,779	1,040,440	38.54%	393,516	14.58%	867,509	32.13%	357,303	13.23%	41,011	1.52%	0	0.00%
Mobility limitations	6,100,788	1,579,270	25.89%	1,971,865	32.32%	1,458,272	23.90%	1,044,664	17.12%	46,717	0.77%	0	0.00%
No mobility limitations	5,022,436	769,564	15.32%	1,025,191	20.41%	1,631,588	32.49%	1,596,093	31.78%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	10,102,749	660,171	6.53%	1,552,943	15.37%	2,576,084	25.50%	5,313,552	52.60%	0	0.00%	0	0.00%
<b>Sleep disorder</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,149,701	182,758	15.90%	394,538	34.32%	290,247	25.25%	282,157	24.54%	0	0.00%	0	0.00%
30 to 34	1,546,562	227,412	14.70%	387,513	25.06%	422,732	27.33%	508,904	32.91%	0	0.00%	0	0.00%
35 to 39	1,921,406	490,570	25.53%	585,494	30.47%	553,711	28.82%	261,957	13.63%	29,674	1.54%	0	0.00%
40 to 44	1,615,309	498,735	30.88%	321,396	19.90%	590,666	36.57%	163,501	10.12%	41,011	2.54%	0	0.00%
45 to 49	3,088,647	791,959	25.64%	374,343	12.12%	914,548	29.61%	1,007,796	32.63%	0	0.00%	0	0.00%
50 to 54	3,764,397	993,798	26.40%	1,036,860	27.54%	940,953	25.00%	792,787	21.06%	0	0.00%	0	0.00%
55 or older	1,387,354	282,654	20.37%	405,756	29.25%	333,803	24.06%	365,140	26.32%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	9,377,790	557,675	5.95%	1,421,684	15.16%	2,480,966	26.46%	4,917,465	52.44%	0	0.00%	0	0.00%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	482,168	90,416	18.75%	113,728	23.59%	125,094	25.94%	152,930	31.72%	0	0.00%	0	0.00%
None	2,498,228	156,990	6.28%	535,830	21.45%	794,348	31.80%	1,011,059	40.47%	0	0.00%	0	0.00%
1	2,331,034	122,457	5.25%	479,207	20.56%	773,227	33.17%	956,143	41.02%	0	0.00%	0	0.00%
2	3,932,623	505,278	12.85%	507,219	12.90%	1,295,652	32.95%	1,594,800	40.55%	29,674	0.75%	0	0.00%
More than 2	14,681,699	3,174,304	21.62%	3,307,530	22.53%	3,545,130	24.15%	4,596,680	31.31%	58,054	0.40%	0	0.00%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	601,966	98,042	16.29%	113,728	18.89%	125,094	20.78%	265,102	44.04%	0	0.00%	0	0.00%
29 years old or younger	15,812,185	3,384,828	21.41%	3,351,216	21.19%	4,143,186	26.20%	4,874,901	30.83%	58,054	0.37%	0	0.00%
30 - 40 years old	800,094	81,195	10.15%	151,943	18.99%	408,559	51.06%	158,397	19.80%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,498,228	156,990	6.28%	535,830	21.45%	794,348	31.80%	1,011,059	40.47%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	3,293,674	271,350	8.24%	536,990	16.30%	684,905	20.79%	1,770,755	53.76%	29,674	0.90%	0	0.00%
No pregnancies resulting in live births	903,674	57,039	6.31%	237,878	26.32%	377,359	41.76%	231,398	25.61%	0	0.00%	0	0.00%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,331,696	461,482	19.79%	337,613	14.48%	500,427	21.46%	1,002,500	42.99%	29,674	1.27%	0	0.00%
29 years old or younger	9,114,520	1,731,194	18.99%	1,987,929	21.81%	2,096,297	23.00%	3,299,101	36.20%	0	0.00%	0	0.00%
30 - 40 years old	7,943,374	1,404,960	17.69%	1,568,717	19.75%	2,489,873	31.35%	2,421,770	30.49%	58,054	0.73%	0	0.00%
41 years or older	1,134,260	237,780	20.96%	275,548	24.29%	275,148	24.26%	345,783	30.49%	0	0.00%	0	0.00%
N/A - Never pregnant	2,498,228	156,990	6.28%	535,830	21.45%	794,348	31.80%	1,011,059	40.47%	0	0.00%	0	0.00%
No pregnancies resulting in live births	903,674	57,039	6.31%	237,878	26.32%	377,359	41.76%	231,398	25.61%	0	0.00%	0	0.00%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	7,531,384	307,831	4.09%	928,173	12.32%	2,131,199	28.30%	4,164,181	55.29%	0	0.00%	0	0.00%
Yes	8,240,673	2,138,509	25.95%	2,076,538	25.20%	2,160,092	26.21%	1,818,817	22.07%	46,717	0.57%	0	0.00%
No	8,153,694	1,603,105	19.66%	1,938,803	23.78%	2,242,160	27.50%	2,328,614	28.56%	41,011	0.50%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	7,564,128	307,831	4.07%	928,173	12.27%	2,163,942	28.61%	4,164,181	55.05%	0	0.00%	0	0.00%
35 years old or younger	3,024,492	624,515	20.65%	836,804	27.67%	745,276	24.64%	817,898	27.04%	0	0.00%	0	0.00%
36 - 50 years old	3,911,183	1,149,224	29.38%	819,057	20.94%	1,097,666	28.06%	798,518	20.42%	46,717	1.19%	0	0.00%
51 years or older	1,144,392	364,770	31.87%	370,059	32.34%	240,998	21.06%	168,565	14.73%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	8,153,694	1,603,105	19.66%	1,938,803	23.78%	2,242,160	27.50%	2,328,614	28.56%	41,011	0.50%	0	0.00%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,493,988	226,549	15.16%	286,321	19.16%	333,221	22.30%	647,897	43.37%	0	0.00%	0	0.00%
Yes	8,240,673	2,138,509	25.95%	2,076,538	25.20%	2,160,092	26.21%	1,818,817	22.07%	46,717	0.57%	0	0.00%
No	14,191,090	1,684,387	11.87%	2,580,655	18.19%	4,040,138	28.47%	5,844,898	41.19%	41,011	0.29%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	1,456,241	127,219	8.74%	346,788	23.81%	402,862	27.66%	579,372	39.79%	0	0.00%	0	0.00%
30-39	3,216,701	409,152	12.72%	586,788	18.24%	861,197	26.77%	1,359,564	42.27%	0	0.00%	0	0.00%
40-49	4,170,880	396,675	9.51%	691,508	16.58%	843,551	20.22%	2,239,146	53.69%	0	0.00%	0	0.00%
50-59	5,527,419	568,771	10.29%	1,182,892	21.40%	1,287,109	23.29%	2,488,647	45.02%	0	0.00%	0	0.00%
60-64	1,283,925	277,830	21.64%	317,784	24.75%	286,907	22.35%	401,404	31.26%	0	0.00%	0	0.00%
65-69	2,590,075	361,246	13.95%	729,006	28.15%	484,158	18.69%	1,015,664	39.21%	0	0.00%	0	0.00%
70-74	2,088,104	532,076	25.48%	479,051	22.94%	671,522	32.16%	405,455	19.42%	0	0.00%	0	0.00%
75-79	2,415,697	636,147	26.33%	673,241	27.87%	472,798	19.57%	633,511	26.22%	0	0.00%	0	0.00%
80-84	1,087,353	351,926	32.37%	337,716	31.06%	267,679	24.62%	130,032	11.96%	0	0.00%	0	0.00%
85+	1,051,109	370,102	35.21%	304,513	28.97%	220,832	21.01%	155,662	14.81%	0	0.00%	0	0.00%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,422,888	204,482	14.37%	328,300	23.07%	378,064	26.57%	512,043	35.99%	0	0.00%	0	0.00%
Other Hispanic	1,020,738	21,114	2.07%	69,122	6.77%	354,915	34.77%	575,588	56.39%	0	0.00%	0	0.00%
Non-Hispanic White	18,440,395	3,030,757	16.44%	4,154,767	22.53%	3,929,034	21.31%	7,325,836	39.73%	0	0.00%	0	0.00%
Non-Hispanic Black	3,321,231	518,273	15.60%	811,462	24.43%	1,011,927	30.47%	979,568	29.49%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	809,297	256,519	31.70%	285,635	35.29%	124,676	15.41%	142,467	17.60%	0	0.00%	0	0.00%
<b>Education</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	912,462	164,382	18.02%	150,755	16.52%	307,513	33.70%	289,812	31.76%	0	0.00%	0	0.00%
Less Than 9th Grade	2,200,338	521,670	23.71%	805,388	36.60%	378,143	17.19%	495,136	22.50%	0	0.00%	0	0.00%
9-11th Grade	3,463,925	838,070	24.19%	942,745	27.22%	892,891	25.78%	790,219	22.81%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	18,399,215	2,471,771	13.43%	3,747,041	20.37%	4,220,067	22.94%	7,960,335	43.26%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Marital status</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	12,323,620	1,571,915	12.76%	2,501,007	20.29%	3,291,943	26.71%	4,958,754	40.24%	0	0.00%	0	0.00%
Widowed	4,599,609	1,237,637	26.91%	1,250,311	27.18%	1,066,871	23.19%	1,044,790	22.71%	0	0.00%	0	0.00%
Divorced	3,746,586	677,474	18.08%	764,764	20.41%	533,553	14.24%	1,770,796	47.26%	0	0.00%	0	0.00%
Separated	419,989	159,387	37.95%	22,284	5.31%	124,804	29.72%	113,513	27.03%	0	0.00%	0	0.00%
Never Married	2,647,160	238,548	9.01%	695,982	26.29%	534,472	20.19%	1,178,157	44.51%	0	0.00%	0	0.00%
Living with Partner	1,192,059	113,027	9.48%	414,937	34.81%	246,973	20.72%	417,122	34.99%	0	0.00%	0	0.00%
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,169,561	238,134	20.36%	491,702	42.04%	215,476	18.42%	224,249	19.17%	0	0.00%	0	0.00%
Poverty income ratio less than 1	4,158,086	660,718	15.89%	911,831	21.93%	911,529	21.92%	1,674,008	40.26%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	19,686,902	3,132,292	15.91%	4,245,754	21.57%	4,671,610	23.73%	7,637,245	38.79%	0	0.00%	0	0.00%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	431,430	126,093	29.23%	229,095	53.10%	70,102	16.25%	6,140	1.42%	0	0.00%	0	0.00%
Normal: less than 25	6,694,721	986,046	14.73%	1,509,502	22.55%	1,546,860	23.11%	2,652,314	39.62%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	7,688,386	1,217,847	15.84%	1,494,558	19.44%	1,549,987	20.16%	3,425,993	44.56%	0	0.00%	0	0.00%
Obese: 30 or higher	10,200,012	1,701,159	16.68%	2,416,132	23.69%	2,631,666	25.80%	3,451,054	33.83%	0	0.00%	0	0.00%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	628,980	108,805	17.30%	263,351	41.87%	70,182	11.16%	186,641	29.67%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	17,412,020	2,463,907	14.15%	3,225,675	18.53%	3,893,271	22.36%	7,829,167	44.96%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	4,725,769	820,403	17.36%	1,346,010	28.48%	1,495,709	31.65%	1,063,647	22.51%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	2,247,780	638,029	28.38%	814,251	36.22%	339,453	15.10%	456,046	20.29%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Diabetes</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,455,291	778,350	22.53%	1,384,459	40.07%	666,465	19.29%	626,018	18.12%	0	0.00%	0	0.00%
No	21,559,257	3,252,795	15.09%	4,264,828	19.78%	5,132,151	23.80%	8,909,484	41.33%	0	0.00%	0	0.00%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	12,859,201	2,227,464	17.32%	2,701,452	21.01%	3,282,801	25.53%	4,647,484	36.14%	0	0.00%	0	0.00%
No	12,155,348	1,803,681	14.84%	2,947,835	24.25%	2,515,814	20.70%	4,888,017	40.21%	0	0.00%	0	0.00%
<b>High blood pressure</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	10,715,400	2,119,245	19.78%	2,938,725	27.43%	2,764,838	25.80%	2,892,591	26.99%	0	0.00%	0	0.00%
No	14,082,853	1,911,899	13.58%	2,680,733	19.04%	2,957,203	21.00%	6,533,018	46.39%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,460,263	1,593,665	29.19%	1,647,550	30.17%	1,287,297	23.58%	931,751	17.06%	0	0.00%	0	0.00%
Mobility limitations	4,852,008	818,448	16.87%	1,540,705	31.75%	1,217,238	25.09%	1,275,617	26.29%	0	0.00%	0	0.00%
No mobility limitations	5,878,766	973,403	16.56%	1,310,682	22.30%	1,830,850	31.14%	1,763,831	30.00%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	8,823,512	645,629	7.32%	1,150,350	13.04%	1,463,230	16.58%	5,564,302	63.06%	0	0.00%	0	0.00%
<b>Sleep disorder</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	633,447	260,536	41.13%	122,448	19.33%	212,130	33.49%	38,333	6.05%	0	0.00%	0	0.00%
Younger than 30	1,102,451	227,530	20.64%	201,484	18.28%	333,626	30.26%	339,810	30.82%	0	0.00%	0	0.00%
30 to 34	2,166,179	259,028	11.96%	599,065	27.66%	270,637	12.49%	1,037,448	47.89%	0	0.00%	0	0.00%
35 to 39	2,300,079	746,059	32.44%	697,491	30.32%	490,054	21.31%	366,475	15.93%	0	0.00%	0	0.00%
40 to 44	2,136,702	308,228	14.43%	575,686	26.94%	472,578	22.12%	780,210	36.51%	0	0.00%	0	0.00%
45 to 49	3,579,247	441,682	12.34%	951,364	26.58%	922,661	25.78%	1,263,540	35.30%	0	0.00%	0	0.00%
50 to 54	2,980,449	726,406	24.37%	591,502	19.85%	810,751	27.20%	851,790	28.58%	0	0.00%	0	0.00%
55 or older	2,030,084	265,156	13.06%	572,552	28.20%	464,000	22.86%	728,376	35.88%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	8,031,301	779,330	9.70%	1,300,276	16.19%	1,822,179	22.69%	4,129,517	51.42%	0	0.00%	0	0.00%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	626,612	260,536	41.58%	97,314	15.53%	140,325	22.39%	128,438	20.50%	0	0.00%	0	0.00%
None	2,939,167	493,252	16.78%	657,439	22.37%	757,736	25.78%	1,030,740	35.07%	0	0.00%	0	0.00%
1	2,664,303	210,348	7.90%	589,789	22.14%	751,207	28.20%	1,112,959	41.77%	0	0.00%	0	0.00%
2	4,860,387	402,035	8.27%	1,078,841	22.20%	1,178,624	24.25%	2,200,886	45.28%	0	0.00%	0	0.00%
More than 2	13,924,079	2,664,974	19.14%	3,225,904	23.17%	2,970,723	21.34%	5,062,478	36.36%	0	0.00%	0	0.00%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	550,922	260,536	47.29%	97,314	17.66%	148,133	26.89%	44,940	8.16%	0	0.00%	0	0.00%
29 years old or younger	16,012,220	2,766,856	17.28%	3,709,173	23.16%	3,380,182	21.11%	6,156,010	38.45%	0	0.00%	0	0.00%
30 - 40 years old	839,379	65,332	7.78%	178,572	21.27%	202,705	24.15%	392,769	46.79%	0	0.00%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,939,167	493,252	16.78%	657,439	22.37%	757,736	25.78%	1,030,740	35.07%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	3,876,784	421,596	10.87%	851,553	21.97%	1,120,699	28.91%	1,482,935	38.25%	0	0.00%	0	0.00%
No pregnancies resulting in live births	733,146	23,572	3.22%	155,237	21.17%	180,924	24.68%	373,413	50.93%	0	0.00%	0	0.00%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,055,229	468,711	22.81%	403,015	19.61%	687,308	33.44%	496,196	24.14%	0	0.00%	0	0.00%
29 years old or younger	12,007,836	1,795,201	14.95%	3,057,038	25.46%	2,495,569	20.78%	4,660,027	38.81%	0	0.00%	0	0.00%
30 - 40 years old	6,573,739	1,151,548	17.52%	1,248,934	19.00%	1,529,743	23.27%	2,643,514	40.21%	0	0.00%	0	0.00%
41 years or older	675,009	92,052	13.64%	104,011	15.41%	147,335	21.83%	331,611	49.13%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,939,167	493,252	16.78%	657,439	22.37%	757,736	25.78%	1,030,740	35.07%	0	0.00%	0	0.00%
No pregnancies resulting in live births	733,146	23,572	3.22%	155,237	21.17%	180,924	24.68%	373,413	50.93%	0	0.00%	0	0.00%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,629,010	755,679	11.40%	900,320	13.58%	1,712,398	25.83%	3,260,613	49.19%	0	0.00%	0	0.00%
Yes	8,638,932	1,767,511	20.46%	2,013,798	23.31%	1,960,157	22.69%	2,897,466	33.54%	0	0.00%	0	0.00%
No	9,683,313	1,507,954	15.57%	2,735,169	28.25%	2,062,767	21.30%	3,377,423	34.88%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	23,925,751	4,049,445	16.93%	4,943,515	20.66%	6,533,452	27.31%	8,311,612	34.74%	87,728	0.37%	0	0.00%
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	6,692,303	755,679	11.29%	900,320	13.45%	1,775,692	26.53%	3,260,613	48.72%	0	0.00%	0	0.00%
35 years old or younger	3,606,573	684,257	18.97%	904,079	25.07%	574,591	15.93%	1,443,647	40.03%	0	0.00%	0	0.00%
36 - 50 years old	3,491,696	675,308	19.34%	625,133	17.90%	1,001,731	28.69%	1,189,524	34.07%	0	0.00%	0	0.00%
51 years or older	1,455,269	407,946	28.03%	399,193	27.43%	383,835	26.38%	264,295	18.16%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	9,683,313	1,507,954	15.57%	2,735,169	28.25%	2,062,767	21.30%	3,377,423	34.88%	0	0.00%	0	0.00%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2003-2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,014,549	4,031,145	16.12%	5,649,287	22.58%	5,798,615	23.18%	9,535,501	38.12%	0	0.00%	0	0.00%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	536,508	260,536	48.56%	97,314	18.14%	140,325	26.16%	38,333	7.14%	0	0.00%	0	0.00%
Yes	8,638,932	1,767,511	20.46%	2,013,798	23.31%	1,960,157	22.69%	2,897,466	33.54%	0	0.00%	0	0.00%
No	15,775,816	2,003,098	12.70%	3,538,175	22.43%	3,634,841	23.04%	6,599,702	41.83%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	1,590,931	51,417	3.23%	49,889	3.14%	390,566	24.55%	1,044,732	65.67%	0	0.00%	54,327	3.41%
30-39	3,254,841	174,857	5.37%	408,537	12.55%	925,482	28.43%	1,745,965	53.64%	0	0.00%	0	0.00%
40-49	4,526,778	582,984	12.88%	269,964	5.96%	1,669,758	36.89%	2,004,072	44.27%	0	0.00%	0	0.00%
50-59	5,282,835	274,036	5.19%	1,008,235	19.09%	1,404,127	26.58%	2,596,437	49.15%	0	0.00%	0	0.00%
60-64	2,330,182	178,276	7.65%	236,198	10.14%	694,725	29.81%	1,156,209	49.62%	64,774	2.78%	0	0.00%
65-69	2,237,833	118,900	5.31%	527,510	23.57%	917,555	41.00%	673,868	30.11%	0	0.00%	0	0.00%
70-74	2,329,200	205,129	8.81%	555,465	23.85%	666,455	28.61%	902,150	38.73%	0	0.00%	0	0.00%
75-79	1,408,061	107,194	7.61%	230,219	16.35%	554,435	39.38%	516,213	36.66%	0	0.00%	0	0.00%
80-84	1,261,821	362,359	28.72%	223,067	17.68%	199,390	15.80%	428,940	33.99%	48,066	3.81%	0	0.00%
85+	940,262	273,383	29.08%	219,407	23.33%	265,050	28.19%	182,421	19.40%	0	0.00%	0	0.00%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,628,304	68,706	4.22%	258,855	15.90%	553,774	34.01%	746,968	45.87%	0	0.00%	0	0.00%
Other Hispanic	879,742	211,073	23.99%	21,615	2.46%	307,306	34.93%	339,748	38.62%	0	0.00%	0	0.00%
Non-Hispanic White	18,283,765	1,744,892	9.54%	2,663,137	14.57%	5,491,887	30.04%	8,271,010	45.24%	112,839	0.62%	0	0.00%
Non-Hispanic Black	3,525,787	273,774	7.76%	571,993	16.22%	992,258	28.14%	1,687,760	47.87%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	903,325	30,091	3.33%	237,540	26.30%	342,318	37.90%	239,048	26.46%	0	0.00%	54,327	6.01%
<b>Education</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	604,008	104,723	17.34%	97,703	16.18%	188,913	31.28%	212,669	35.21%	0	0.00%	0	0.00%
Less Than 9th Grade	1,661,873	335,097	20.16%	332,373	20.00%	478,600	28.80%	515,802	31.04%	0	0.00%	0	0.00%
9-11th Grade	3,465,061	365,639	10.55%	420,678	12.14%	1,206,066	34.81%	1,472,678	42.50%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	19,457,204	1,523,077	7.83%	2,902,386	14.92%	5,781,189	29.71%	9,083,385	46.68%	112,839	0.58%	54,327	0.28%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Marital status</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	14,167,387	1,301,676	9.19%	2,203,970	15.56%	4,734,397	33.42%	5,873,017	41.45%	0	0.00%	54,327	0.38%
Widowed	3,356,052	560,898	16.71%	677,546	20.19%	943,529	28.11%	1,126,014	33.55%	48,066	1.43%	0	0.00%
Divorced	3,357,375	217,987	6.49%	352,515	10.50%	963,714	28.70%	1,823,159	54.30%	0	0.00%	0	0.00%
Separated	877,219	78,678	8.97%	101,684	11.59%	319,167	36.38%	377,690	43.06%	0	0.00%	0	0.00%
Never Married	2,205,102	106,087	4.81%	132,170	5.99%	604,879	27.43%	1,361,966	61.76%	0	0.00%	0	0.00%
Living with Partner	1,131,141	63,211	5.59%	285,256	25.22%	121,857	10.77%	596,043	52.69%	64,774	5.73%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,390,005	0	0.00%	130,529	9.39%	502,691	36.16%	756,784	54.44%	0	0.00%	0	0.00%
Poverty income ratio less than 1	3,182,392	357,082	11.22%	603,087	18.95%	1,031,725	32.42%	1,136,170	35.70%	0	0.00%	54,327	1.71%
Poverty income ratio 1 or greater	20,648,525	1,971,454	9.55%	3,019,525	14.62%	6,153,127	29.80%	9,391,580	45.48%	112,839	0.55%	0	0.00%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	491,432	199,171	40.53%	66,223	13.48%	212,472	43.24%	13,566	2.76%	0	0.00%	0	0.00%
Normal: less than 25	6,846,555	285,338	4.17%	1,117,840	16.33%	1,359,713	19.86%	3,970,826	58.00%	112,839	1.65%	0	0.00%
Overweight: 25.0 - 29.9	6,576,612	650,587	9.89%	749,148	11.39%	2,252,283	34.25%	2,870,267	43.64%	0	0.00%	54,327	0.83%
Obese: 30 or higher	11,306,323	1,193,441	10.56%	1,819,930	16.10%	3,863,076	34.17%	4,429,876	39.18%	0	0.00%	0	0.00%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	667,970	67,292	10.07%	41,921	6.28%	114,696	17.17%	379,287	56.78%	64,774	9.70%	0	0.00%
Glycohemoglobin < 5.7%	17,400,028	1,555,994	8.94%	2,491,140	14.32%	5,025,597	28.88%	8,272,970	47.55%	0	0.00%	54,327	0.31%
Glycohemoglobin 5.7% - 6.4%	5,047,775	587,059	11.63%	877,480	17.38%	1,799,172	35.64%	1,784,065	35.34%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	2,105,149	118,191	5.61%	342,600	16.27%	748,079	35.54%	848,213	40.29%	48,066	2.28%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Diabetes</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,250,395	298,540	9.18%	594,961	18.30%	1,289,374	39.67%	1,019,454	31.36%	48,066	1.48%	0	0.00%
No	21,970,527	2,029,996	9.24%	3,158,180	14.37%	6,398,169	29.12%	10,265,081	46.72%	64,774	0.29%	54,327	0.25%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	10,359,802	655,774	6.33%	1,683,107	16.25%	2,963,590	28.61%	4,938,229	47.67%	64,774	0.63%	54,327	0.52%
No	14,852,634	1,672,762	11.26%	2,070,034	13.94%	4,723,953	31.81%	6,337,819	42.67%	48,066	0.32%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	10,776,742	1,185,355	11.00%	1,589,507	14.75%	3,551,775	32.96%	4,450,105	41.29%	0	0.00%	0	0.00%
No	14,356,119	1,111,448	7.74%	2,163,634	15.07%	4,135,768	28.81%	6,778,102	47.21%	112,839	0.79%	54,327	0.38%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	4,973,568	784,641	15.78%	765,588	15.39%	1,960,926	39.43%	1,414,348	28.44%	48,066	0.97%	0	0.00%
Mobility limitations	3,829,182	522,256	13.64%	497,427	12.99%	1,432,974	37.42%	1,376,525	35.95%	0	0.00%	0	0.00%
No mobility limitations	6,568,299	547,734	8.34%	1,426,420	21.72%	1,725,656	26.27%	2,803,715	42.69%	64,774	0.99%	0	0.00%
N/A - Respondent not asked mobility limitation questions	9,849,874	473,906	4.81%	1,063,705	10.80%	2,567,988	26.07%	5,689,947	57.77%	0	0.00%	54,327	0.55%
<b>Sleep disorder</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,532,603	363,772	14.36%	448,084	17.69%	728,066	28.75%	992,682	39.20%	0	0.00%	0	0.00%
No	22,688,319	1,964,764	8.66%	3,305,057	14.57%	6,959,478	30.67%	10,291,853	45.36%	112,839	0.50%	54,327	0.24%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	3,512,561	446,817	12.72%	675,081	19.22%	994,829	28.32%	1,395,834	39.74%	0	0.00%	0	0.00%
Once a day	2,391,390	169,722	7.10%	345,015	14.43%	617,325	25.81%	1,259,328	52.66%	0	0.00%	0	0.00%
2 or more time a week	2,694,274	144,848	5.38%	298,996	11.10%	997,091	37.01%	1,253,338	46.52%	0	0.00%	0	0.00%
Once a week	1,733,866	207,951	11.99%	379,208	21.87%	434,598	25.07%	712,109	41.07%	0	0.00%	0	0.00%
1-3 times a month	4,212,110	316,239	7.51%	863,587	20.50%	1,537,013	36.49%	1,430,498	33.96%	64,774	1.54%	0	0.00%
Never	10,350,050	1,011,226	9.77%	1,146,654	11.08%	3,020,875	29.19%	5,171,294	49.96%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	472,461	127,763	27.04%	88,557	18.74%	147,480	31.22%	108,661	23.00%	0	0.00%	0	0.00%
Never	24,157,788	2,061,377	8.53%	3,516,844	14.56%	7,401,054	30.64%	11,113,740	46.00%	64,774	0.27%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,324,106	238,535	18.01%	249,026	18.81%	251,549	19.00%	584,997	44.18%	0	0.00%	0	0.00%
30 to 34	1,348,019	185,678	13.77%	69,692	5.17%	540,151	40.07%	552,498	40.99%	0	0.00%	0	0.00%
35 to 39	2,097,662	266,879	12.72%	506,384	24.14%	457,093	21.79%	867,306	41.35%	0	0.00%	0	0.00%
40 to 44	2,901,226	251,834	8.68%	572,578	19.74%	1,019,581	35.14%	1,057,233	36.44%	0	0.00%	0	0.00%
45 to 49	4,183,891	399,480	9.55%	390,459	9.33%	1,637,628	39.14%	1,756,324	41.98%	0	0.00%	0	0.00%
50 to 54	3,495,074	313,958	8.98%	796,670	22.79%	878,580	25.14%	1,441,092	41.23%	64,774	1.85%	0	0.00%
55 or older	1,010,289	73,331	7.26%	193,225	19.13%	336,398	33.30%	359,270	35.56%	48,066	4.76%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	8,704,440	535,102	6.15%	975,107	11.20%	2,474,088	28.42%	4,665,816	53.60%	0	0.00%	54,327	0.62%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
None	1,773,752	45,315	2.55%	104,424	5.89%	455,264	25.67%	1,120,684	63.18%	48,066	2.71%	0	0.00%
1	3,109,925	173,090	5.57%	368,029	11.83%	524,325	16.86%	2,044,481	65.74%	0	0.00%	0	0.00%
2	5,878,129	578,549	9.84%	923,971	15.72%	1,752,563	29.81%	2,568,720	43.70%	0	0.00%	54,327	0.92%
More than 2	14,459,116	1,531,582	10.59%	2,356,717	16.30%	4,955,393	34.27%	5,550,651	38.39%	64,774	0.45%	0	0.00%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	5,294,183	312,147	5.90%	649,609	12.27%	1,556,226	29.40%	2,728,134	51.53%	48,066	0.91%	0	0.00%
1	4,287,844	171,398	4.00%	627,126	14.63%	851,655	19.86%	2,637,665	61.51%	0	0.00%	0	0.00%
2	5,683,911	639,679	11.25%	749,569	13.19%	2,055,562	36.16%	2,184,774	38.44%	0	0.00%	54,327	0.96%
More than 2	9,910,261	1,205,312	12.16%	1,726,837	17.42%	3,224,099	32.53%	3,689,239	37.23%	64,774	0.65%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	8,584,518	801,541	9.34%	1,638,234	19.08%	2,190,475	25.52%	3,954,269	46.06%	0	0.00%	0	0.00%
1	1,829,322	176,831	9.67%	240,940	13.17%	639,280	34.95%	772,271	42.22%	0	0.00%	0	0.00%
2	1,397,266	118,114	8.45%	228,155	16.33%	575,127	41.16%	475,871	34.06%	0	0.00%	0	0.00%
More than 2	488,819	25,289	5.17%	48,446	9.91%	227,349	46.51%	187,735	38.41%	0	0.00%	0	0.00%
N/A - Vaginal delivery/deliveries only	12,876,274	1,206,763	9.37%	1,597,367	12.41%	4,055,312	31.49%	5,849,666	45.43%	112,839	0.88%	54,327	0.42%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	646,120	38,208	5.91%	128,928	19.95%	49,851	7.72%	429,134	66.42%	0	0.00%	0	0.00%
29 years old or younger	16,623,518	1,824,184	10.97%	2,723,340	16.38%	5,705,239	34.32%	6,251,654	37.61%	64,774	0.39%	54,327	0.33%
30 - 40 years old	780,039	50,514	6.48%	17,797	2.28%	390,688	50.09%	321,040	41.16%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,773,752	45,315	2.55%	104,424	5.89%	455,264	25.67%	1,120,684	63.18%	48,066	2.71%	0	0.00%
Only 1 live birth - see age at last live birth	4,188,366	293,040	7.00%	678,628	16.20%	751,288	17.94%	2,465,410	58.86%	0	0.00%	0	0.00%
No pregnancies resulting in live births	1,067,171	66,013	6.19%	56,987	5.34%	280,976	26.33%	663,195	62.15%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	646,120	38,208	5.91%	128,928	19.95%	49,851	7.72%	429,134	66.42%	0	0.00%	0	0.00%
29 years old or younger	11,919,979	1,188,248	9.97%	1,731,658	14.53%	3,641,019	30.55%	5,239,953	43.96%	64,774	0.54%	54,327	0.46%
30 - 40 years old	9,123,265	904,691	9.92%	1,611,427	17.66%	2,940,871	32.23%	3,666,276	40.19%	0	0.00%	0	0.00%
41 years or older	451,924	35,126	7.77%	95,068	21.04%	195,249	43.20%	126,482	27.99%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,773,752	45,315	2.55%	104,424	5.89%	455,264	25.67%	1,120,684	63.18%	48,066	2.71%	0	0.00%
No pregnancies resulting in live births	1,067,171	66,013	6.19%	56,987	5.34%	280,976	26.33%	663,195	62.15%	0	0.00%	0	0.00%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	7,570,700	511,016	6.75%	922,253	12.18%	2,026,997	26.77%	4,056,106	53.58%	0	0.00%	54,327	0.72%
Yes	8,113,279	1,153,294	14.21%	1,446,488	17.83%	2,807,940	34.61%	2,705,557	33.35%	0	0.00%	0	0.00%
No	9,503,575	664,226	6.99%	1,351,032	14.22%	2,852,606	30.02%	4,522,871	47.59%	112,839	1.19%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	7,604,068	511,016	6.72%	955,621	12.57%	2,026,997	26.66%	4,056,106	53.34%	0	0.00%	54,327	0.71%
35 years old or younger	2,718,788	466,699	17.17%	462,448	17.01%	750,876	27.62%	1,038,765	38.21%	0	0.00%	0	0.00%
36 - 50 years old	4,555,075	503,335	11.05%	738,421	16.21%	1,854,148	40.71%	1,459,171	32.03%	0	0.00%	0	0.00%
51 years or older	602,808	54,551	9.05%	169,726	28.16%	170,910	28.35%	207,622	34.44%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	9,503,575	664,226	6.99%	1,351,032	14.22%	2,852,606	30.02%	4,522,871	47.59%	112,839	1.19%	0	0.00%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2005-2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,220,922	2,328,536	9.23%	3,753,141	14.88%	7,687,543	30.48%	11,284,535	44.74%	112,839	0.45%	54,327	0.22%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	8,113,279	1,153,294	14.21%	1,446,488	17.83%	2,807,940	34.61%	2,705,557	33.35%	0	0.00%	0	0.00%
No	17,074,275	1,175,242	6.88%	2,273,285	13.31%	4,879,603	28.58%	8,578,978	50.25%	112,839	0.66%	54,327	0.32%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	1,070,655	57,669	5.39%	74,087	6.92%	399,705	37.33%	539,193	50.36%	0	0.00%	0	0.00%
30-39	2,291,102	174,834	7.63%	456,303	19.92%	566,659	24.73%	1,080,097	47.14%	13,209	0.58%	0	0.00%
40-49	5,124,808	364,407	7.11%	365,538	7.13%	1,691,752	33.01%	2,703,112	52.75%	0	0.00%	0	0.00%
50-59	6,035,135	409,241	6.78%	799,327	13.24%	1,949,214	32.30%	2,877,354	47.68%	0	0.00%	0	0.00%
60-64	2,365,499	142,995	6.05%	346,245	14.64%	822,976	34.79%	1,053,284	44.53%	0	0.00%	0	0.00%
65-69	2,249,997	179,744	7.99%	382,682	17.01%	845,592	37.58%	841,979	37.42%	0	0.00%	0	0.00%
70-74	2,543,230	151,675	5.96%	224,435	8.82%	1,247,676	49.06%	912,865	35.89%	6,578	0.26%	0	0.00%
75-79	1,844,535	181,477	9.84%	402,313	21.81%	612,716	33.22%	648,029	35.13%	0	0.00%	0	0.00%
80+	2,594,583	406,878	15.68%	447,422	17.24%	900,723	34.72%	824,260	31.77%	15,300	0.59%	0	0.00%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,553,678	146,864	9.45%	141,272	9.09%	575,778	37.06%	683,186	43.97%	6,578	0.42%	0	0.00%
Other Hispanic	1,301,346	157,593	12.11%	244,087	18.76%	391,071	30.05%	508,596	39.08%	0	0.00%	0	0.00%
Non-Hispanic White	18,979,877	1,522,320	8.02%	2,534,637	13.35%	6,678,339	35.19%	8,244,582	43.44%	0	0.00%	0	0.00%
Non-Hispanic Black	3,610,806	199,556	5.53%	589,459	16.32%	1,060,751	29.38%	1,732,530	47.98%	28,510	0.79%	0	0.00%
Other Race - Including Multi-Racial	805,075	42,586	5.29%	24,947	3.10%	331,074	41.12%	406,468	50.49%	0	0.00%	0	0.00%
<b>Education</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	780,652	9,596	1.23%	74,990	9.61%	335,486	42.98%	360,580	46.19%	0	0.00%	0	0.00%
Less Than 9th Grade	2,242,619	274,889	12.26%	524,214	23.38%	640,657	28.57%	796,281	35.51%	6,578	0.29%	0	0.00%
9-11th Grade	3,881,059	314,872	8.11%	747,988	19.27%	1,703,764	43.90%	1,099,136	28.32%	15,300	0.39%	0	0.00%
High School Grad/GED or Equivalent	19,334,836	1,469,563	7.60%	2,175,593	11.25%	6,357,106	32.88%	9,319,365	48.20%	13,209	0.07%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Marital status</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	14,074,127	989,427	7.03%	1,781,166	12.66%	4,921,080	34.97%	6,375,876	45.30%	6,578	0.05%	0	0.00%
Widowed	3,526,660	466,931	13.24%	486,903	13.81%	1,210,693	34.33%	1,346,834	38.19%	15,300	0.43%	0	0.00%
Divorced	4,100,623	375,873	9.17%	736,335	17.96%	1,393,368	33.98%	1,595,047	38.90%	0	0.00%	0	0.00%
Separated	606,331	57,107	9.42%	113,877	18.78%	162,123	26.74%	273,224	45.06%	0	0.00%	0	0.00%
Never Married	2,474,011	115,346	4.66%	326,504	13.20%	817,455	33.04%	1,201,496	48.56%	13,209	0.53%	0	0.00%
Living with Partner	1,208,744	64,236	5.31%	48,466	4.01%	488,727	40.43%	607,316	50.24%	0	0.00%	0	0.00%
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,785,020	233,274	13.07%	177,562	9.95%	518,773	29.06%	848,832	47.55%	6,578	0.37%	0	0.00%
Poverty income ratio less than 1	3,819,925	311,457	8.15%	537,956	14.08%	1,059,086	27.73%	1,911,426	50.04%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	20,645,837	1,524,188	7.38%	2,818,882	13.65%	7,459,153	36.13%	8,815,105	42.70%	28,510	0.14%	0	0.00%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	426,688	70,259	16.47%	54,408	12.75%	85,520	20.04%	203,291	47.64%	13,209	3.10%	0	0.00%
Normal: less than 25	7,329,434	443,702	6.05%	1,254,496	17.12%	2,112,442	28.82%	3,512,216	47.92%	6,578	0.09%	0	0.00%
Overweight: 25.0 - 29.9	7,888,995	642,051	8.14%	827,589	10.49%	3,044,879	38.60%	3,374,476	42.77%	0	0.00%	0	0.00%
Obese: 30 or higher	10,605,665	912,908	8.61%	1,397,908	13.18%	3,794,171	35.77%	4,485,378	42.29%	15,300	0.14%	0	0.00%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	781,126	39,042	5.00%	71,249	9.12%	353,872	45.30%	288,454	36.93%	28,510	3.65%	0	0.00%
Glycohemoglobin < 5.7%	13,407,599	856,033	6.38%	1,538,401	11.47%	4,652,099	34.70%	6,361,066	47.44%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	9,014,889	716,166	7.94%	1,494,524	16.58%	2,803,924	31.10%	3,993,697	44.30%	6,578	0.07%	0	0.00%
Glycohemoglobin 6.5% or higher	3,047,167	457,678	15.02%	430,227	14.12%	1,227,116	40.27%	932,145	30.59%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Diabetes</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,408,790	658,550	14.94%	725,708	16.46%	1,778,895	40.35%	1,232,428	27.95%	13,209	0.30%	0	0.00%
No	21,841,991	1,410,370	6.46%	2,808,693	12.86%	7,258,117	33.23%	10,342,934	47.35%	21,878	0.10%	0	0.00%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	12,368,068	822,239	6.65%	1,740,063	14.07%	4,213,133	34.06%	5,557,545	44.93%	35,088	0.28%	0	0.00%
No	13,882,714	1,246,680	8.98%	1,794,338	12.92%	4,823,879	34.75%	6,017,817	43.35%	0	0.00%	0	0.00%
<b>High blood pressure</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	12,649,611	971,375	7.68%	1,995,541	15.78%	4,647,142	36.74%	5,000,465	39.53%	35,088	0.28%	0	0.00%
No	13,601,171	1,097,544	8.07%	1,538,860	11.31%	4,389,870	32.28%	6,574,897	48.34%	0	0.00%	0	0.00%
<b>Mobility limitation</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,900,280	602,956	15.46%	477,790	12.25%	1,204,037	30.87%	1,595,708	40.91%	19,788	0.51%	0	0.00%
Mobility limitations	4,407,431	352,877	8.01%	790,909	17.94%	1,479,904	33.58%	1,768,441	40.12%	15,300	0.35%	0	0.00%
No mobility limitations	7,499,869	648,097	8.64%	1,025,267	13.67%	2,972,095	39.63%	2,854,410	38.06%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	10,443,202	464,989	4.45%	1,240,435	11.88%	3,380,975	32.37%	5,356,803	51.29%	0	0.00%	0	0.00%
<b>Sleep disorder</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,406,317	339,525	14.11%	326,645	13.57%	855,643	35.56%	884,504	36.76%	0	0.00%	0	0.00%
No	23,801,450	1,729,395	7.27%	3,191,877	13.41%	8,154,233	34.26%	10,690,858	44.92%	35,088	0.15%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	2,692,415	287,085	10.66%	434,687	16.14%	787,503	29.25%	1,183,141	43.94%	0	0.00%	0	0.00%
Once a day	2,607,295	197,240	7.56%	266,283	10.21%	610,667	23.42%	1,533,104	58.80%	0	0.00%	0	0.00%
2 or more time a week	2,294,811	61,726	2.69%	545,021	23.75%	1,175,358	51.22%	512,705	22.34%	0	0.00%	0	0.00%
Once a week	2,067,749	100,898	4.88%	266,802	12.90%	425,524	20.58%	1,267,947	61.32%	6,578	0.32%	0	0.00%
1-3 times a month	5,053,236	455,107	9.01%	626,511	12.40%	2,013,696	39.85%	1,957,921	38.75%	0	0.00%	0	0.00%
Never	11,298,450	888,412	7.86%	1,328,999	11.76%	3,980,037	35.23%	5,085,702	45.01%	15,300	0.14%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	963,213	181,653	18.86%	114,343	11.87%	401,818	41.72%	265,399	27.55%	0	0.00%	0	0.00%
Never	24,615,330	1,718,650	6.98%	3,297,536	13.40%	8,482,148	34.46%	11,095,118	45.07%	21,878	0.09%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,092,036	110,448	10.11%	130,903	11.99%	558,490	51.14%	276,895	25.36%	15,300	1.40%	0	0.00%
30 to 34	1,445,999	61,912	4.28%	115,457	7.98%	666,127	46.07%	602,503	41.67%	0	0.00%	0	0.00%
35 to 39	1,852,481	143,885	7.77%	247,163	13.34%	725,485	39.16%	735,947	39.73%	0	0.00%	0	0.00%
40 to 44	3,215,975	361,620	11.24%	521,427	16.21%	1,016,694	31.61%	1,303,024	40.52%	13,209	0.41%	0	0.00%
45 to 49	4,624,738	384,067	8.30%	1,002,044	21.67%	1,551,252	33.54%	1,687,375	36.49%	0	0.00%	0	0.00%
50 to 54	3,240,354	257,770	7.95%	366,273	11.30%	956,240	29.51%	1,653,493	51.03%	6,578	0.20%	0	0.00%
55 or older	2,238,244	149,784	6.69%	300,632	13.43%	981,432	43.85%	806,395	36.03%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	8,469,331	565,586	6.68%	850,501	10.04%	2,543,513	30.03%	4,509,730	53.25%	0	0.00%	0	0.00%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	2,580,354	0	0.00%	518,925	20.11%	823,051	31.90%	1,238,378	47.99%	0	0.00%	0	0.00%
1	2,621,413	318,534	12.15%	357,134	13.62%	810,972	30.94%	1,134,773	43.29%	0	0.00%	0	0.00%
2	6,031,539	446,117	7.40%	719,441	11.93%	1,843,223	30.56%	3,009,549	49.90%	13,209	0.22%	0	0.00%
More than 2	14,959,853	1,304,269	8.72%	1,938,902	12.96%	5,502,143	36.78%	6,192,661	41.40%	21,878	0.15%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	5,804,859	136,836	2.36%	958,341	16.51%	2,046,033	35.25%	2,663,649	45.89%	0	0.00%	0	0.00%
1	3,566,141	449,230	12.60%	479,530	13.45%	1,081,085	30.32%	1,556,296	43.64%	0	0.00%	0	0.00%
2	6,946,976	342,362	4.93%	757,951	10.91%	2,589,711	37.28%	3,243,742	46.69%	13,209	0.19%	0	0.00%
More than 2	9,816,697	1,140,491	11.62%	1,311,677	13.36%	3,230,977	32.91%	4,111,674	41.88%	21,878	0.22%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	7,177,469	553,368	7.71%	704,367	9.81%	2,851,312	39.73%	3,068,422	42.75%	0	0.00%	0	0.00%
1	2,221,278	154,837	6.97%	411,377	18.52%	753,601	33.93%	901,463	40.58%	0	0.00%	0	0.00%
2	1,299,049	93,258	7.18%	172,852	13.31%	475,060	36.57%	557,879	42.95%	0	0.00%	0	0.00%
More than 2	732,226	59,590	8.14%	211,416	28.87%	170,073	23.23%	291,148	39.76%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	14,682,052	1,207,866	8.23%	2,007,487	13.67%	4,755,383	32.39%	6,676,228	45.47%	35,088	0.24%	0	0.00%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,606,796	0	0.00%	665,891	18.46%	1,399,755	38.81%	1,541,151	42.73%	0	0.00%	0	0.00%
Yes	4,326,233	295,405	6.83%	433,005	10.01%	1,288,992	29.79%	2,308,831	53.37%	0	0.00%	0	0.00%
No	18,280,301	1,773,515	9.70%	2,430,698	13.30%	6,315,620	34.55%	7,725,380	42.26%	35,088	0.19%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,644,248	0	0.00%	670,697	18.40%	1,432,400	39.31%	1,541,151	42.29%	0	0.00%	0	0.00%
35 years old or younger	3,922,334	287,491	7.33%	419,159	10.69%	1,231,235	31.39%	1,984,448	50.59%	0	0.00%	0	0.00%
36 years or older	395,986	0	0.00%	13,846	3.50%	57,757	14.59%	324,383	81.92%	0	0.00%	0	0.00%
Refused/Don't know	*	*		*		*		*		*		*	
N/A - Did not have baby weighing 9 lbs or more	18,280,301	1,773,515	9.70%	2,430,698	13.30%	6,315,620	34.55%	7,725,380	42.26%	35,088	0.19%	0	0.00%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,026,443	0	0.00%	146,966	14.32%	576,704	56.18%	302,773	29.50%	0	0.00%	0	0.00%
29 years old or younger	17,896,821	1,622,417	9.07%	2,343,623	13.10%	6,242,646	34.88%	7,672,835	42.87%	15,300	0.09%	0	0.00%
30 - 40 years old	1,154,942	34,925	3.02%	69,371	6.01%	346,847	30.03%	684,012	59.22%	19,788	1.71%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,580,354	0	0.00%	518,925	20.11%	823,051	31.90%	1,238,378	47.99%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	3,505,690	411,577	11.74%	433,148	12.36%	983,599	28.06%	1,677,365	47.85%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,433,545	76,525	5.34%	165,644	11.55%	673,226	46.96%	518,150	36.14%	0	0.00%	0	0.00%
29 years old or younger	12,999,816	1,057,469	8.13%	1,743,717	13.41%	4,700,179	36.16%	5,483,152	42.18%	15,300	0.12%	0	0.00%
30 - 40 years old	8,310,231	761,422	9.16%	1,065,371	12.82%	2,623,200	31.57%	3,840,450	46.21%	19,788	0.24%	0	0.00%
41 years or older	717,691	161,733	22.54%	0	0.00%	98,033	13.66%	457,925	63.81%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,580,354	0	0.00%	518,925	20.11%	823,051	31.90%	1,238,378	47.99%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	9,363,305	947,699	10.12%	1,828,651	19.53%	3,870,395	41.34%	2,716,560	29.01%	0	0.00%	0	0.00%
No	16,849,984	1,121,220	6.65%	1,705,750	10.12%	5,166,617	30.66%	8,821,309	52.35%	35,088	0.21%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
35 years old or younger	2,492,721	181,340	7.27%	207,270	8.32%	1,435,032	57.57%	669,079	26.84%	0	0.00%	0	0.00%
36 - 50 years old	5,476,908	669,830	12.23%	1,433,039	26.17%	1,536,364	28.05%	1,837,674	33.55%	0	0.00%	0	0.00%
51 years or older	1,199,334	96,529	8.05%	153,683	12.81%	739,316	61.64%	209,807	17.49%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	16,849,984	1,121,220	6.65%	1,705,750	10.12%	5,166,617	30.66%	8,821,309	52.35%	35,088	0.21%	0	0.00%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	5,305,756	574,711	10.83%	1,108,546	20.89%	2,123,904	40.03%	1,498,594	28.24%	0	0.00%	0	0.00%
No	20,579,534	1,440,131	7.00%	2,327,108	11.31%	6,825,035	33.16%	9,952,171	48.36%	35,088	0.17%	0	0.00%
Don't know	222,347	54,077	24.32%	71,845	32.31%	70,809	31.85%	25,615	11.52%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	365,492	54,077	14.80%	98,747	27.02%	88,072	24.10%	124,596	34.09%	0	0.00%	0	0.00%
35 years old or younger	843,180	108,866	12.91%	7,145	0.85%	417,608	49.53%	309,560	36.71%	0	0.00%	0	0.00%
36 - 50 years old	3,322,455	433,018	13.03%	917,412	27.61%	991,129	29.83%	980,895	29.52%	0	0.00%	0	0.00%
51 years or older	1,024,360	32,827	3.20%	146,823	14.33%	636,571	62.14%	208,139	20.32%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	20,579,534	1,440,131	7.00%	2,327,108	11.31%	6,825,035	33.16%	9,952,171	48.36%	35,088	0.17%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2007-2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	26,250,782	2,068,919	7.88%	3,534,401	13.46%	9,037,012	34.43%	11,575,362	44.10%	35,088	0.13%	0	0.00%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	9,166,381	947,699	10.34%	1,741,607	19.00%	3,853,413	42.04%	2,623,662	28.62%	0	0.00%	0	0.00%
No	17,046,908	1,121,220	6.58%	1,792,794	10.52%	5,183,599	30.41%	8,914,207	52.29%	35,088	0.21%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Age at screening</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	1,840,192	214,233	11.64%	112,275	6.10%	247,855	13.47%	1,265,829	68.79%	0	0.00%	0	0.00%
30-39	2,687,312	147,353	5.48%	261,237	9.72%	777,725	28.94%	1,500,996	55.85%	0	0.00%	0	0.00%
40-49	4,635,591	375,460	8.10%	390,467	8.42%	1,860,418	40.13%	2,009,245	43.34%	0	0.00%	0	0.00%
50-59	5,211,647	152,046	2.92%	450,343	8.64%	2,132,634	40.92%	2,463,862	47.28%	12,763	0.24%	0	0.00%
60-64	2,437,163	115,823	4.75%	413,879	16.98%	957,213	39.28%	950,248	38.99%	0	0.00%	0	0.00%
65-69	2,072,827	100,141	4.83%	338,229	16.32%	820,620	39.59%	813,837	39.26%	0	0.00%	0	0.00%
70-74	2,365,064	188,223	7.96%	469,098	19.83%	752,744	31.83%	954,999	40.38%	0	0.00%	0	0.00%
75-79	1,777,635	89,419	5.03%	436,551	24.56%	582,606	32.77%	623,538	35.08%	45,521	2.56%	0	0.00%
80+	2,295,448	495,267	21.58%	432,156	18.83%	854,621	37.23%	513,403	22.37%	0	0.00%	0	0.00%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	2,092,224	138,898	6.64%	167,095	7.99%	702,381	33.57%	1,071,088	51.19%	12,763	0.61%	0	0.00%
Other Hispanic	1,184,251	86,503	7.30%	103,106	8.71%	395,748	33.42%	598,894	50.57%	0	0.00%	0	0.00%
Non-Hispanic White	17,480,786	1,337,885	7.65%	2,417,510	13.83%	6,501,905	37.19%	7,177,964	41.06%	45,521	0.26%	0	0.00%
Non-Hispanic Black	3,812,462	296,784	7.78%	428,347	11.24%	1,195,040	31.35%	1,892,292	49.63%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	848,083	17,896	2.11%	188,179	22.19%	205,765	24.26%	436,243	51.44%	0	0.00%	0	0.00%
<b>Education</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	635,598	81,956	12.89%	203,802	32.06%	99,184	15.60%	250,656	39.44%	0	0.00%	0	0.00%
Less Than 9th Grade	1,844,541	145,008	7.86%	203,628	11.04%	715,772	38.80%	767,371	41.60%	12,763	0.69%	0	0.00%
9-11th Grade	3,504,140	420,423	12.00%	722,662	20.62%	1,172,858	33.47%	1,188,197	33.91%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	19,358,370	1,224,555	6.33%	2,162,529	11.17%	6,989,184	36.10%	8,936,581	46.16%	45,521	0.24%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Marital status</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	13,148,028	728,509	5.54%	1,870,262	14.22%	5,237,195	39.83%	5,312,062	40.40%	0	0.00%	0	0.00%
Widowed	3,965,487	507,798	12.81%	726,838	18.33%	1,159,784	29.25%	1,512,782	38.15%	58,284	1.47%	0	0.00%
Divorced	2,951,626	207,258	7.02%	175,314	5.94%	1,016,932	34.45%	1,552,121	52.59%	0	0.00%	0	0.00%
Separated	849,896	120,458	14.17%	41,500	4.88%	224,711	26.44%	463,228	54.50%	0	0.00%	0	0.00%
Never Married	2,776,325	248,959	8.97%	370,848	13.36%	809,484	29.16%	1,347,034	48.52%	0	0.00%	0	0.00%
Living with Partner	1,455,784	64,984	4.46%	119,474	8.21%	374,563	25.73%	896,763	61.60%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,897,549	106,659	5.62%	149,069	7.86%	687,116	36.21%	954,705	50.31%	0	0.00%	0	0.00%
Poverty income ratio less than 1	4,290,978	512,084	11.93%	424,566	9.89%	1,510,704	35.21%	1,843,624	42.97%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	19,229,279	1,259,222	6.55%	2,730,601	14.20%	6,803,018	35.38%	8,378,153	43.57%	58,284	0.30%	0	0.00%
<b>Body mass index</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	5,714,127	327,711	5.74%	478,676	8.38%	1,697,781	29.71%	3,209,959	56.18%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	7,234,091	478,436	6.61%	953,575	13.18%	2,542,233	35.14%	3,259,846	45.06%	0	0.00%	0	0.00%
Obese: 30 or higher	12,207,800	1,035,081	8.48%	1,755,096	14.38%	4,676,826	38.31%	4,682,513	38.36%	58,284	0.48%	0	0.00%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	853,886	52,028	6.09%	77,106	9.03%	327,810	38.39%	396,942	46.49%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	13,935,553	861,988	6.19%	1,727,368	12.40%	4,668,468	33.50%	6,664,966	47.83%	12,763	0.09%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	7,795,151	691,216	8.87%	929,622	11.93%	2,848,516	36.54%	3,280,277	42.08%	45,521	0.58%	0	0.00%
Glycohemoglobin 6.5% or higher	2,833,217	272,734	9.63%	570,141	20.12%	1,156,044	40.80%	834,298	29.45%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Diabetes</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,128,037	304,411	7.37%	767,401	18.59%	1,719,175	41.65%	1,337,050	32.39%	0	0.00%	0	0.00%
No	21,289,769	1,573,555	7.39%	2,536,835	11.92%	7,281,663	34.20%	9,839,432	46.22%	58,284	0.27%	0	0.00%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,091,913	685,901	6.18%	1,540,744	13.89%	4,005,225	36.11%	4,860,042	43.82%	0	0.00%	0	0.00%
No	14,325,894	1,192,064	8.32%	1,763,492	12.31%	4,995,613	34.87%	6,316,440	44.09%	58,284	0.41%	0	0.00%
<b>High blood pressure</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	11,606,108	1,172,975	10.11%	1,962,996	16.91%	4,456,101	38.39%	4,014,036	34.59%	0	0.00%	0	0.00%
No	13,796,321	704,990	5.11%	1,341,240	9.72%	4,544,737	32.94%	7,147,070	51.80%	58,284	0.42%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	5,124,229	659,000	12.86%	1,174,687	22.92%	1,361,281	26.57%	1,929,260	37.65%	0	0.00%	0	0.00%
Mobility limitations	3,894,488	423,231	10.87%	732,892	18.82%	1,672,294	42.94%	1,053,308	27.05%	12,763	0.33%	0	0.00%
No mobility limitations	6,222,816	353,596	5.68%	621,730	9.99%	2,499,835	40.17%	2,702,134	43.42%	45,521	0.73%	0	0.00%
N/A - Respondent not asked mobility limitation questions	10,176,273	442,138	4.34%	774,927	7.62%	3,467,428	34.07%	5,491,779	53.97%	0	0.00%	0	0.00%
<b>Sleep disorder</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,780,568	218,717	7.87%	248,085	8.92%	1,336,524	48.07%	964,479	34.69%	12,763	0.46%	0	0.00%
No	22,630,326	1,659,248	7.33%	3,056,151	13.50%	7,664,314	33.87%	10,205,091	45.09%	45,521	0.20%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	2,801,521	308,696	11.02%	550,648	19.66%	1,048,783	37.44%	893,394	31.89%	0	0.00%	0	0.00%
Once a day	2,380,873	236,545	9.94%	168,423	7.07%	703,914	29.57%	1,271,990	53.43%	0	0.00%	0	0.00%
2 or more time a week	2,678,116	136,194	5.09%	405,070	15.13%	1,137,485	42.47%	999,367	37.32%	0	0.00%	0	0.00%
Once a week	1,874,622	105,454	5.63%	282,617	15.08%	773,653	41.27%	712,897	38.03%	0	0.00%	0	0.00%
1-3 times a month	3,679,787	252,943	6.87%	612,153	16.64%	1,457,391	39.61%	1,357,300	36.89%	0	0.00%	0	0.00%
Never	11,808,270	781,626	6.62%	1,159,214	9.82%	3,867,611	32.75%	5,941,533	50.32%	58,284	0.49%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	688,088	20,873	3.03%	122,855	17.85%	328,560	47.75%	215,800	31.36%	0	0.00%	0	0.00%
Never	24,040,280	1,743,296	7.25%	2,913,362	12.12%	8,461,875	35.20%	10,863,462	45.19%	58,284	0.24%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	1,039,094	196,339	18.90%	293,059	28.20%	290,345	27.94%	259,350	24.96%	0	0.00%	0	0.00%
30 to 34	1,006,929	109,041	10.83%	262,053	26.02%	381,820	37.92%	254,014	25.23%	0	0.00%	0	0.00%
35 to 39	1,347,858	153,375	11.38%	303,273	22.50%	431,273	32.00%	459,937	34.12%	0	0.00%	0	0.00%
40 to 44	2,873,502	170,101	5.92%	540,067	18.79%	1,136,364	39.55%	968,686	33.71%	58,284	2.03%	0	0.00%
45 to 49	3,356,466	428,126	12.76%	276,098	8.23%	1,026,873	30.59%	1,625,369	48.43%	0	0.00%	0	0.00%
50 to 54	4,611,844	221,568	4.80%	595,025	12.90%	2,143,395	46.48%	1,651,856	35.82%	0	0.00%	0	0.00%
55 or older	1,942,557	252,978	13.02%	280,014	14.41%	790,173	40.68%	619,391	31.89%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	9,032,022	346,437	3.84%	704,636	7.80%	2,769,150	30.66%	5,211,799	57.70%	0	0.00%	0	0.00%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
None	2,652,878	59,378	2.24%	215,777	8.13%	630,928	23.78%	1,746,795	65.85%	0	0.00%	0	0.00%
1	2,655,546	210,350	7.92%	259,124	9.76%	738,272	27.80%	1,447,799	54.52%	0	0.00%	0	0.00%
2	5,674,202	509,005	8.97%	632,139	11.14%	1,998,519	35.22%	2,534,538	44.67%	0	0.00%	0	0.00%
More than 2	14,435,181	1,099,232	7.61%	2,197,196	15.22%	5,633,118	39.02%	5,447,350	37.74%	58,284	0.40%	0	0.00%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	5,869,186	237,411	4.05%	468,407	7.98%	1,893,682	32.26%	3,269,686	55.71%	0	0.00%	0	0.00%
1	3,586,898	315,885	8.81%	478,398	13.34%	1,003,491	27.98%	1,789,125	49.88%	0	0.00%	0	0.00%
2	6,391,261	448,068	7.01%	749,306	11.72%	2,568,183	40.18%	2,625,704	41.08%	0	0.00%	0	0.00%
More than 2	9,434,881	847,280	8.98%	1,608,126	17.04%	3,515,744	37.26%	3,405,447	36.09%	58,284	0.62%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	7,632,579	402,974	5.28%	853,219	11.18%	3,147,665	41.24%	3,228,721	42.30%	0	0.00%	0	0.00%
1	2,299,292	164,703	7.16%	206,810	8.99%	801,881	34.88%	1,125,898	48.97%	0	0.00%	0	0.00%
2	1,312,038	100,991	7.70%	69,160	5.27%	550,740	41.98%	591,147	45.06%	0	0.00%	0	0.00%
More than 2	623,369	119,655	19.19%	110,561	17.74%	187,144	30.02%	206,009	33.05%	0	0.00%	0	0.00%
N/A - Vaginal delivery/deliveries only	13,317,326	1,060,321	7.96%	1,991,579	14.95%	4,293,669	32.24%	5,913,473	44.40%	58,284	0.44%	0	0.00%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,861,897	140,520	3.64%	233,642	6.05%	1,138,073	29.47%	2,349,662	60.84%	0	0.00%	0	0.00%
Yes	4,855,510	442,082	9.10%	621,208	12.79%	1,780,933	36.68%	1,998,523	41.16%	12,763	0.26%	0	0.00%
No	16,538,839	1,289,542	7.80%	2,398,620	14.50%	5,999,149	36.27%	6,806,006	41.15%	45,521	0.28%	0	0.00%
Don't know	161,561	5,822	3.60%	50,766	31.42%	82,684	51.18%	22,290	13.80%	0	0.00%	0	0.00%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	4,023,458	146,341	3.64%	284,407	7.07%	1,220,757	30.34%	2,371,952	58.95%	0	0.00%	0	0.00%
35 years old or younger	4,345,629	412,144	9.48%	576,202	13.26%	1,454,732	33.48%	1,889,788	43.49%	12,763	0.29%	0	0.00%
36 years or older	491,599	29,938	6.09%	26,725	5.44%	326,200	66.35%	108,735	22.12%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	16,538,839	1,289,542	7.80%	2,398,620	14.50%	5,999,149	36.27%	6,806,006	41.15%	45,521	0.28%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,209,019	81,142	6.71%	17,865	1.48%	507,145	41.95%	602,867	49.86%	0	0.00%	0	0.00%
29 years old or younger	16,270,236	1,309,118	8.05%	2,494,890	15.33%	6,050,855	37.19%	6,357,089	39.07%	58,284	0.36%	0	0.00%
30 - 40 years old	1,542,350	123,193	7.99%	68,667	4.45%	710,842	46.09%	639,648	41.47%	0	0.00%	0	0.00%
N/A - Never pregnant	2,652,878	59,378	2.24%	215,777	8.13%	630,928	23.78%	1,746,795	65.85%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	3,527,466	305,134	8.65%	487,788	13.83%	1,056,706	29.96%	1,677,837	47.56%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,344,681	101,155	7.52%	17,865	1.33%	542,714	40.36%	682,946	50.79%	0	0.00%	0	0.00%
29 years old or younger	12,042,012	1,044,849	8.68%	1,516,491	12.59%	4,624,209	38.40%	4,810,941	39.95%	45,521	0.38%	0	0.00%
30 - 40 years old	8,339,256	613,746	7.36%	1,432,389	17.18%	2,678,079	32.11%	3,602,279	43.20%	12,763	0.15%	0	0.00%
41 years or older	727,954	21,362	2.93%	71,092	9.77%	454,225	62.40%	181,275	24.90%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	2,652,878	59,378	2.24%	215,777	8.13%	630,928	23.78%	1,746,795	65.85%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	7,934,861	777,429	9.80%	1,420,613	17.90%	3,006,243	37.89%	2,730,576	34.41%	0	0.00%	0	0.00%
No	17,454,893	1,100,537	6.31%	1,855,571	10.63%	5,994,595	34.34%	8,445,906	48.39%	58,284	0.33%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
35 years old or younger	2,062,218	276,677	13.42%	417,071	20.22%	812,482	39.40%	555,988	26.96%	0	0.00%	0	0.00%
36 - 50 years old	4,064,475	333,204	8.20%	796,040	19.59%	1,369,883	33.70%	1,565,348	38.51%	0	0.00%	0	0.00%
51 years or older	1,532,708	117,210	7.65%	141,444	9.23%	768,154	50.12%	505,901	33.01%	0	0.00%	0	0.00%
Refused/Don't know	275,460	50,337	18.27%	66,058	23.98%	55,724	20.23%	103,340	37.52%	0	0.00%	0	0.00%
N/A - Did not have procedure	17,454,893	1,100,537	6.31%	1,855,571	10.63%	5,994,595	34.34%	8,445,906	48.39%	58,284	0.33%	0	0.00%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	428,996	29,321	6.83%	97,071	22.63%	134,349	31.32%	168,255	39.22%	0	0.00%	0	0.00%
Yes	3,927,401	306,909	7.81%	521,062	13.27%	1,769,811	45.06%	1,329,619	33.85%	0	0.00%	0	0.00%
No	20,752,776	1,514,332	7.30%	2,638,100	12.71%	6,921,767	33.35%	9,620,293	46.36%	58,284	0.28%	0	0.00%
Don't know	308,633	27,403	8.88%	48,004	15.55%	174,911	56.67%	58,315	18.89%	0	0.00%	0	0.00%
<b>Age when ovaries removed</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	737,629	56,724	7.69%	145,075	19.67%	309,260	41.93%	226,570	30.72%	0	0.00%	0	0.00%
35 years old or younger	891,251	113,244	12.71%	244,989	27.49%	307,371	34.49%	225,648	25.32%	0	0.00%	0	0.00%
36 - 50 years old	2,089,328	105,335	5.04%	227,551	10.89%	865,988	41.45%	890,455	42.62%	0	0.00%	0	0.00%
51 years or older	766,122	37,290	4.87%	48,522	6.33%	570,133	74.42%	110,177	14.38%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	20,752,776	1,514,332	7.30%	2,638,100	12.71%	6,921,767	33.35%	9,620,293	46.36%	58,284	0.28%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.045: How frequently does this (referring to KI.Q.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among women, 2009-2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
<b>Total</b>	25,417,806	1,877,965	7.39%	3,304,236	13.00%	9,000,838	35.41%	11,176,482	43.97%	58,284	0.23%	0	0.00%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	7,918,952	777,429	9.82%	1,420,613	17.94%	2,990,334	37.76%	2,730,576	34.48%	0	0.00%	0	0.00%
No	17,470,803	1,100,537	6.30%	1,855,571	10.62%	6,010,504	34.40%	8,445,906	48.34%	58,284	0.33%	0	0.00%
Don't know	*	*		*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.