

Table F04 - Prevalence of mixed urinary incontinence among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*
20-29	10,228,786	865,887	8.47%	4.56%	12.37%	9,362,899	91.53%
30-39	13,268,013	1,851,360	13.95%	8.55%	19.36%	11,416,653	86.05%
40-49	14,571,797	3,090,696	21.21%	14.99%	27.43%	11,481,101	78.79%
50-59	10,626,450	3,216,813	30.27%	24.36%	36.19%	7,409,637	69.73%
60-64	4,060,065	1,470,922	36.23%	24.65%	47.81%	2,589,143	63.77%
65-69	3,563,926	1,308,014	36.70%	23.45%	49.96%	2,255,912	63.30%
70-74	2,889,907	1,075,711	37.22%	26.30%	48.14%	1,814,196	62.78%
75-79	2,640,163	1,302,804	49.35%	35.48%	63.21%	1,337,359	50.65%
80-84	2,019,122	1,019,337	50.48%	36.29%	64.68%	999,785	49.52%
85+	1,326,554	603,277	45.48%	23.06%	67.90%	723,277	54.52%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	4,135,524	750,000	18.14%	14.01%	22.26%	3,385,524	81.86%
Other Hispanic	4,212,300	706,449	16.77%	7.89%	25.65%	3,505,851	83.23%
Non-Hispanic White	45,120,970	12,269,488	27.19%	23.65%	30.73%	32,851,482	72.81%
Non-Hispanic Black	8,591,298	1,350,620	15.72%	10.58%	20.86%	7,240,678	84.28%
Other Race - Including Multi-Racial	3,355,177	728,264	21.71%	5.59%	37.83%	2,626,913	78.29%
Education	-	-	-	-	-	-	-
Missing	1,845,302	418,835	22.70%	8.38%	37.02%	1,426,468	77.30%
Less Than 9th Grade	4,848,546	1,470,075	30.32%	23.01%	37.63%	3,378,471	69.68%
9-11th Grade	7,883,954	1,923,216	24.39%	15.51%	33.28%	5,960,738	75.61%
High School Grad/GED or Equivalent	50,643,207	11,982,670	23.66%	20.74%	26.58%	38,660,537	76.34%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	522,575	47,932	9.17%	(4.79%)	23.14%	474,643	90.83%
Married	36,467,967	8,843,655	24.25%	20.03%	28.48%	27,624,312	75.75%
Widowed	6,833,659	3,032,474	44.38%	33.85%	54.90%	3,801,185	55.62%
Divorced	6,321,172	1,487,072	23.53%	16.09%	30.96%	4,834,100	76.47%
Separated	2,838,399	573,193	20.19%	10.21%	30.18%	2,265,206	79.81%
Never Married	8,156,151	1,557,478	19.10%	13.33%	24.86%	6,598,673	80.90%
Living with Partner	4,228,639	238,882	5.65%	1.77%	9.53%	3,989,757	94.35%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,303,863	906,283	21.06%	12.53%	29.58%	3,397,580	78.94%
Poverty income ratio less than 1	10,728,977	2,576,319	24.01%	19.42%	28.61%	8,152,658	75.99%
Poverty income ratio 1 or greater	50,382,429	12,322,220	24.46%	21.07%	27.85%	38,060,209	75.54%
Body mass index	-	-	-	-	-	-	-
Missing	2,764,968	1,337,501	48.37%	35.30%	61.45%	1,427,467	51.63%
Normal: less than 25	25,994,755	3,798,719	14.61%	10.52%	18.71%	22,196,036	85.39%
Overweight: 25.0 - 29.9	16,404,435	3,967,388	24.18%	20.89%	27.48%	12,437,047	75.82%
Obese: 30 or higher	20,251,111	6,701,213	33.09%	27.98%	38.20%	13,549,898	66.91%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,659,395	465,563	17.51%	6.31%	28.70%	2,193,832	82.49%
Glycohemoglobin < 5.7%	50,268,577	10,396,089	20.68%	17.85%	23.51%	39,872,488	79.32%
Glycohemoglobin 5.7% - 6.4%	8,889,667	3,699,229	41.61%	31.75%	51.48%	5,190,438	58.39%
Glycohemoglobin 6.5% or higher	3,597,630	1,243,941	34.58%	22.59%	46.57%	2,353,689	65.42%
Diabetes	-	-	-	-	-	-	-
Yes	4,894,686	1,790,767	36.59%	29.60%	43.57%	3,103,919	63.41%
No	60,520,583	14,014,054	23.16%	20.22%	26.09%	46,506,528	76.84%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	27,981,243	7,897,959	28.23%	24.17%	32.28%	20,083,283	71.77%
No	37,409,879	7,906,862	21.14%	17.15%	25.12%	29,503,017	78.86%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	17,400,790	6,409,561	36.83%	32.54%	41.13%	10,991,229	63.17%
No	47,825,607	9,368,304	19.59%	16.16%	23.01%	38,457,303	80.41%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	4,317,024	2,043,494	47.34%	35.68%	58.99%	2,273,529	52.66%
Mobility limitations	8,999,340	4,558,791	50.66%	42.40%	58.92%	4,440,548	49.34%
No mobility limitations	10,998,833	3,150,098	28.64%	21.63%	35.65%	7,848,735	71.36%
N/A - Respondent not asked mobility limitation questions	41,100,073	6,052,438	14.73%	11.51%	17.94%	35,047,635	85.27%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	2,382,790	907,900	38.10%	17.80%	58.41%	1,474,891	61.90%
30 to 34	2,506,237	1,143,471	45.63%	28.43%	62.82%	1,362,766	54.37%
35 to 39	3,227,642	1,540,359	47.72%	34.04%	61.41%	1,687,283	52.28%
40 to 44	3,562,377	1,204,708	33.82%	23.54%	44.09%	2,357,669	66.18%
45 to 49	6,566,057	1,934,005	29.45%	20.63%	38.28%	4,632,052	70.55%
50 to 54	6,165,689	2,860,332	46.39%	37.31%	55.47%	3,305,357	53.61%
55 or older	2,311,702	906,063	39.19%	26.41%	51.97%	1,405,639	60.81%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	38,600,701	5,290,940	13.71%	10.57%	16.84%	33,309,761	86.29%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,058,764	330,438	31.21%	13.38%	49.04%	728,327	68.79%
None	12,604,005	1,160,358	9.21%	4.18%	14.23%	11,443,647	90.79%
1	6,782,022	1,320,661	19.47%	10.32%	28.62%	5,461,361	80.53%
2	11,805,355	2,545,521	21.56%	14.18%	28.94%	9,259,834	78.44%
More than 2	33,165,122	10,447,844	31.50%	28.44%	34.56%	22,717,278	68.50%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	1,506,903	338,064	22.43%	8.68%	36.18%	1,168,839	77.57%
29 years old or younger	37,129,102	11,692,201	31.49%	28.05%	34.93%	25,436,901	68.51%
30 - 40 years old	2,206,891	502,895	22.79%	13.68%	31.89%	1,703,996	77.21%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	12,604,005	1,160,358	9.21%	4.18%	14.23%	11,443,647	90.79%
Only 1 live birth - see age at last live birth	9,788,163	1,670,908	17.07%	9.52%	24.62%	8,117,255	82.93%
No pregnancies resulting in live births	2,129,064	440,395	20.68%	0.85%	40.52%	1,688,669	79.32%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	5,053,414	1,625,729	32.17%	24.22%	40.12%	3,427,685	67.83%
29 years old or younger	27,815,094	6,261,416	22.51%	18.82%	26.20%	21,553,678	77.49%
30 - 40 years old	16,692,249	5,801,233	34.75%	29.67%	39.83%	10,891,016	65.25%
41 years or older	1,061,385	515,691	48.59%	19.64%	77.53%	545,694	51.41%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	12,604,005	1,160,358	9.21%	4.18%	14.23%	11,443,647	90.79%
No pregnancies resulting in live births	2,129,064	440,395	20.68%	0.85%	40.52%	1,688,669	79.32%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	30,350,668	4,111,794	13.55%	10.12%	16.98%	26,238,874	86.45%
Yes	13,627,313	6,490,733	47.63%	41.41%	53.85%	7,136,580	52.37%
No	21,408,395	5,202,294	24.30%	20.35%	28.25%	16,206,101	75.70%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	30,412,304	4,144,538	13.63%	10.20%	17.05%	26,267,766	86.37%
35 years old or younger	4,832,906	2,397,707	49.61%	34.84%	64.38%	2,435,199	50.39%
36 - 50 years old	7,030,034	3,064,520	43.59%	37.79%	49.40%	3,965,514	56.41%
51 years or older	1,574,680	893,529	56.74%	40.83%	72.66%	681,151	43.26%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	21,408,395	5,202,294	24.30%	20.35%	28.25%	16,206,101	75.70%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	65,415,269	15,804,821	24.16%	21.42%	26.90%	49,610,447	75.84%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	5,458,760	995,547	18.24%	7.23%	29.25%	4,463,213	81.76%
Yes	13,627,313	6,490,733	47.63%	41.41%	53.85%	7,136,580	52.37%
No	46,300,303	8,318,541	17.97%	15.28%	20.66%	37,981,762	82.03%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Age at screening	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*
20-29	10,435,336	1,069,318	10.25%	4.81%	15.68%	9,366,018	89.75%
30-39	12,056,295	2,177,492	18.06%	12.12%	24.00%	9,878,803	81.94%
40-49	12,872,412	2,802,393	21.77%	15.59%	27.95%	10,070,019	78.23%
50-59	11,731,103	3,786,816	32.28%	22.15%	42.41%	7,944,287	67.72%
60-64	3,567,033	744,245	20.86%	12.67%	29.06%	2,822,787	79.14%
65-69	4,192,276	1,889,426	45.07%	34.38%	55.76%	2,302,850	54.93%
70-74	3,464,356	1,355,350	39.12%	21.98%	56.27%	2,109,005	60.88%
75-79	3,951,361	1,884,208	47.69%	33.82%	61.55%	2,067,154	52.31%
80-84	2,272,389	696,601	30.66%	20.66%	40.65%	1,575,789	69.35%
85+	1,609,520	719,197	44.68%	28.91%	60.46%	890,323	55.32%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	4,363,237	935,381	21.44%	17.55%	25.33%	3,427,855	78.56%
Other Hispanic	2,364,116	486,064	20.56%	(2.93%)	44.05%	1,878,053	79.44%
Non-Hispanic White	48,112,095	13,317,584	27.68%	24.04%	31.32%	34,794,511	72.32%
Non-Hispanic Black	8,436,007	1,832,356	21.72%	16.52%	26.92%	6,603,651	78.28%
Other Race - Including Multi-Racial	3,280,639	553,660	16.88%	6.83%	26.92%	2,726,978	83.12%
Education	-	-	-	-	-	-	-
Missing	2,358,304	628,734	26.66%	5.56%	47.76%	1,729,570	73.34%
Less Than 9th Grade	4,928,907	1,592,065	32.30%	22.50%	42.10%	3,336,842	67.70%
9-11th Grade	7,291,632	2,365,883	32.45%	24.78%	40.11%	4,925,749	67.55%
High School Grad/GED or Equivalent	51,860,949	12,499,754	24.10%	20.33%	27.87%	39,361,195	75.90%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	483,813	70,791	14.63%	(6.04%)	35.31%	413,021	85.37%
Married	35,857,103	8,527,336	23.78%	19.53%	28.03%	27,329,767	76.22%
Widowed	7,652,616	3,138,007	41.01%	33.72%	48.29%	4,514,609	58.99%
Divorced	7,668,992	2,303,313	30.03%	17.40%	42.67%	5,365,680	69.97%
Separated	1,751,848	330,310	18.85%	4.99%	32.72%	1,421,538	81.15%
Never Married	9,286,835	1,821,157	19.61%	13.05%	26.17%	7,465,679	80.39%
Living with Partner	3,802,076	934,132	24.57%	12.19%	36.95%	2,867,944	75.43%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	3,695,371	917,960	24.84%	14.86%	34.82%	2,777,411	75.16%
Poverty income ratio less than 1	9,181,112	2,715,318	29.58%	21.48%	37.67%	6,465,794	70.42%
Poverty income ratio 1 or greater	53,679,611	13,491,769	25.13%	22.33%	27.93%	40,187,842	74.87%
Body mass index	-	-	-	-	-	-	-
Missing	1,141,703	380,953	33.37%	15.44%	51.29%	760,751	66.63%
Normal: less than 25	26,111,136	4,170,345	15.97%	12.65%	19.29%	21,940,791	84.03%
Overweight: 25.0 - 29.9	18,737,939	5,401,920	28.83%	23.02%	34.63%	13,336,019	71.17%
Obese: 30 or higher	20,565,316	7,171,829	34.87%	29.32%	40.42%	13,393,487	65.13%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,707,367	525,109	19.40%	8.58%	30.21%	2,182,258	80.60%
Glycohemoglobin < 5.7%	50,618,555	11,526,805	22.77%	19.56%	25.98%	39,091,750	77.23%
Glycohemoglobin 5.7% - 6.4%	8,862,362	3,265,237	36.84%	29.38%	44.31%	5,597,125	63.16%
Glycohemoglobin 6.5% or higher	4,367,810	1,807,896	41.39%	27.88%	54.90%	2,559,915	58.61%
Diabetes	-	-	-	-	-	-	-
Yes	5,635,836	2,640,736	46.86%	36.74%	56.98%	2,995,100	53.14%
No	60,920,258	14,484,310	23.78%	20.22%	27.33%	46,435,948	76.22%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,979,526	8,895,854	30.70%	27.52%	33.88%	20,083,672	69.30%
No	37,545,466	8,229,192	21.92%	18.05%	25.79%	29,316,273	78.08%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	19,776,048	7,830,526	39.60%	35.05%	44.15%	11,945,522	60.40%
No	46,471,487	9,236,292	19.88%	16.90%	22.85%	37,235,195	80.12%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	8,138,547	3,928,403	48.27%	36.43%	60.10%	4,210,144	51.73%
Mobility limitations	8,513,495	3,526,104	41.42%	34.63%	48.20%	4,987,391	58.58%
No mobility limitations	13,742,930	4,012,628	29.20%	24.60%	33.79%	9,730,303	70.80%
N/A - Respondent not asked mobility limitation questions	36,161,121	5,657,910	15.65%	12.09%	19.20%	30,503,211	84.35%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	1,811,649	559,041	30.86%	16.47%	45.24%	1,252,608	69.14%
Younger than 30	1,944,450	829,914	42.68%	21.32%	64.04%	1,114,536	57.32%
30 to 34	2,619,844	1,472,085	56.19%	36.54%	75.83%	1,147,759	43.81%
35 to 39	3,032,591	1,368,595	45.13%	32.67%	57.59%	1,663,996	54.87%
40 to 44	4,088,802	1,290,772	31.57%	24.85%	38.29%	2,798,030	68.43%
45 to 49	6,668,327	2,326,600	34.89%	24.52%	45.27%	4,341,727	65.11%
50 to 54	6,167,527	2,137,574	34.66%	28.95%	40.37%	4,029,952	65.34%
55 or older	3,407,624	1,567,949	46.01%	32.46%	59.56%	1,839,675	53.99%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	36,597,196	5,517,908	15.08%	11.28%	18.88%	31,079,288	84.92%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	1,576,425	552,206	35.03%	16.95%	53.10%	1,024,219	64.97%
None	13,990,772	1,925,961	13.77%	9.33%	18.20%	12,064,812	86.23%
1	9,044,711	1,640,502	18.14%	11.83%	24.44%	7,404,209	81.86%
2	15,080,531	3,363,908	22.31%	13.36%	31.25%	11,716,623	77.69%
More than 2	26,863,655	9,642,470	35.89%	31.15%	40.64%	17,221,185	64.11%
Number of vaginal deliveries	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Number of cesarean deliveries	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,017,125	469,910	23.30%	11.20%	35.40%	1,547,215	76.70%
29 years old or younger	33,824,215	11,264,545	33.30%	29.26%	37.35%	22,559,670	66.70%
30 - 40 years old	2,728,903	611,062	22.39%	8.81%	35.98%	2,117,841	77.61%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	13,990,772	1,925,961	13.77%	9.33%	18.20%	12,064,812	86.23%
Only 1 live birth - see age at last live birth	10,586,329	2,476,078	23.39%	17.01%	29.77%	8,110,251	76.61%
No pregnancies resulting in live births	3,231,023	314,560	9.74%	1.56%	17.91%	2,916,463	90.26%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	4,771,090	1,339,158	28.07%	21.33%	34.81%	3,431,931	71.93%
29 years old or younger	27,811,033	8,649,452	31.10%	26.07%	36.13%	19,161,581	68.90%
30 - 40 years old	15,311,229	4,459,148	29.12%	21.48%	36.77%	10,852,081	70.88%
41 years or older	1,271,146	406,344	31.97%	12.96%	50.98%	864,802	68.03%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	13,990,772	1,925,961	13.77%	9.33%	18.20%	12,064,812	86.23%
No pregnancies resulting in live births	3,231,023	314,560	9.74%	1.56%	17.91%	2,916,463	90.26%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	29,252,666	4,632,721	15.84%	11.43%	20.24%	24,619,945	84.16%
Yes	13,070,994	6,081,079	46.52%	39.91%	53.14%	6,989,915	53.48%
No	24,169,140	6,347,952	26.26%	21.57%	30.96%	17,821,188	73.74%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	29,315,960	4,696,015	16.02%	11.58%	20.46%	24,619,945	83.98%
35 years old or younger	4,655,861	2,559,529	54.97%	42.82%	67.13%	2,096,332	45.03%
36 - 50 years old	6,536,547	2,332,869	35.69%	28.49%	42.88%	4,203,677	64.31%
51 years or older	1,770,027	1,170,291	66.12%	53.29%	78.94%	599,737	33.88%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	24,169,140	6,347,952	26.26%	21.57%	30.96%	17,821,188	73.74%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	66,556,094	17,125,046	25.73%	22.49%	28.97%	49,431,048	74.27%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	1,495,246	462,101	30.90%	16.13%	45.68%	1,033,144	69.10%
Yes	13,070,994	6,081,079	46.52%	39.91%	53.14%	6,989,915	53.48%
No	51,926,561	10,518,572	20.26%	16.53%	23.98%	41,407,989	79.74%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Age at screening	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*
20-29	9,713,260	893,822	9.20%	6.07%	12.34%	8,819,438	90.80%
30-39	12,237,794	2,413,067	19.72%	14.07%	25.37%	9,824,727	80.28%
40-49	15,375,729	3,011,318	19.58%	13.26%	25.91%	12,364,410	80.42%
50-59	11,797,704	3,551,009	30.10%	21.69%	38.51%	8,246,695	69.90%
60-64	3,939,408	1,615,042	41.00%	31.78%	50.21%	2,324,366	59.00%
65-69	2,880,342	1,445,560	50.19%	34.82%	65.55%	1,434,782	49.81%
70-74	2,927,279	1,281,380	43.77%	29.92%	57.63%	1,645,899	56.23%
75-79	2,046,420	623,950	30.49%	16.70%	44.28%	1,422,470	69.51%
80-84	1,597,467	577,603	36.16%	22.66%	49.65%	1,019,865	63.84%
85+	1,238,200	577,631	46.65%	28.30%	65.00%	660,569	53.35%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	4,427,370	1,109,098	25.05%	16.84%	33.26%	3,318,272	74.95%
Other Hispanic	2,246,765	721,473	32.11%	18.00%	46.23%	1,525,292	67.89%
Non-Hispanic White	46,204,694	11,981,645	25.93%	22.22%	29.64%	34,223,049	74.07%
Non-Hispanic Black	7,679,717	1,613,676	21.01%	16.10%	25.92%	6,066,042	78.99%
Other Race - Including Multi-Racial	3,539,135	605,149	17.10%	6.74%	27.46%	2,933,986	82.90%
Education	-	-	-	-	-	-	-
Missing	2,008,700	464,437	23.12%	3.27%	42.97%	1,544,262	76.88%
Less Than 9th Grade	3,371,673	1,084,493	32.16%	19.77%	44.56%	2,287,180	67.84%
9-11th Grade	7,145,504	2,414,946	33.80%	26.58%	41.01%	4,730,558	66.20%
High School Grad/GED or Equivalent	51,397,840	12,067,164	23.48%	20.37%	26.59%	39,330,677	76.52%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	702,937	6,942	0.99%	(0.82%)	2.79%	695,994	99.01%
Married	36,522,683	9,637,080	26.39%	22.74%	30.03%	26,885,603	73.61%
Widowed	5,184,392	1,464,705	28.25%	19.79%	36.71%	3,719,688	71.75%
Divorced	7,993,304	2,333,186	29.19%	17.82%	40.55%	5,660,118	70.81%
Separated	2,186,999	671,051	30.68%	18.77%	42.60%	1,515,949	69.32%
Never Married	6,816,446	1,118,558	16.41%	9.84%	22.98%	5,697,888	83.59%
Living with Partner	4,604,945	799,518	17.36%	8.11%	26.61%	3,805,427	82.64%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	1,955,077	767,615	39.26%	23.98%	54.54%	1,187,462	60.74%
Poverty income ratio less than 1	7,476,362	1,951,618	26.10%	19.81%	32.40%	5,524,744	73.90%
Poverty income ratio 1 or greater	54,666,242	13,311,807	24.35%	20.89%	27.81%	41,354,435	75.65%
Body mass index	-	-	-	-	-	-	-
Missing	641,120	283,810	44.27%	16.23%	72.31%	357,310	55.73%
Normal: less than 25	26,202,495	3,879,860	14.81%	11.53%	18.08%	22,322,636	85.19%
Overweight: 25.0 - 29.9	16,187,974	4,233,810	26.15%	18.89%	33.42%	11,954,164	73.85%
Obese: 30 or higher	21,066,092	7,633,560	36.24%	31.85%	40.62%	13,432,532	63.76%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,261,270	269,142	11.90%	(0.27%)	24.07%	1,992,128	88.10%
Glycohemoglobin < 5.7%	49,210,821	11,106,533	22.57%	18.45%	26.69%	38,104,287	77.43%
Glycohemoglobin 5.7% - 6.4%	9,635,236	3,407,294	35.36%	28.58%	42.15%	6,227,942	64.64%
Glycohemoglobin 6.5% or higher	2,990,354	1,248,070	41.74%	26.84%	56.63%	1,742,284	58.26%
Diabetes	-	-	-	-	-	-	-
Yes	4,746,187	2,111,704	44.49%	33.29%	55.70%	2,634,482	55.51%
No	59,351,494	13,919,335	23.45%	20.07%	26.83%	45,432,159	76.55%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	26,200,596	6,709,387	25.61%	21.54%	29.67%	19,491,209	74.39%
No	37,888,599	9,313,167	24.58%	21.21%	27.95%	28,575,432	75.42%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	18,389,067	7,174,118	39.01%	33.32%	44.71%	11,214,949	60.99%
No	45,578,646	8,788,681	19.28%	16.37%	22.20%	36,789,965	80.72%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,817,778	3,546,155	52.01%	41.84%	62.18%	3,271,623	47.99%
Mobility limitations	5,505,206	2,857,484	51.91%	37.62%	66.19%	2,647,722	48.09%
No mobility limitations	12,057,482	3,697,868	30.67%	25.80%	35.54%	8,359,614	69.33%
N/A - Respondent not asked mobility limitation questions	39,717,215	5,929,533	14.93%	12.44%	17.42%	33,787,682	85.07%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	4,281,896	2,104,978	49.16%	28.46%	69.86%	2,176,918	50.84%
No	59,750,152	13,926,062	23.31%	19.95%	26.66%	45,824,091	76.69%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	469,165	217,472	46.35%	13.38%	79.32%	251,692	53.65%
2 or more times a day	6,684,840	2,264,836	33.88%	25.13%	42.63%	4,420,004	66.12%
Once a day	6,029,406	1,598,991	26.52%	17.11%	35.93%	4,430,415	73.48%
2 or more time a week	6,609,756	1,864,901	28.21%	20.30%	36.13%	4,744,854	71.79%
Once a week	3,873,415	1,208,792	31.21%	22.00%	40.41%	2,664,622	68.79%
1-3 times a month	8,196,624	2,831,900	34.55%	27.35%	41.75%	5,364,724	65.45%
Never	32,229,199	6,044,147	18.75%	14.62%	22.89%	26,185,052	81.25%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	469,165	217,472	46.35%	13.38%	79.32%	251,692	53.65%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	8,196,624	2,831,900	34.55%	39.41%	108.00%	5,364,724	65.45%
Never	32,229,199	6,044,147	18.75%	21.57%	27.54%	26,185,052	81.25%
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	247,985	0	0.00%	0.00%	0.00%	247,985	100.00%
Younger than 30	2,130,259	705,283	33.11%	17.51%	48.70%	1,424,975	66.89%
30 to 34	2,199,959	1,084,648	49.30%	28.75%	69.85%	1,115,311	50.70%
35 to 39	3,273,968	1,294,136	39.53%	27.04%	52.02%	1,979,832	60.47%
40 to 44	3,851,926	1,902,720	49.40%	36.17%	62.63%	1,949,206	50.60%
45 to 49	6,710,667	2,402,712	35.80%	26.64%	44.97%	4,307,956	64.20%
50 to 54	5,518,164	2,106,359	38.17%	29.56%	46.78%	3,411,805	61.83%
55 or older	1,801,695	610,703	33.90%	15.68%	52.11%	1,190,992	66.10%
Don't know	510,000	117,977	23.13%	(8.86%)	55.12%	392,022	76.87%
N/A - Had period in last 12 months	37,853,058	5,806,502	15.34%	11.52%	19.16%	32,046,557	84.66%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	12,740,275	888,335	6.97%	2.57%	11.38%	11,851,940	93.03%
1	8,811,688	1,928,859	21.89%	14.70%	29.08%	6,882,829	78.11%
2	13,332,004	3,864,818	28.99%	23.35%	34.63%	9,467,186	71.01%
More than 2	29,159,737	9,349,029	32.06%	27.47%	36.65%	19,810,708	67.94%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	758,347	13,522	1.78%	(0.97%)	4.53%	744,825	98.22%
None	23,700,106	3,164,418	13.35%	8.82%	17.88%	20,535,688	86.65%
1	9,648,777	2,452,026	25.41%	19.32%	31.50%	7,196,751	74.59%
2	12,766,467	4,184,624	32.78%	27.60%	37.96%	8,581,843	67.22%
More than 2	17,200,955	6,216,450	36.14%	30.13%	42.15%	10,984,505	63.86%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	781,376	13,522	1.73%	(0.93%)	4.39%	767,854	98.27%
None	18,080,385	5,375,140	29.73%	25.77%	33.68%	12,705,244	70.27%
1	6,576,652	1,215,953	18.49%	8.22%	28.76%	5,360,699	81.51%
2	3,441,378	992,489	28.84%	12.96%	44.72%	2,448,889	71.16%
More than 2	1,470,928	420,712	28.60%	18.53%	38.67%	1,050,216	71.40%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	33,673,827	8,013,224	23.80%	19.86%	27.73%	25,660,603	76.20%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
N/A for these years of data	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
N/A for these years of data	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,162,561	213,101	9.85%	1.14%	18.57%	1,949,460	90.15%
29 years old or younger	33,112,084	11,205,699	33.84%	30.14%	37.55%	21,906,385	66.16%
30 - 40 years old	2,191,100	500,186	22.83%	11.61%	34.05%	1,690,914	77.17%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	12,740,275	888,335	6.97%	2.57%	11.38%	11,851,940	93.03%
Only 1 live birth - see age at last live birth	10,593,985	2,512,490	23.72%	17.35%	30.08%	8,081,495	76.28%
No pregnancies resulting in live births	3,066,109	613,953	20.02%	5.46%	34.58%	2,452,156	79.98%
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	2,206,610	213,101	9.66%	1.03%	18.29%	1,993,509	90.34%
29 years old or younger	27,846,240	8,094,572	29.07%	24.27%	33.87%	19,751,668	70.93%
30 - 40 years old	17,127,051	5,732,119	33.47%	26.36%	40.58%	11,394,932	66.53%
41 years or older	773,419	334,602	43.26%	24.78%	61.75%	438,817	56.74%
Refused/Don't know	337,977	154,359	45.67%	11.74%	79.60%	183,619	54.33%
N/A - Never pregnant	12,740,275	888,335	6.97%	2.57%	11.38%	11,851,940	93.03%
No pregnancies resulting in live births	3,066,109	613,953	20.02%	5.46%	34.58%	2,452,156	79.98%
**Had hysterectomy	-	-	-	-	-	-	-
Missing	31,341,024	5,072,440	16.18%	11.93%	20.44%	26,268,584	83.82%
Yes	13,027,346	5,480,242	42.07%	34.51%	49.62%	7,547,104	57.93%
No	19,695,943	5,444,989	27.65%	21.35%	33.94%	14,250,954	72.35%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	31,374,392	5,105,808	16.27%	12.10%	20.45%	26,268,584	83.73%
35 years old or younger	4,332,804	1,828,256	42.20%	32.01%	52.38%	2,504,548	57.80%
36 - 50 years old	6,944,676	3,040,872	43.79%	31.64%	55.93%	3,903,803	56.21%
51 years or older	1,073,411	425,441	39.63%	17.78%	61.48%	647,971	60.37%
Refused/Don't know	676,455	185,673	27.45%	6.54%	48.36%	490,782	72.55%
N/A - Did not have procedure	19,695,943	5,444,989	27.65%	21.35%	33.94%	14,250,954	72.35%
Had ovaries removed	-	-	-	-	-	-	-
N/A for these years of data	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
Age when ovaries were removed	-	-	-	-	-	-	-
N/A for these years of data	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	64,097,681	16,031,040	25.01%	21.94%	28.08%	48,066,641	74.99%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	13,027,346	5,480,242	42.07%	34.51%	49.62%	7,547,104	57.93%
No	51,036,967	10,517,429	20.61%	17.38%	23.83%	40,519,538	79.39%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
Age at screening	-	-	-	-	-	-	-
10-19	540,596	36,049	6.67%	(7.79%)	21.13%	504,547	93.33%
20-29	10,407,479	721,666	6.93%	4.21%	9.66%	9,685,813	93.07%
30-39	11,949,661	1,418,125	11.87%	7.66%	16.08%	10,531,536	88.13%
40-49	12,022,008	3,345,274	27.83%	20.71%	34.94%	8,676,733	72.17%
50-59	11,519,273	3,796,914	32.96%	26.57%	39.35%	7,722,358	67.04%
60-64	4,842,588	1,607,880	33.20%	19.64%	46.77%	3,234,708	66.80%
65-69	3,429,847	1,308,396	38.15%	25.95%	50.34%	2,121,451	61.85%
70-74	2,992,498	1,500,804	50.15%	35.50%	64.81%	1,491,693	49.85%
75-79	2,416,031	853,664	35.33%	21.36%	49.31%	1,562,367	64.67%
80+	3,600,797	1,531,628	42.54%	33.40%	51.68%	2,069,169	57.46%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	4,666,441	1,095,740	23.48%	19.40%	27.56%	3,570,701	76.52%
Other Hispanic	3,451,035	811,063	23.50%	17.52%	29.49%	2,639,972	76.50%
Non-Hispanic White	43,724,749	11,925,113	27.27%	23.96%	30.59%	31,799,636	72.73%
Non-Hispanic Black	8,037,695	1,730,961	21.54%	16.10%	26.97%	6,306,734	78.46%
Other Race - Including Multi-Racial	3,840,857	557,525	14.52%	6.05%	22.98%	3,283,333	85.48%
Education	-	-	-	-	-	-	-
Missing	2,003,818	435,383	21.73%	8.10%	35.35%	1,568,435	78.27%
Less Than 9th Grade	3,803,704	1,446,291	38.02%	26.93%	49.12%	2,357,413	61.98%
9-11th Grade	8,863,362	2,679,159	30.23%	24.65%	35.80%	6,184,202	69.77%
High School Grad/GED or Equivalent	49,038,278	11,547,951	23.55%	20.91%	26.18%	37,490,327	76.45%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,012,617	122,357	12.08%	(0.21%)	24.38%	890,260	87.92%
Married	35,143,140	8,955,357	25.48%	22.50%	28.46%	26,187,783	74.52%
Widowed	5,684,784	2,303,243	40.52%	34.17%	46.86%	3,381,541	59.48%
Divorced	7,659,214	2,508,020	32.75%	23.55%	41.94%	5,151,194	67.25%
Separated	1,264,113	319,192	25.25%	14.03%	36.48%	944,921	74.75%
Never Married	8,706,281	1,269,406	14.58%	10.36%	18.80%	7,436,874	85.42%
Living with Partner	4,250,628	642,825	15.12%	3.33%	26.92%	3,607,803	84.88%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,702,316	1,046,215	22.25%	14.82%	29.67%	3,656,101	77.75%
Poverty income ratio less than 1	9,468,126	2,547,305	26.90%	20.80%	33.01%	6,920,821	73.10%
Poverty income ratio 1 or greater	49,550,335	12,526,880	25.28%	22.02%	28.54%	37,023,454	74.72%
Body mass index	-	-	-	-	-	-	-
Missing	636,095	318,183	50.02%	30.23%	69.81%	317,912	49.98%
Normal: less than 25	23,877,081	3,854,729	16.14%	12.29%	20.00%	20,022,352	83.86%
Overweight: 25.0 - 29.9	18,729,571	4,808,312	25.67%	20.87%	30.48%	13,921,259	74.33%
Obese: 30 or higher	20,478,030	7,139,177	34.86%	29.00%	40.73%	13,338,854	65.14%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,746,785	529,746	19.29%	10.62%	27.95%	2,217,039	80.71%
Glycohemoglobin < 5.7%	44,063,042	8,310,037	18.86%	15.84%	21.88%	35,753,005	81.14%
Glycohemoglobin 5.7% - 6.4%	13,018,408	5,316,067	40.84%	33.92%	47.75%	7,702,341	59.16%
Glycohemoglobin 6.5% or higher	3,892,542	1,964,551	50.47%	39.49%	61.45%	1,927,992	49.53%
Diabetes	-	-	-	-	-	-	-
Yes	5,570,473	3,038,053	54.54%	43.27%	65.80%	2,532,419	45.46%
No	58,150,304	13,082,347	22.50%	20.13%	24.86%	45,067,957	77.50%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	25,021,667	7,879,105	31.49%	28.49%	34.49%	17,142,562	68.51%
No	38,634,927	8,241,296	21.33%	18.16%	24.50%	30,393,632	78.67%
Refused	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	18,481,930	7,791,866	42.16%	36.38%	47.93%	10,690,064	57.84%
No	45,155,263	8,328,535	18.44%	16.11%	20.77%	36,826,728	81.56%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,960,650	2,679,157	38.49%	32.03%	44.95%	4,281,493	61.51%
Mobility limitations	6,601,825	2,946,374	44.63%	33.00%	56.26%	3,655,451	55.37%
No mobility limitations	11,901,621	3,997,096	33.58%	26.97%	40.20%	7,904,526	66.42%
N/A - Respondent not asked mobility limitation questions	38,256,681	6,497,774	16.98%	14.62%	19.35%	31,758,907	83.02%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	4,475,421	1,620,643	36.21%	22.39%	50.03%	2,854,778	63.79%
No	59,202,341	14,456,743	24.42%	22.15%	26.68%	44,745,598	75.58%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	355,534	110,561	31.10%	5.64%	56.55%	244,972	68.90%
2 or more times a day	5,511,973	1,765,177	32.02%	19.56%	44.49%	3,746,796	67.98%
Once a day	6,521,669	1,679,704	25.76%	20.63%	30.88%	4,841,965	74.24%
2 or more time a week	4,319,998	1,406,278	32.55%	25.75%	39.35%	2,913,721	67.45%
Once a week	3,706,436	1,179,704	31.83%	17.14%	46.52%	2,526,732	68.17%
1-3 times a month	9,322,159	3,577,364	38.37%	31.28%	45.47%	5,744,795	61.63%
Never	33,783,053	6,362,996	18.83%	16.08%	21.59%	27,420,057	81.17%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	355,534	110,561	31.10%	5.64%	56.55%	244,972	68.90%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	9,322,159	3,577,364	38.37%	74.74%	96.01%	5,744,795	61.63%
Never	33,783,053	6,362,996	18.83%	21.88%	26.26%	27,420,057	81.17%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	2,180,972	664,548	30.47%	19.28%	41.66%	1,516,424	69.53%
30 to 34	2,642,586	811,954	30.73%	20.81%	40.64%	1,830,632	69.27%
35 to 39	3,020,012	1,177,595	38.99%	24.19%	53.80%	1,842,417	61.01%
40 to 44	4,465,609	2,094,617	46.91%	30.99%	62.82%	2,370,992	53.09%
45 to 49	6,840,162	2,858,596	41.79%	33.78%	49.81%	3,981,566	58.21%
50 to 54	5,693,359	1,887,666	33.16%	20.59%	45.73%	3,805,693	66.84%
55 or older	2,192,579	957,261	43.66%	27.98%	59.34%	1,235,318	56.34%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	36,274,207	5,660,727	15.61%	12.40%	18.81%	30,613,480	84.39%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	12,326,560	1,122,336	9.11%	4.74%	13.47%	11,204,224	90.89%
1	8,591,668	1,067,812	12.43%	6.60%	18.26%	7,523,856	87.57%
2	14,254,557	3,942,273	27.66%	22.21%	33.11%	10,312,284	72.34%
More than 2	28,325,067	9,987,980	35.26%	30.44%	40.08%	18,337,087	64.74%
Refused/Don't know	*	*	*	*	*	*	*
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	22,390,483	2,755,450	12.31%	9.49%	15.12%	19,635,033	87.69%
1	10,803,713	2,046,904	18.95%	14.47%	23.42%	8,756,809	81.05%
2	13,891,632	4,872,465	35.07%	30.65%	39.50%	9,019,166	64.93%
More than 2	16,226,054	6,404,360	39.47%	33.98%	44.96%	9,821,694	60.53%
Refused/Don't know	*	*	*	*	*	*	*
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	704,810	83,452	11.84%	(4.86%)	28.54%	621,358	88.16%
None	15,473,644	4,927,208	31.84%	25.73%	37.96%	10,546,436	68.16%
1	6,577,027	1,499,851	22.80%	14.30%	31.30%	5,077,176	77.20%
2	3,530,151	667,959	18.92%	12.43%	25.41%	2,862,192	81.08%
More than 2	1,789,085	469,914	26.27%	14.15%	38.38%	1,319,171	73.73%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	35,631,740	8,457,696	23.74%	20.54%	26.93%	27,174,044	76.26%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	15,202,557	1,584,247	10.42%	6.64%	14.20%	13,618,310	89.58%
Yes	7,349,801	2,648,494	36.03%	24.95%	47.12%	4,701,307	63.97%
No	41,015,264	11,859,894	28.92%	26.88%	30.95%	29,155,370	71.08%
Don't know	*	*	*	*	*	*	*
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	15,355,712	1,612,013	10.50%	6.73%	14.26%	13,743,699	89.50%
35 years old or younger	6,823,426	2,437,103	35.72%	24.75%	46.69%	4,386,323	64.28%
36 years or older	441,504	211,390	47.88%	8.89%	86.87%	230,114	52.12%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	41,015,264	11,859,894	28.92%	26.88%	30.95%	29,155,370	71.08%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	2,875,997	461,911	16.06%	3.82%	28.30%	2,414,086	83.94%
29 years old or younger	33,626,548	11,742,913	34.92%	31.85%	37.99%	21,883,635	65.08%
30 - 40 years old	2,819,868	847,019	30.04%	15.36%	44.72%	1,972,849	69.96%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	12,326,560	1,122,336	9.11%	4.74%	13.47%	11,204,224	90.89%
Only 1 live birth - see age at last live birth	11,642,383	1,922,469	16.51%	12.09%	20.93%	9,719,914	83.49%
No pregnancies resulting in live births	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	3,658,154	699,969	19.13%	7.85%	30.42%	2,958,185	80.87%
29 years old or younger	29,162,615	8,658,228	29.69%	27.19%	32.19%	20,504,387	70.31%
30 - 40 years old	16,829,708	5,068,967	30.12%	25.96%	34.28%	11,760,741	69.88%
41 years or older	1,159,139	477,670	41.21%	13.67%	68.75%	681,469	58.79%
Refused/Don't know	172,695	69,479	40.23%	17.00%	63.46%	103,216	59.77%
N/A - Never pregnant	12,326,560	1,122,336	9.11%	4.74%	13.47%	11,204,224	90.89%
No pregnancies resulting in live births	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	13,398,126	6,037,172	45.06%	37.96%	52.16%	7,360,953	54.94%
No	50,164,852	10,045,735	20.03%	17.34%	22.71%	40,119,117	79.97%
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	4,275,376	1,583,845	37.05%	28.96%	45.13%	2,691,531	62.95%
36 - 50 years old	7,299,884	3,583,144	49.08%	39.09%	59.08%	3,716,740	50.92%
51 years or older	1,482,342	787,468	53.12%	38.97%	67.27%	694,874	46.88%
Refused/Don't know	340,524	82,715	24.29%	1.10%	47.48%	257,809	75.71%
N/A - Did not have procedure	50,164,852	10,045,735	20.03%	17.34%	22.71%	40,119,117	79.97%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
Had ovaries removed	-	-	-	-	-	-	-
Missing	1,092,070	102,210	9.36%	(0.92%)	19.64%	989,860	90.64%
Yes	7,328,472	3,767,504	51.41%	39.33%	63.49%	3,560,968	48.59%
No	54,875,109	12,160,172	22.16%	19.27%	25.04%	42,714,937	77.84%
Don't know	425,126	90,515	21.29%	(3.50%)	46.08%	334,611	78.71%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	1,517,196	192,725	12.70%	0.90%	24.51%	1,324,471	87.30%
35 years old or younger	1,726,917	559,628	32.41%	15.49%	49.32%	1,167,289	67.59%
36 - 50 years old	3,982,739	2,516,219	63.18%	48.80%	77.56%	1,466,519	36.82%
51 years or older	1,529,540	669,899	43.80%	26.35%	61.24%	859,641	56.20%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	54,875,109	12,160,172	22.16%	19.27%	25.04%	42,714,937	77.84%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	63,720,777	16,120,401	25.30%	22.81%	27.78%	47,600,376	74.70%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	12,857,364	5,876,298	45.70%	38.06%	53.35%	6,981,066	54.30%
No	50,705,615	10,206,610	20.13%	17.49%	22.77%	40,499,004	79.87%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	61,324,726	15,244,554	24.86%	21.52%	28.19%	46,080,172	75.14%
Age at screening	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*
20-29	9,182,876	1,050,429	11.44%	7.17%	15.71%	8,132,448	88.56%
30-39	10,827,135	1,512,213	13.97%	8.64%	19.29%	9,314,922	86.03%
40-49	12,834,378	3,079,378	23.99%	15.71%	32.28%	9,755,000	76.01%
50-59	12,132,357	3,604,696	29.71%	21.75%	37.68%	8,527,661	70.29%
60-64	4,114,535	1,188,817	28.89%	18.79%	39.00%	2,925,719	71.11%
65-69	4,118,935	1,201,036	29.16%	18.46%	39.86%	2,917,899	70.84%
70-74	3,002,101	1,508,514	50.25%	41.00%	59.50%	1,493,587	49.75%
75-79	2,209,146	1,011,119	45.77%	32.86%	58.67%	1,198,027	54.23%
80+	2,781,244	1,056,735	38.00%	25.35%	50.64%	1,724,509	62.00%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	4,751,227	1,368,385	28.80%	23.75%	33.85%	3,382,842	71.20%
Other Hispanic	2,941,248	773,709	26.31%	20.29%	32.33%	2,167,540	73.69%
Non-Hispanic White	42,012,748	10,926,122	26.01%	21.71%	30.31%	31,086,626	73.99%
Non-Hispanic Black	7,332,322	1,539,393	20.99%	16.46%	25.53%	5,792,929	79.01%
Other Race - Including Multi-Racial	4,287,181	636,947	14.86%	8.15%	21.56%	3,650,235	85.14%
Education	-	-	-	-	-	-	-
Missing	1,546,877	520,135	33.62%	12.00%	55.25%	1,026,742	66.38%
Less Than 9th Grade	3,507,730	1,153,281	32.88%	23.32%	42.44%	2,354,449	67.12%
9-11th Grade	7,382,674	2,202,673	29.84%	21.74%	37.93%	5,180,000	70.16%
High School Grad/GED or Equivalent	48,831,529	11,348,885	23.24%	20.38%	26.10%	37,482,644	76.76%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Married	33,296,032	7,666,301	23.02%	18.69%	27.36%	25,629,731	76.98%
Widowed	5,662,311	2,413,928	42.63%	32.96%	52.30%	3,248,384	57.37%
Divorced	7,307,490	1,895,267	25.94%	19.17%	32.70%	5,412,223	74.06%
Separated	1,560,127	505,887	32.43%	12.08%	52.77%	1,054,240	67.57%
Never Married	8,320,591	1,638,849	19.70%	14.76%	24.63%	6,681,742	80.30%
Living with Partner	4,918,871	955,419	19.42%	11.82%	27.03%	3,963,452	80.58%
Refused	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,140,393	1,159,918	28.01%	19.81%	36.22%	2,980,475	71.99%
Poverty income ratio less than 1	9,383,987	2,838,689	30.25%	25.67%	34.83%	6,545,297	69.75%
Poverty income ratio 1 or greater	47,800,347	11,245,947	23.53%	19.97%	27.08%	36,554,399	76.47%
Body mass index	-	-	-	-	-	-	-
Missing	229,381	81,507	35.53%	(4.49%)	75.55%	147,874	64.47%
Normal: less than 25	23,484,279	3,160,969	13.46%	10.14%	16.78%	20,323,310	86.54%
Overweight: 25.0 - 29.9	17,118,171	4,367,857	25.52%	19.04%	31.99%	12,750,314	74.48%
Obese: 30 or higher	20,492,896	7,634,222	37.25%	32.73%	41.78%	12,858,674	62.75%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	61,324,726	15,244,554	24.86%	21.52%	28.19%	46,080,172	75.14%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,233,997	622,135	27.85%	19.55%	36.14%	1,611,862	72.15%
Glycohemoglobin < 5.7%	40,914,707	8,080,148	19.75%	16.92%	22.58%	32,834,559	80.25%
Glycohemoglobin 5.7% - 6.4%	14,351,734	4,763,152	33.19%	26.60%	39.78%	9,588,582	66.81%
Glycohemoglobin 6.5% or higher	3,824,288	1,779,119	46.52%	34.01%	59.04%	2,045,169	53.48%
Diabetes	-	-	-	-	-	-	-
Yes	5,248,803	2,761,401	52.61%	41.63%	63.59%	2,487,401	47.39%
No	56,075,924	12,483,153	22.26%	19.17%	25.35%	43,592,771	77.74%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	25,120,527	7,139,872	28.42%	23.49%	33.36%	17,980,655	71.58%
No	36,204,200	8,104,683	22.39%	18.39%	26.38%	28,099,517	77.61%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	17,374,564	7,202,646	41.46%	35.11%	47.80%	10,171,919	58.54%
No	43,921,653	8,041,909	18.31%	15.17%	21.45%	35,879,745	81.69%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	7,091,805	3,155,221	44.49%	34.55%	54.43%	3,936,585	55.51%
Mobility limitations	5,396,051	2,588,193	47.96%	39.42%	56.51%	2,807,858	52.04%
No mobility limitations	12,056,842	3,400,316	28.20%	20.88%	35.52%	8,656,526	71.80%
N/A - Respondent not asked mobility limitation questions	36,780,028	6,100,825	16.59%	13.27%	19.90%	30,679,203	83.41%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	4,002,837	1,895,680	47.36%	36.04%	58.67%	2,107,156	52.64%
No	57,321,890	13,348,874	23.29%	19.67%	26.90%	43,973,016	76.71%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	309,763	68,507	22.12%	(11.00%)	55.23%	241,256	77.88%
2 or more times a day	5,500,105	2,124,897	38.63%	27.47%	49.79%	3,375,208	61.37%
Once a day	4,580,303	1,165,160	25.44%	16.92%	33.96%	3,415,143	74.56%
2 or more time a week	4,802,662	1,920,037	39.98%	25.45%	54.51%	2,882,625	60.02%
Once a week	4,279,562	1,324,816	30.96%	21.13%	40.79%	2,954,745	69.04%
1-3 times a month	7,784,885	2,122,618	27.27%	19.56%	34.98%	5,662,267	72.73%
Never	34,050,166	6,518,520	19.14%	14.78%	23.51%	27,531,647	80.86%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	309,763	68,507	22.12%	(11.00%)	55.23%	241,256	77.88%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	4,279,562	1,324,816	30.96%	36.73%	109.10%	2,954,745	69.04%
1-3 times a month	7,784,885	2,122,618	27.27%	27.45%	83.82%	5,662,267	72.73%
Never	34,050,166	6,518,520	19.14%	20.63%	27.25%	27,531,647	80.86%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	61,324,726	15,244,554	24.86%	21.52%	28.19%	46,080,172	75.14%
Age of patient at last menstruation period	-	-	-	-	-	-	-
Missing	393,627	17,966	4.56%	(5.04%)	14.17%	375,661	95.44%
Younger than 30	1,885,261	715,877	37.97%	20.35%	55.59%	1,169,384	62.03%
30 to 34	2,334,729	877,327	37.58%	25.18%	49.97%	1,457,401	62.42%
35 to 39	2,973,028	956,227	32.16%	17.93%	46.40%	2,016,801	67.84%
40 to 44	4,188,179	1,708,308	40.79%	31.07%	50.51%	2,479,871	59.21%
45 to 49	5,551,597	2,038,965	36.73%	29.76%	43.70%	3,512,632	63.27%
50 to 54	7,510,861	2,802,318	37.31%	27.62%	47.00%	4,708,543	62.69%
55 or older	2,636,009	841,907	31.94%	19.21%	44.66%	1,794,102	68.06%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	33,604,604	5,202,384	15.48%	11.82%	19.14%	28,402,221	84.52%
Total number of pregnancies	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	13,728,142	1,414,386	10.30%	7.31%	13.30%	12,313,755	89.70%
1	7,722,862	1,541,865	19.96%	12.34%	27.59%	6,180,997	80.04%
2	13,807,785	3,219,980	23.32%	15.20%	31.44%	10,587,805	76.68%
More than 2	26,035,419	9,068,323	34.83%	30.30%	39.36%	16,967,096	65.17%
Number of vaginal deliveries	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	23,605,641	2,986,399	12.65%	9.77%	15.53%	20,619,242	87.35%
1	8,552,135	2,307,493	26.98%	17.22%	36.74%	6,244,642	73.02%
2	13,113,075	4,187,295	31.93%	21.35%	42.51%	8,925,779	68.07%
More than 2	15,938,346	5,719,506	35.89%	30.00%	41.77%	10,218,840	64.11%
Number of cesarean deliveries	-	-	-	-	-	-	-
Missing	352,535	141,483	40.13%	13.87%	66.39%	211,053	59.87%
None	14,692,704	4,826,984	32.85%	25.58%	40.13%	9,865,720	67.15%
1	5,736,418	1,282,999	22.37%	13.79%	30.95%	4,453,419	77.63%
2	3,377,938	789,384	23.37%	17.34%	29.40%	2,588,554	76.63%
More than 2	1,507,399	347,799	23.07%	7.71%	38.43%	1,159,600	76.93%
N/A - Vaginal delivery/deliveries only	35,657,733	7,855,906	22.03%	18.41%	25.65%	27,801,826	77.97%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-
Missing	17,112,302	1,971,708	11.52%	8.25%	14.79%	15,140,595	88.48%
Yes	8,462,133	2,912,553	34.42%	24.56%	44.28%	5,549,580	65.58%
No	35,547,194	10,271,527	28.90%	24.41%	33.38%	25,275,666	71.10%
Don't know	203,097	88,766	43.71%	8.02%	79.39%	114,331	56.29%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-
Missing	17,315,400	2,060,474	11.90%	8.56%	15.24%	15,254,926	88.10%
35 years old or younger	7,789,451	2,702,581	34.70%	24.28%	45.11%	5,086,871	65.30%
36 years or older	634,627	209,972	33.09%	17.09%	49.08%	424,654	66.91%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	35,547,194	10,271,527	28.90%	24.41%	33.38%	25,275,666	71.10%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	61,324,726	15,244,554	24.86%	21.52%	28.19%	46,080,172	75.14%
Age of patient at 1st live birth	-	-	-	-	-	-	-
Missing	3,384,161	557,322	16.47%	(1.60%)	34.53%	2,826,839	83.53%
29 years old or younger	31,512,241	10,167,428	32.27%	28.53%	36.00%	21,344,813	67.73%
30 - 40 years old	2,835,231	701,314	24.74%	15.53%	33.94%	2,133,917	75.26%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	13,728,142	1,414,386	10.30%	7.31%	13.30%	12,313,755	89.70%
Only 1 live birth - see age at last live birth	9,574,194	2,229,956	23.29%	15.50%	31.09%	7,344,238	76.71%
No pregnancies resulting in live births	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-
Missing	4,320,151	632,132	14.63%	0.04%	29.23%	3,688,019	85.37%
29 years old or younger	25,376,563	7,637,899	30.10%	25.76%	34.44%	17,738,664	69.90%
30 - 40 years old	16,177,542	4,926,139	30.45%	24.60%	36.30%	11,251,403	69.55%
41 years or older	1,341,703	382,962	28.54%	15.19%	41.90%	958,741	71.46%
Refused/Don't know	189,008	76,886	40.68%	(5.02%)	86.37%	112,122	59.32%
N/A - Never pregnant	13,728,142	1,414,386	10.30%	7.31%	13.30%	12,313,755	89.70%
No pregnancies resulting in live births	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	12,492,753	5,498,472	44.01%	36.63%	51.39%	6,994,282	55.99%
No	48,781,724	9,746,083	19.98%	16.44%	23.52%	39,035,641	80.02%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	3,552,051	1,575,761	44.36%	30.48%	58.24%	1,976,290	55.64%
36 - 50 years old	6,406,216	2,779,609	43.39%	33.88%	52.90%	3,626,607	56.61%
51 years or older	2,142,725	892,836	41.67%	28.80%	54.54%	1,249,889	58.33%
Refused/Don't know	391,761	250,266	63.88%	37.44%	90.33%	141,495	36.12%
N/A - Did not have procedure	48,781,724	9,746,083	19.98%	16.44%	23.52%	39,035,641	80.02%
Had ovaries removed	-	-	-	-	-	-	-
Missing	968,464	305,179	31.51%	12.88%	50.14%	663,285	68.49%
Yes	5,959,260	2,709,779	45.47%	36.49%	54.45%	3,249,481	54.53%
No	53,987,400	12,049,691	22.32%	19.03%	25.61%	41,937,709	77.68%
Refused	*	*	*	*	*	*	*
Don't know	398,051	179,905	45.20%	5.00%	85.39%	218,146	54.80%
Age when ovaries were removed	-	-	-	-	-	-	-
Missing	1,378,066	485,084	35.20%	15.30%	55.11%	892,982	64.80%
35 years old or younger	1,352,174	593,844	43.92%	25.98%	61.85%	758,330	56.08%
36 - 50 years old	2,947,862	1,474,066	50.00%	35.01%	65.00%	1,473,797	50.00%
51 years or older	1,474,095	499,347	33.87%	21.85%	45.90%	974,748	66.13%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	53,987,400	12,049,691	22.32%	19.03%	25.61%	41,937,709	77.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F04 - Prevalence of mixed urinary incontinence among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	61,324,726	15,244,554	24.86%	21.52%	28.19%	46,080,172	75.14%
***Had hysterectomy	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	12,313,977	5,482,562	44.52%	37.24%	51.81%	6,831,414	55.48%
No	48,960,501	9,761,992	19.94%	16.38%	23.50%	39,198,509	80.06%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.