

Table F05 - Prevalence of insensate urine loss among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Age at screening</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	*	*	*	*	*	*	*
20-29	12,825,480	415,755	3.24%	0.98%	5.51%	12,409,725	96.76%
30-39	18,943,946	1,128,840	5.96%	3.28%	8.63%	17,815,107	94.04%
40-49	22,500,594	1,842,607	8.19%	5.81%	10.57%	20,657,988	91.81%
50-59	17,533,327	1,854,103	10.57%	7.61%	13.54%	15,679,224	89.43%
60-64	5,474,602	613,297	11.20%	4.89%	17.52%	4,861,305	88.80%
65-69	5,272,171	1,211,096	22.97%	12.21%	33.73%	4,061,075	77.03%
70-74	4,310,387	633,916	14.71%	9.95%	19.47%	3,676,471	85.29%
75-79	4,298,734	934,115	21.73%	15.89%	27.57%	3,364,619	78.27%
80-84	2,888,291	670,865	23.23%	14.57%	31.89%	2,217,427	76.77%
85+	2,142,995	868,482	40.53%	16.46%	64.60%	1,274,513	59.47%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	6,044,522	466,028	7.71%	6.32%	9.10%	5,578,494	92.29%
Other Hispanic	5,721,649	624,075	10.91%	3.95%	17.86%	5,097,575	89.09%
Non-Hispanic White	69,144,500	7,728,283	11.18%	9.59%	12.77%	61,416,217	88.82%
Non-Hispanic Black	10,623,716	817,735	7.70%	5.19%	10.21%	9,805,981	92.30%
Other Race - Including Multi-Racial	4,925,900	573,081	11.63%	3.95%	19.31%	4,352,818	88.37%
<b>Education</b>	-	-	-	-	-	-	-
Missing	2,566,250	292,886	11.41%	0.97%	21.86%	2,273,364	88.59%
Less Than 9th Grade	6,983,938	1,103,900	15.81%	10.61%	21.00%	5,880,038	84.19%
9-11th Grade	11,493,555	1,441,527	12.54%	7.59%	17.50%	10,052,028	87.46%
High School Grad/GED or Equivalent	75,209,472	7,368,596	9.80%	8.33%	11.27%	67,840,876	90.20%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	606,658	70,897	11.69%	( 6.73%)	30.11%	535,761	88.31%
Married	56,394,435	5,132,815	9.10%	7.73%	10.48%	51,261,620	90.90%
Widowed	9,963,613	2,256,948	22.65%	16.45%	28.86%	7,706,666	77.35%
Divorced	9,618,647	1,223,384	12.72%	7.59%	17.85%	8,395,263	87.28%
Separated	3,664,817	515,018	14.05%	3.82%	24.29%	3,149,798	85.95%
Never Married	10,695,115	810,829	7.58%	4.21%	10.95%	9,884,287	92.42%
Living with Partner	5,458,082	175,177	3.21%	( 1.50%)	7.92%	5,282,905	96.79%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	6,220,599	571,455	9.19%	3.86%	14.51%	5,649,144	90.81%
Poverty income ratio less than 1	14,133,124	1,806,760	12.78%	8.96%	16.60%	12,326,364	87.22%
Poverty income ratio 1 or greater	76,106,564	7,830,987	10.29%	9.07%	11.51%	68,275,577	89.71%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	3,885,282	1,357,937	34.95%	21.30%	48.60%	2,527,345	65.05%
Normal: less than 25	36,632,098	2,397,610	6.55%	3.73%	9.36%	34,234,488	93.45%
Overweight: 25.0 - 29.9	25,530,785	2,411,701	9.45%	6.91%	11.98%	23,119,085	90.55%
Obese: 30 or higher	30,412,122	4,041,955	13.29%	10.34%	16.24%	26,370,168	86.71%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,329,836	544,549	16.35%	6.78%	25.92%	2,785,287	83.65%
Glycohemoglobin < 5.7%	75,071,947	6,417,332	8.55%	7.13%	9.97%	68,654,615	91.45%
Glycohemoglobin 5.7% - 6.4%	13,369,576	2,414,705	18.06%	12.90%	23.22%	10,954,871	81.94%
Glycohemoglobin 6.5% or higher	4,688,929	832,617	17.76%	9.51%	26.01%	3,856,313	82.24%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,416,102	1,368,017	21.32%	11.69%	30.96%	5,048,085	78.68%
No	90,044,185	8,841,185	9.82%	8.53%	11.11%	81,203,000	90.18%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,810,795	5,045,434	12.07%	9.47%	14.67%	36,765,361	87.93%
No	54,556,728	5,163,768	9.46%	7.16%	11.77%	49,392,960	90.54%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	27,368,244	4,981,130	18.20%	15.00%	21.40%	22,387,114	81.80%
No	68,702,093	5,137,322	7.48%	5.89%	9.07%	63,564,771	92.52%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	6,316,505	1,900,937	30.09%	22.58%	37.61%	4,415,568	69.91%
Mobility limitations	13,246,790	3,395,578	25.63%	17.19%	34.08%	9,851,212	74.37%
No mobility limitations	17,642,704	1,691,514	9.59%	6.08%	13.09%	15,951,190	90.41%
N/A - Respondent not asked mobility limitation questions	59,254,288	3,221,173	5.44%	3.68%	7.19%	56,033,115	94.56%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Younger than 30	3,837,297	558,722	14.56%	4.81%	24.31%	3,278,575	85.44%
30 to 34	3,723,416	892,121	23.96%	13.52%	34.40%	2,831,295	76.04%
35 to 39	4,762,950	1,082,167	22.72%	10.44%	35.00%	3,680,783	77.28%
40 to 44	5,719,731	946,416	16.55%	9.76%	23.33%	4,773,315	83.45%
45 to 49	10,167,841	1,540,105	15.15%	11.00%	19.29%	8,627,736	84.85%
50 to 54	9,442,873	1,443,291	15.28%	10.67%	19.90%	7,999,582	84.72%
55 or older	3,441,979	505,816	14.70%	5.50%	23.89%	2,936,163	85.30%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	55,153,584	3,110,803	5.64%	3.97%	7.31%	52,042,780	94.36%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-
Missing	1,404,672	225,704	16.07%	6.00%	26.13%	1,178,969	83.93%
None	15,771,213	785,906	4.98%	2.00%	7.96%	14,985,307	95.02%
1	10,550,872	873,893	8.28%	3.26%	13.30%	9,676,979	91.72%
2	18,643,815	1,699,897	9.12%	6.29%	11.94%	16,943,918	90.88%
More than 2	50,089,715	6,623,803	13.22%	11.43%	15.01%	43,465,912	86.78%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-
Missing	2,107,547	373,716	17.73%	8.00%	27.46%	1,733,831	82.27%
29 years old or younger	56,165,230	7,153,482	12.74%	11.06%	14.41%	49,011,748	87.26%
30 - 40 years old	3,432,833	382,069	11.13%	2.33%	19.93%	3,050,764	88.87%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	785,906	4.98%	2.00%	7.96%	14,985,307	95.02%
Only 1 live birth - see age at last live birth	15,380,871	1,229,049	7.99%	3.65%	12.33%	14,151,822	92.01%
No pregnancies resulting in live births	3,535,523	269,051	7.61%	4.80%	10.42%	3,266,472	92.39%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-
Missing	7,125,329	1,076,321	15.11%	10.08%	20.13%	6,049,008	84.89%
29 years old or younger	41,975,139	4,171,000	9.94%	7.90%	11.97%	37,804,139	90.06%
30 - 40 years old	25,834,321	3,289,445	12.73%	10.64%	14.83%	22,544,876	87.27%
41 years or older	2,115,817	617,481	29.18%	15.55%	42.82%	1,498,336	70.82%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	15,771,213	785,906	4.98%	2.00%	7.96%	14,985,307	95.02%
No pregnancies resulting in live births	3,535,523	269,051	7.61%	4.80%	10.42%	3,266,472	92.39%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	42,912,018	2,347,291	5.47%	3.64%	7.30%	40,564,728	94.53%
Yes	21,008,876	4,499,930	21.42%	16.00%	26.84%	16,508,946	78.58%
No	32,510,501	3,361,982	10.34%	8.47%	12.21%	29,148,518	89.66%
Don't know	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-
Missing	42,973,654	2,347,291	5.46%	3.63%	7.29%	40,626,364	94.54%
35 years old or younger	7,522,654	1,573,100	20.91%	11.37%	30.45%	5,949,554	79.09%
36 - 50 years old	10,892,423	2,194,473	20.15%	14.15%	26.14%	8,697,949	79.85%
51 years or older	2,336,250	645,998	27.65%	11.88%	43.42%	1,690,251	72.35%
Refused/Don't know	224,806	86,358	38.41%	7.13%	69.70%	138,448	61.59%
N/A - Did not have procedure	32,510,501	3,361,982	10.34%	8.47%	12.21%	29,148,518	89.66%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-
N/A for these years of data	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	96,460,287	10,209,202	10.58%	9.21%	11.95%	86,251,085	89.42%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	8,390,140	901,320	10.74%	4.10%	17.39%	7,488,820	89.26%
Yes	21,008,876	4,499,930	21.42%	16.00%	26.84%	16,508,946	78.58%
No	67,032,379	4,807,952	7.17%	5.88%	8.47%	62,224,427	92.83%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	696,430	0	0.00%	0.00%	0.00%	696,430	100.00%
20-29	12,963,783	263,027	2.03%	( 0.07%)	4.13%	12,700,756	97.97%
30-39	17,035,030	1,017,807	5.97%	1.99%	9.96%	16,017,223	94.03%
40-49	20,812,873	1,115,444	5.36%	2.24%	8.48%	19,697,428	94.64%
50-59	18,372,452	2,203,978	12.00%	7.91%	16.08%	16,168,474	88.00%
60-64	5,613,252	557,625	9.93%	4.11%	15.76%	5,055,626	90.07%
65-69	6,373,059	899,721	14.12%	7.55%	20.69%	5,473,338	85.88%
70-74	5,290,152	944,503	17.85%	8.51%	27.20%	4,345,649	82.15%
75-79	5,606,134	1,580,493	28.19%	19.43%	36.95%	4,025,641	71.81%
80-84	3,339,588	572,824	17.15%	6.05%	28.25%	2,766,764	82.85%
85+	2,125,548	639,643	30.09%	15.92%	44.26%	1,485,905	69.91%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	6,689,585	493,109	7.37%	5.07%	9.67%	6,196,475	92.63%
Other Hispanic	3,490,447	180,203	5.16%	( 1.10%)	11.43%	3,310,244	94.84%
Non-Hispanic White	71,779,337	7,792,030	10.86%	9.03%	12.68%	63,987,307	89.14%
Non-Hispanic Black	11,514,225	1,033,942	8.98%	5.65%	12.31%	10,480,283	91.02%
Other Race - Including Multi-Racial	4,754,706	295,781	6.22%	1.93%	10.52%	4,458,925	93.78%
<b>Education</b>	-	-	-	-	-	-	-
Missing	3,272,227	211,763	6.47%	2.03%	10.91%	3,060,464	93.53%
Less Than 9th Grade	7,084,184	1,333,289	18.82%	13.65%	23.99%	5,750,895	81.18%
9-11th Grade	10,222,174	1,388,920	13.59%	8.48%	18.70%	8,833,254	86.41%
High School Grad/GED or Equivalent	77,525,020	6,835,984	8.82%	7.32%	10.32%	70,689,037	91.18%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	538,429	61,946	11.50%	( 6.28%)	29.29%	476,483	88.50%
Married	55,195,562	5,049,762	9.15%	7.19%	11.10%	50,145,801	90.85%
Widowed	11,447,343	2,163,670	18.90%	12.92%	24.88%	9,283,673	81.10%
Divorced	11,142,010	1,261,828	11.32%	5.76%	16.89%	9,880,182	88.68%
Separated	2,410,096	89,084	3.70%	1.11%	6.28%	2,321,012	96.30%
Never Married	12,255,856	740,154	6.04%	3.75%	8.33%	11,515,702	93.96%
Living with Partner	5,186,193	428,621	8.26%	( 0.20%)	16.73%	4,757,572	91.74%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	5,148,576	488,212	9.48%	5.60%	13.36%	4,660,364	90.52%
Poverty income ratio less than 1	13,079,770	1,301,227	9.95%	6.87%	13.03%	11,778,543	90.05%
Poverty income ratio 1 or greater	79,999,954	8,005,627	10.01%	8.33%	11.68%	71,994,328	89.99%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	1,422,592	330,560	23.24%	6.21%	40.26%	1,092,031	76.76%
Normal: less than 25	36,253,820	3,003,922	8.29%	6.19%	10.38%	33,249,898	91.71%
Overweight: 25.0 - 29.9	27,727,409	2,588,365	9.34%	6.29%	12.38%	25,139,044	90.66%
Obese: 30 or higher	32,824,479	3,872,219	11.80%	8.90%	14.70%	28,952,261	88.20%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,306,134	324,055	9.80%	2.25%	17.36%	2,982,080	90.20%
Glycohemoglobin < 5.7%	75,244,125	6,051,022	8.04%	5.88%	10.20%	69,193,103	91.96%
Glycohemoglobin 5.7% - 6.4%	13,399,092	1,994,999	14.89%	10.68%	19.10%	11,404,093	85.11%
Glycohemoglobin 6.5% or higher	6,278,949	1,424,989	22.69%	15.71%	29.68%	4,853,960	77.31%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	8,142,154	1,873,337	23.01%	17.44%	28.58%	6,268,817	76.99%
No	90,086,145	7,921,728	8.79%	7.00%	10.59%	82,164,417	91.21%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	43,295,090	4,168,551	9.63%	7.25%	12.01%	39,126,538	90.37%
No	54,897,405	5,621,812	10.24%	8.51%	11.98%	49,275,593	89.76%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,594,115	4,702,215	15.37%	12.29%	18.45%	25,891,900	84.63%
No	67,061,499	5,014,484	7.48%	5.35%	9.61%	62,047,016	92.52%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	11,782,793	3,026,711	25.69%	20.26%	31.11%	8,756,083	74.31%
Mobility limitations	13,102,646	2,119,706	16.18%	13.18%	19.17%	10,982,940	83.82%
No mobility limitations	20,844,953	2,270,954	10.89%	8.55%	13.24%	18,573,999	89.11%
N/A - Respondent not asked mobility limitation questions	52,497,907	2,377,695	4.53%	2.63%	6.43%	50,120,212	95.47%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-
Missing	2,539,111	560,933	22.09%	8.77%	35.42%	1,978,179	77.91%
Younger than 30	2,891,152	342,915	11.86%	2.71%	21.01%	2,548,236	88.14%
30 to 34	4,500,186	917,553	20.39%	12.93%	27.84%	3,582,633	79.61%
35 to 39	5,581,951	1,025,173	18.37%	12.21%	24.53%	4,556,777	81.63%
40 to 44	6,423,717	981,417	15.28%	9.18%	21.38%	5,442,299	84.72%
45 to 49	10,508,053	1,526,930	14.53%	9.44%	19.62%	8,981,124	85.47%
50 to 54	9,282,520	1,433,048	15.44%	11.02%	19.86%	7,849,471	84.56%
55 or older	4,536,403	749,382	16.52%	9.13%	23.91%	3,787,022	83.48%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	51,728,762	2,234,157	4.32%	2.45%	6.18%	49,494,605	95.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-
Missing	1,870,938	481,048	25.71%	7.06%	44.37%	1,389,890	74.29%
None	17,857,228	1,126,090	6.31%	3.19%	9.42%	16,731,139	93.69%
1	13,698,789	1,003,647	7.33%	4.47%	10.18%	12,695,142	92.67%
2	22,315,868	1,488,337	6.67%	3.89%	9.45%	20,827,531	93.33%
More than 2	42,485,476	5,695,944	13.41%	10.52%	16.29%	36,789,532	86.59%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-
Missing	2,503,448	488,637	19.52%	4.73%	34.30%	2,014,811	80.48%
29 years old or younger	52,967,302	6,268,550	11.83%	9.48%	14.19%	46,698,752	88.17%
30 - 40 years old	4,126,416	268,754	6.51%	2.70%	10.33%	3,857,663	93.49%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	1,126,090	6.31%	3.19%	9.42%	16,731,139	93.69%
Only 1 live birth - see age at last live birth	16,014,683	1,442,083	9.00%	5.56%	12.45%	14,572,600	91.00%
No pregnancies resulting in live births	4,581,496	192,717	4.21%	0.56%	7.85%	4,388,779	95.79%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-
Missing	6,856,024	820,318	11.96%	5.04%	18.89%	6,035,706	88.04%
29 years old or younger	42,675,691	4,553,522	10.67%	8.34%	13.01%	38,122,169	89.33%
30 - 40 years old	24,296,503	2,887,669	11.89%	8.86%	14.91%	21,408,834	88.11%
41 years or older	1,773,198	184,327	10.40%	2.34%	18.45%	1,588,871	89.60%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,857,228	1,126,090	6.31%	3.19%	9.42%	16,731,139	93.69%
No pregnancies resulting in live births	4,581,496	192,717	4.21%	0.56%	7.85%	4,388,779	95.79%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	41,299,409	2,311,924	5.60%	3.26%	7.93%	38,987,485	94.40%
Yes	21,118,755	4,083,201	19.33%	15.43%	23.24%	17,035,554	80.67%
No	35,742,140	3,331,945	9.32%	7.03%	11.61%	32,410,195	90.68%
Don't know	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-
Missing	41,367,405	2,379,920	5.75%	3.54%	7.96%	38,987,485	94.25%
35 years old or younger	7,565,019	1,632,885	21.58%	13.00%	30.17%	5,932,134	78.42%
36 - 50 years old	10,924,810	1,484,894	13.59%	9.05%	18.13%	9,439,916	86.41%
51 years or older	2,435,004	941,864	38.68%	22.30%	55.06%	1,493,140	61.32%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	35,742,140	3,331,945	9.32%	7.03%	11.61%	32,410,195	90.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F05 - Prevalence of insensate urine loss among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-
N/A for these years of data	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	98,228,300	9,795,065	9.97%	8.29%	11.65%	88,433,234	90.03%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	1,940,125	481,048	24.79%	6.40%	43.18%	1,459,077	75.21%
Yes	21,118,755	4,083,201	19.33%	15.43%	23.24%	17,035,554	80.67%
No	75,101,424	5,162,821	6.87%	5.02%	8.73%	69,938,603	93.13%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	398,858	0	0.00%	0.00%	0.00%	398,858	100.00%
20-29	12,701,673	331,541	2.61%	0.01%	5.21%	12,370,132	97.39%
30-39	17,627,319	701,154	3.98%	1.94%	6.01%	16,926,165	96.02%
40-49	23,800,262	1,977,424	8.31%	4.47%	12.15%	21,822,839	91.69%
50-59	19,102,084	1,795,103	9.40%	4.70%	14.10%	17,306,981	90.60%
60-64	6,594,843	749,267	11.36%	4.96%	17.77%	5,845,576	88.64%
65-69	5,449,338	819,993	15.05%	6.97%	23.13%	4,629,345	84.95%
70-74	4,951,447	823,366	16.63%	8.93%	24.33%	4,128,081	83.37%
75-79	4,097,250	396,041	9.67%	2.24%	17.10%	3,701,210	90.33%
80-84	2,838,189	646,336	22.77%	16.26%	29.29%	2,191,853	77.23%
85+	1,982,190	350,980	17.71%	6.16%	29.26%	1,631,210	82.29%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	6,995,209	418,689	5.99%	3.25%	8.72%	6,576,520	94.01%
Other Hispanic	3,345,447	301,074	9.00%	1.04%	16.95%	3,044,374	91.00%
Non-Hispanic White	72,230,454	6,598,771	9.14%	7.25%	11.02%	65,631,682	90.86%
Non-Hispanic Black	11,495,230	859,810	7.48%	4.43%	10.53%	10,635,420	92.52%
Other Race - Including Multi-Racial	5,477,113	412,860	7.54%	1.37%	13.71%	5,064,253	92.46%
<b>Education</b>	-	-	-	-	-	-	-
Missing	2,515,028	128,011	5.09%	( 0.16%)	10.34%	2,387,017	94.91%
Less Than 9th Grade	5,390,275	855,100	15.86%	7.83%	23.90%	4,535,175	84.14%
9-11th Grade	11,067,500	1,365,111	12.33%	6.46%	18.21%	9,702,390	87.67%
High School Grad/GED or Equivalent	80,359,910	6,242,983	7.77%	5.81%	9.73%	74,116,928	92.23%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	745,740	0	0.00%	0.00%	0.00%	745,740	100.00%
Married	57,178,183	4,356,667	7.62%	5.61%	9.63%	52,821,516	92.38%
Widowed	8,886,413	1,284,310	14.45%	10.55%	18.35%	7,602,103	85.55%
Divorced	12,389,946	1,274,206	10.28%	6.11%	14.46%	11,115,740	89.72%
Separated	3,088,185	436,218	14.13%	2.83%	25.42%	2,651,967	85.87%
Never Married	10,375,951	777,819	7.50%	4.32%	10.67%	9,598,132	92.50%
Living with Partner	6,702,495	461,984	6.89%	1.82%	11.97%	6,240,511	93.11%
Refused	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	3,202,329	268,772	8.39%	0.58%	16.21%	2,933,557	91.61%
Poverty income ratio less than 1	10,867,636	1,120,508	10.31%	6.14%	14.48%	9,747,128	89.69%
Poverty income ratio 1 or greater	85,473,489	7,201,925	8.43%	6.56%	10.29%	78,271,564	91.57%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	922,023	207,838	22.54%	4.97%	40.11%	714,185	77.46%
Normal: less than 25	38,293,758	1,662,769	4.34%	2.51%	6.18%	36,630,989	95.66%
Overweight: 25.0 - 29.9	25,176,708	2,518,496	10.00%	6.23%	13.78%	22,658,212	90.00%
Obese: 30 or higher	35,150,965	4,202,102	11.95%	9.41%	14.50%	30,948,862	88.05%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,122,733	235,888	7.55%	( 1.14%)	16.25%	2,886,845	92.45%
Glycohemoglobin < 5.7%	75,014,443	5,515,343	7.35%	5.46%	9.24%	69,499,100	92.65%
Glycohemoglobin 5.7% - 6.4%	15,520,258	2,325,504	14.98%	10.16%	19.81%	13,194,755	85.02%
Glycohemoglobin 6.5% or higher	5,886,019	514,470	8.74%	3.53%	13.96%	5,371,549	91.26%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	8,450,575	1,461,519	17.29%	12.73%	21.86%	6,989,056	82.71%
No	91,092,878	7,129,686	7.83%	6.12%	9.53%	83,963,192	92.17%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,933,184	3,904,173	9.31%	5.92%	12.70%	38,029,011	90.69%
No	57,601,783	4,687,032	8.14%	6.16%	10.11%	52,914,751	91.86%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	31,026,049	4,391,598	14.15%	10.87%	17.44%	26,634,451	85.85%
No	68,249,544	4,199,606	6.15%	3.80%	8.50%	64,049,938	93.85%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	10,750,218	1,797,364	16.72%	10.02%	23.42%	8,952,854	83.28%
Mobility limitations	8,792,109	1,947,722	22.15%	13.27%	31.04%	6,844,387	77.85%
No mobility limitations	20,548,923	1,997,206	9.72%	7.96%	11.47%	18,551,716	90.28%
N/A - Respondent not asked mobility limitation questions	59,452,203	2,848,912	4.79%	2.96%	6.63%	56,603,291	95.21%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,664,983	994,987	14.93%	6.38%	23.48%	5,669,996	85.07%
No	92,800,223	7,583,603	8.17%	6.43%	9.92%	85,216,620	91.83%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	653,429	79,099	12.11%	( 2.52%)	26.73%	574,330	87.89%
2 or more times a day	11,136,447	1,624,460	14.59%	9.77%	19.41%	9,511,987	85.41%
Once a day	9,094,735	740,751	8.14%	4.06%	12.22%	8,353,984	91.86%
2 or more time a week	10,514,091	1,127,433	10.72%	5.16%	16.29%	9,386,658	89.28%
Once a week	6,191,809	509,854	8.23%	2.52%	13.95%	5,681,955	91.77%
1-3 times a month	13,475,962	1,585,059	11.76%	6.13%	17.40%	11,890,903	88.24%
Never	48,438,928	2,924,548	6.04%	3.98%	8.10%	45,514,380	93.96%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	653,429	79,099	12.11%	( 2.52%)	26.73%	574,330	87.89%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	10,514,091	1,127,433	10.72%	( 9.42%)	53.54%	9,386,658	89.28%
Once a week	*	*	*	*	*	*	*
1-3 times a month	13,475,962	1,585,059	11.76%	4.98%	41.20%	11,890,903	88.24%
Never	48,438,928	2,924,548	6.04%	6.72%	9.92%	45,514,380	93.96%
Don't know	*	*	*	*	*	*	*
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-
Missing	480,340	5,529	1.15%	( 0.47%)	2.77%	474,811	98.85%
Younger than 30	3,249,527	608,691	18.73%	10.19%	27.27%	2,640,836	81.27%
30 to 34	3,113,296	512,551	16.46%	3.32%	29.61%	2,600,745	83.54%
35 to 39	5,187,920	845,331	16.29%	10.34%	22.25%	4,342,589	83.71%
40 to 44	6,708,828	1,166,113	17.38%	9.34%	25.43%	5,542,714	82.62%
45 to 49	10,897,419	1,220,611	11.20%	7.77%	14.63%	9,676,809	88.80%
50 to 54	11,102,718	1,218,090	10.97%	6.09%	15.86%	9,884,628	89.03%
55 or older	3,608,524	409,726	11.35%	4.24%	18.47%	3,198,798	88.65%
Don't know	491,202	50,748	10.33%	( 2.50%)	23.17%	440,454	89.67%
N/A - Had period in last 12 months	54,703,679	2,553,814	4.67%	2.75%	6.59%	52,149,866	95.33%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,220,150	796,270	4.91%	1.84%	7.98%	15,423,880	95.09%
1	12,829,048	1,143,023	8.91%	4.43%	13.39%	11,686,025	91.09%
2	20,502,899	1,377,198	6.72%	3.13%	10.30%	19,125,701	93.28%
More than 2	49,937,378	5,274,713	10.56%	8.64%	12.49%	44,662,665	89.44%
Refused/Don't know	*	*	*	*	*	*	*
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-
Missing	993,141	36,421	3.67%	( 0.48%)	7.81%	956,720	96.33%
None	32,005,010	1,830,851	5.72%	3.18%	8.27%	30,174,158	94.28%
1	14,883,888	1,101,117	7.40%	3.11%	11.69%	13,782,771	92.60%
2	21,190,707	1,912,599	9.03%	5.33%	12.72%	19,278,108	90.97%
More than 2	30,447,679	3,710,217	12.19%	9.79%	14.58%	26,737,463	87.81%
Refused/Don't know	*	*	*	*	*	*	*
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-
Missing	1,016,170	36,421	3.58%	( 0.45%)	7.62%	979,749	96.42%
None	29,973,541	3,211,569	10.71%	8.67%	12.76%	26,761,972	89.29%
1	9,918,573	635,809	6.41%	1.00%	11.82%	9,282,765	93.59%
2	5,209,552	244,973	4.70%	0.14%	9.27%	4,964,579	95.30%
More than 2	1,827,082	177,546	9.72%	3.54%	15.90%	1,649,536	90.28%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	51,417,052	4,284,887	8.33%	5.74%	10.92%	47,132,165	91.67%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-
N/A for these years of data	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-
N/A for these years of data	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-
Missing	3,595,632	239,907	6.67%	( 0.26%)	13.60%	3,355,725	93.33%
29 years old or younger	55,606,083	5,817,235	10.46%	8.09%	12.83%	49,788,848	89.54%
30 - 40 years old	3,767,991	335,847	8.91%	2.83%	14.99%	3,432,145	91.09%
Refused/Don't know	288,861	12,615	4.37%	( 4.97%)	13.70%	276,247	95.63%
N/A - Never pregnant	16,220,150	796,270	4.91%	1.84%	7.98%	15,423,880	95.09%
Only 1 live birth - see age at last live birth	16,009,516	946,721	5.91%	2.92%	8.91%	15,062,794	94.09%
No pregnancies resulting in live births	4,055,220	442,610	10.91%	2.40%	19.43%	3,612,610	89.09%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-
Missing	3,639,681	239,907	6.59%	( 0.28%)	13.46%	3,399,774	93.41%
29 years old or younger	42,705,521	3,852,200	9.02%	6.19%	11.85%	38,853,321	90.98%
30 - 40 years old	31,105,003	3,133,784	10.07%	7.41%	12.74%	27,971,219	89.93%
41 years or older	1,331,628	113,819	8.55%	( 0.56%)	17.66%	1,217,809	91.45%
Refused/Don't know	486,249	12,615	2.59%	( 3.01%)	8.20%	473,635	97.41%
N/A - Never pregnant	16,220,150	796,270	4.91%	1.84%	7.98%	15,423,880	95.09%
No pregnancies resulting in live births	4,055,220	442,610	10.91%	2.40%	19.43%	3,612,610	89.09%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	44,433,141	2,214,911	4.98%	2.57%	7.40%	42,218,230	95.02%
Yes	20,930,164	3,111,137	14.86%	11.85%	17.88%	17,819,027	85.14%
No	34,146,780	3,265,157	9.56%	6.76%	12.36%	30,881,623	90.44%
Don't know	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-
Missing	44,466,510	2,214,911	4.98%	2.57%	7.39%	42,251,598	95.02%
35 years old or younger	6,450,871	1,256,247	19.47%	11.33%	27.61%	5,194,624	80.53%
36 - 50 years old	11,493,877	1,506,680	13.11%	8.80%	17.42%	9,987,197	86.89%
51 years or older	2,247,812	237,579	10.57%	0.36%	20.78%	2,010,233	89.43%
Refused/Don't know	737,604	110,630	15.00%	( 6.36%)	36.36%	626,973	85.00%
N/A - Did not have procedure	34,146,780	3,265,157	9.56%	6.76%	12.36%	30,881,623	90.44%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-
N/A for these years of data	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-
N/A for these years of data	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	99,543,453	8,591,205	8.63%	6.87%	10.39%	90,952,249	91.37%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,930,164	3,111,137	14.86%	11.85%	17.88%	17,819,027	85.14%
No	78,579,921	5,480,068	6.97%	5.01%	8.94%	73,099,853	93.03%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	723,602	0	0.00%	0.00%	0.00%	723,602	100.00%
20-29	12,424,789	668,375	5.38%	1.64%	9.12%	11,756,414	94.62%
30-39	17,248,808	785,340	4.55%	2.62%	6.49%	16,463,468	95.45%
40-49	20,998,655	1,473,057	7.02%	4.08%	9.95%	19,525,598	92.98%
50-59	20,032,917	1,831,130	9.14%	4.28%	14.00%	18,201,787	90.86%
60-64	7,609,867	531,525	6.98%	1.61%	12.36%	7,078,341	93.02%
65-69	6,578,135	1,142,855	17.37%	5.82%	28.93%	5,435,280	82.63%
70-74	5,285,665	796,825	15.08%	6.74%	23.41%	4,488,840	84.92%
75-79	4,488,645	798,084	17.78%	10.91%	24.65%	3,690,561	82.22%
80+	5,558,404	1,090,291	19.62%	12.95%	26.28%	4,468,113	80.38%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,557,995	507,189	6.71%	5.05%	8.37%	7,050,806	93.29%
Other Hispanic	4,955,929	400,231	8.08%	6.15%	10.00%	4,555,698	91.92%
Non-Hispanic White	71,245,854	7,117,719	9.99%	7.26%	12.72%	64,128,135	90.01%
Non-Hispanic Black	11,964,134	905,114	7.57%	5.55%	9.58%	11,059,019	92.43%
Other Race - Including Multi-Racial	5,225,573	187,229	3.58%	( 0.71%)	7.87%	5,038,345	96.42%
<b>Education</b>	-	-	-	-	-	-	-
Missing	2,863,940	261,913	9.15%	1.31%	16.98%	2,602,027	90.85%
Less Than 9th Grade	6,386,684	1,060,094	16.60%	9.50%	23.70%	5,326,590	83.40%
9-11th Grade	13,227,936	1,091,691	8.25%	5.06%	11.45%	12,136,245	91.75%
High School Grad/GED or Equivalent	78,459,310	6,703,785	8.54%	6.58%	10.51%	71,755,525	91.46%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	1,342,297	5,102	0.38%	( 0.41%)	1.17%	1,337,195	99.62%
Married	57,512,280	5,012,841	8.72%	5.88%	11.55%	52,499,438	91.28%
Widowed	9,335,424	1,467,915	15.72%	10.76%	20.69%	7,867,509	84.28%
Divorced	12,446,378	1,160,194	9.32%	6.28%	12.36%	11,286,184	90.68%
Separated	2,378,944	261,394	10.99%	4.55%	17.42%	2,117,550	89.01%
Never Married	11,745,069	872,982	7.43%	3.10%	11.77%	10,872,087	92.57%
Living with Partner	6,162,712	337,055	5.47%	0.30%	10.64%	5,825,657	94.53%
Refused	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	6,989,889	691,218	9.89%	3.76%	16.01%	6,298,672	90.11%
Poverty income ratio less than 1	13,770,564	1,309,416	9.51%	6.85%	12.17%	12,461,148	90.49%
Poverty income ratio 1 or greater	80,189,033	7,116,849	8.88%	6.69%	11.06%	73,072,184	91.12%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	940,501	143,985	15.31%	1.02%	29.60%	796,516	84.69%
Normal: less than 25	35,488,858	1,988,235	5.60%	3.33%	7.87%	33,500,623	94.40%
Overweight: 25.0 - 29.9	29,208,740	2,599,311	8.90%	6.35%	11.45%	26,609,429	91.10%
Obese: 30 or higher	35,311,387	4,385,952	12.42%	7.02%	17.82%	30,925,435	87.58%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,604,052	496,522	13.78%	4.50%	23.06%	3,107,530	86.22%
Glycohemoglobin < 5.7%	68,091,107	4,357,411	6.40%	4.80%	8.00%	63,733,696	93.60%
Glycohemoglobin 5.7% - 6.4%	22,307,896	2,927,731	13.12%	8.68%	17.57%	19,380,165	86.88%
Glycohemoglobin 6.5% or higher	6,946,431	1,335,819	19.23%	7.67%	30.79%	5,610,612	80.77%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	9,397,306	1,751,046	18.63%	5.81%	31.45%	7,646,259	81.37%
No	91,552,180	7,366,436	8.05%	6.76%	9.34%	84,185,744	91.95%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	41,106,905	3,955,072	9.62%	5.70%	13.54%	37,151,833	90.38%
No	59,778,399	5,162,411	8.64%	6.56%	10.71%	54,615,988	91.36%
Refused	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	32,659,207	4,939,492	15.12%	11.41%	18.84%	27,719,715	84.88%
No	68,206,695	4,177,990	6.13%	4.45%	7.80%	64,028,705	93.87%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	11,577,303	1,628,578	14.07%	8.38%	19.75%	9,948,725	85.93%
Mobility limitations	10,685,927	2,081,480	19.48%	9.49%	29.47%	8,604,447	80.52%
No mobility limitations	20,236,932	2,426,943	11.99%	8.81%	15.17%	17,809,988	88.01%
N/A - Respondent not asked mobility limitation questions	58,449,325	2,980,481	5.10%	3.85%	6.35%	55,468,844	94.90%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,361,300	1,662,416	22.58%	11.05%	34.12%	5,698,884	77.42%
No	93,498,807	7,455,067	7.97%	6.61%	9.33%	86,043,741	92.03%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	583,024	53,256	9.13%	( 6.69%)	24.95%	529,767	90.87%
2 or more times a day	9,899,256	1,117,761	11.29%	6.56%	16.02%	8,781,494	88.71%
Once a day	9,630,096	677,767	7.04%	3.11%	10.97%	8,952,328	92.96%
2 or more time a week	7,557,510	949,718	12.57%	6.38%	18.75%	6,607,792	87.43%
Once a week	7,022,920	978,226	13.93%	6.96%	20.90%	6,044,694	86.07%
1-3 times a month	14,356,537	1,580,930	11.01%	4.79%	17.24%	12,775,607	88.99%
Never	51,673,706	3,727,438	7.21%	5.31%	9.12%	47,946,268	92.79%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	583,024	53,256	9.13%	( 6.69%)	24.95%	529,767	90.87%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	7,557,510	949,718	12.57%	1.97%	71.60%	6,607,792	87.43%
Once a week	*	*	*	*	*	*	*
1-3 times a month	14,356,537	1,580,930	11.01%	20.63%	62.82%	12,775,607	88.99%
Never	51,673,706	3,727,438	7.21%	6.79%	10.07%	47,946,268	92.79%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-
Missing	432,110	0	0.00%	0.00%	0.00%	432,110	100.00%
Younger than 30	3,054,284	423,228	13.86%	( 0.89%)	28.61%	2,631,056	86.14%
30 to 34	3,701,673	383,865	10.37%	1.60%	19.14%	3,317,808	89.63%
35 to 39	5,041,353	525,354	10.42%	2.94%	17.91%	4,515,999	89.58%
40 to 44	7,483,484	920,906	12.31%	6.48%	18.13%	6,562,578	87.69%
45 to 49	11,591,015	1,593,743	13.75%	8.94%	18.56%	9,997,272	86.25%
50 to 54	10,312,194	1,513,834	14.68%	8.55%	20.81%	8,798,360	85.32%
55 or older	5,101,078	763,634	14.97%	5.85%	24.09%	4,337,444	85.03%
Don't know	210,595	0	0.00%	0.00%	0.00%	210,595	100.00%
N/A - Had period in last 12 months	54,021,700	2,992,920	5.54%	4.06%	7.02%	51,028,781	94.46%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	16,821,230	1,207,501	7.18%	3.99%	10.37%	15,613,729	92.82%
1	13,330,257	1,087,943	8.16%	3.72%	12.60%	12,242,314	91.84%
2	22,645,536	1,423,890	6.29%	3.48%	9.09%	21,221,646	93.71%
More than 2	47,871,914	5,398,149	11.28%	8.45%	14.10%	42,473,765	88.72%
Refused/Don't know	*	*	*	*	*	*	*
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-
Missing	674,220	47,585	7.06%	( 5.26%)	19.38%	626,636	92.94%
None	31,904,825	2,265,188	7.10%	5.08%	9.12%	29,639,637	92.90%
1	17,719,654	1,170,086	6.60%	4.62%	8.59%	16,549,568	93.40%
2	22,981,603	1,685,041	7.33%	3.61%	11.06%	21,296,562	92.67%
More than 2	27,654,864	3,949,583	14.28%	10.04%	18.53%	23,705,281	85.72%
Refused/Don't know	*	*	*	*	*	*	*
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-
Missing	964,977	47,585	4.93%	( 2.32%)	12.19%	917,393	95.07%
None	25,236,974	2,594,369	10.28%	7.03%	13.53%	22,642,605	89.72%
1	9,920,178	840,156	8.47%	3.88%	13.06%	9,080,022	91.53%
2	5,518,794	304,810	5.52%	1.25%	9.80%	5,213,985	94.48%
More than 2	2,981,345	240,883	8.08%	0.33%	15.83%	2,740,462	91.92%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	56,312,898	5,089,680	9.04%	6.43%	11.65%	51,223,218	90.96%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-
Missing	21,396,140	1,628,516	7.61%	4.93%	10.29%	19,767,624	92.39%
Yes	13,130,418	1,383,386	10.54%	5.52%	15.55%	11,747,032	89.46%
No	66,243,210	6,088,701	9.19%	6.26%	12.12%	60,154,508	90.81%
Don't know	179,719	16,879	9.39%	(11.72%)	30.50%	162,840	90.61%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-
Missing	21,575,858	1,645,395	7.63%	4.94%	10.32%	19,930,463	92.37%
35 years old or younger	12,104,834	1,359,049	11.23%	5.90%	16.55%	10,745,785	88.77%
36 years or older	932,800	24,336	2.61%	( 2.96%)	8.18%	908,463	97.39%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	66,243,210	6,088,701	9.19%	6.26%	12.12%	60,154,508	90.81%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-
Missing	4,574,909	421,015	9.20%	2.56%	15.84%	4,153,894	90.80%
29 years old or younger	55,782,517	5,727,469	10.27%	7.65%	12.88%	50,055,048	89.73%
30 - 40 years old	5,156,343	316,294	6.13%	1.08%	11.19%	4,840,049	93.87%
41 years or older	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	16,821,230	1,207,501	7.18%	3.99%	10.37%	15,613,729	92.82%
Only 1 live birth - see age at last live birth	18,059,987	1,394,439	7.72%	4.02%	11.42%	16,665,548	92.28%
No pregnancies resulting in live births	491,126	50,764	10.34%	1.38%	19.29%	440,362	89.66%
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-
Missing	5,807,618	497,540	8.57%	2.17%	14.96%	5,310,079	91.43%
29 years old or younger	46,253,779	3,900,864	8.43%	4.95%	11.92%	42,352,915	91.57%
30 - 40 years old	29,451,221	3,248,452	11.03%	8.49%	13.57%	26,202,769	88.97%
41 years or older	1,776,911	140,867	7.93%	( 0.39%)	16.25%	1,636,044	92.07%
Refused/Don't know	347,601	71,494	20.57%	0.45%	40.68%	276,106	79.43%
N/A - Never pregnant	16,821,230	1,207,501	7.18%	3.99%	10.37%	15,613,729	92.82%
No pregnancies resulting in live births	491,126	50,764	10.34%	1.38%	19.29%	440,362	89.66%
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,797,312	3,164,286	14.52%	9.43%	19.60%	18,633,025	85.48%
No	78,994,375	5,953,196	7.54%	6.12%	8.95%	73,041,180	92.46%
Don't know	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	6,116,146	537,281	8.78%	( 0.15%)	17.71%	5,578,865	91.22%
36 - 50 years old	12,694,892	2,064,040	16.26%	10.69%	21.82%	10,630,852	83.74%
51 years or older	2,470,853	539,538	21.84%	8.78%	34.89%	1,931,315	78.16%
Refused/Don't know	515,421	23,428	4.55%	( 3.74%)	12.83%	491,993	95.45%
N/A - Did not have procedure	78,994,375	5,953,196	7.54%	6.12%	8.95%	73,041,180	92.46%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-
Missing	1,803,240	71,844	3.98%	0.83%	7.14%	1,731,396	96.02%
Yes	11,803,029	2,094,025	17.74%	11.57%	23.92%	9,709,004	82.26%
No	86,682,478	6,913,187	7.98%	6.27%	9.68%	79,769,290	92.02%
Don't know	660,739	38,426	5.82%	( 5.61%)	17.24%	622,313	94.18%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-
Missing	2,463,979	110,270	4.48%	0.72%	8.23%	2,353,709	95.52%
35 years old or younger	2,315,265	290,687	12.56%	( 0.32%)	25.43%	2,024,577	87.44%
36 - 50 years old	6,648,470	1,112,276	16.73%	9.93%	23.53%	5,536,194	83.27%
51 years or older	2,575,084	678,554	26.35%	10.75%	41.95%	1,896,530	73.65%
Refused/Don't know	264,211	12,508	4.73%	( 5.30%)	14.77%	251,703	95.27%
N/A - Did not have procedure	86,682,478	6,913,187	7.98%	6.27%	9.68%	79,769,290	92.02%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	100,949,486	9,117,482	9.03%	7.13%	10.93%	91,832,004	90.97%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,220,500	3,134,924	14.77%	9.60%	19.95%	18,085,577	85.23%
No	79,571,187	5,982,559	7.52%	6.10%	8.94%	73,588,628	92.48%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	97,614,302	9,192,844	9.42%	7.77%	11.07%	88,421,458	90.58%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	353,109	0	0.00%	0.00%	0.00%	353,109	100.00%
20-29	12,072,924	519,643	4.30%	1.32%	7.29%	11,553,282	95.70%
30-39	15,950,025	752,154	4.72%	2.49%	6.94%	15,197,871	95.28%
40-49	19,711,852	1,842,589	9.35%	5.64%	13.06%	17,869,263	90.65%
50-59	20,704,188	1,536,416	7.42%	3.37%	11.48%	19,167,772	92.58%
60-64	7,767,115	928,762	11.96%	6.18%	17.74%	6,838,353	88.04%
65-69	6,636,364	1,052,909	15.87%	7.94%	23.79%	5,583,455	84.13%
70-74	5,111,072	749,231	14.66%	10.43%	18.89%	4,361,841	85.34%
75-79	4,127,134	743,102	18.01%	11.62%	24.39%	3,384,032	81.99%
80+	5,180,518	1,068,038	20.62%	9.58%	31.65%	4,112,480	79.38%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,365,117	564,696	7.67%	5.52%	9.82%	6,800,422	92.33%
Other Hispanic	4,864,518	181,284	3.73%	1.81%	5.64%	4,683,235	96.27%
Non-Hispanic White	67,795,316	7,326,980	10.81%	8.62%	12.99%	60,468,336	89.19%
Non-Hispanic Black	11,428,205	865,608	7.57%	3.60%	11.55%	10,562,597	92.43%
Other Race - Including Multi-Racial	6,161,145	254,277	4.13%	1.97%	6.28%	5,906,868	95.87%
<b>Education</b>	-	-	-	-	-	-	-
Missing	2,266,489	339,384	14.97%	3.30%	26.65%	1,927,105	85.03%
Less Than 9th Grade	5,789,901	598,173	10.33%	6.17%	14.49%	5,191,728	89.67%
9-11th Grade	11,482,376	966,081	8.41%	4.93%	11.90%	10,516,295	91.59%
High School Grad/GED or Equivalent	77,874,180	7,274,457	9.34%	7.61%	11.07%	70,599,723	90.66%
Refused	*	*	*	*	*	*	*
Don't Know	156,713	14,748	9.41%	( 7.20%)	26.02%	141,965	90.59%
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	353,109	0	0.00%	0.00%	0.00%	353,109	100.00%
Married	55,880,303	5,484,160	9.81%	8.10%	11.53%	50,396,144	90.19%
Widowed	8,791,969	1,492,588	16.98%	7.81%	26.14%	7,299,381	83.02%
Divorced	11,116,682	910,403	8.19%	4.63%	11.75%	10,206,280	91.81%
Separated	2,211,003	225,238	10.19%	3.95%	16.42%	1,985,764	89.81%
Never Married	11,937,752	804,039	6.74%	3.52%	9.95%	11,133,713	93.26%
Living with Partner	7,120,836	276,416	3.88%	1.59%	6.17%	6,844,420	96.12%
Refused	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	6,365,743	597,493	9.39%	3.92%	14.85%	5,768,250	90.61%
Poverty income ratio less than 1	13,898,370	1,339,274	9.64%	5.96%	13.31%	12,559,096	90.36%
Poverty income ratio 1 or greater	77,350,189	7,256,077	9.38%	7.85%	10.91%	70,094,112	90.62%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	432,161	69,955	16.19%	( 3.09%)	35.47%	362,206	83.81%
Normal: less than 25	34,081,290	2,314,853	6.79%	4.82%	8.76%	31,766,436	93.21%
Overweight: 25.0 - 29.9	27,223,096	2,461,339	9.04%	5.91%	12.17%	24,761,758	90.96%
Obese: 30 or higher	35,877,755	4,346,697	12.12%	8.86%	15.37%	31,531,058	87.88%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	97,614,302	9,192,844	9.42%	7.77%	11.07%	88,421,458	90.58%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,219,727	325,483	10.11%	5.98%	14.24%	2,894,244	89.89%
Glycohemoglobin < 5.7%	62,554,629	4,793,325	7.66%	6.23%	9.10%	57,761,304	92.34%
Glycohemoglobin 5.7% - 6.4%	25,095,519	3,294,171	13.13%	10.74%	15.51%	21,801,348	86.87%
Glycohemoglobin 6.5% or higher	6,744,427	779,864	11.56%	7.21%	15.92%	5,964,563	88.44%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	8,781,262	1,378,672	15.70%	10.57%	20.83%	7,402,590	84.30%
No	88,833,040	7,814,171	8.80%	7.24%	10.35%	81,018,868	91.20%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	39,090,118	4,035,294	10.32%	8.20%	12.44%	35,054,824	89.68%
No	58,524,184	5,157,549	8.81%	6.59%	11.04%	53,366,634	91.19%
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	30,112,832	4,334,527	14.39%	11.19%	17.60%	25,778,305	85.61%
No	67,440,102	4,858,317	7.20%	5.39%	9.02%	62,581,785	92.80%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	11,367,903	1,817,520	15.99%	11.79%	20.18%	9,550,383	84.01%
Mobility limitations	9,501,130	1,773,451	18.67%	12.48%	24.85%	7,727,679	81.33%
No mobility limitations	20,673,589	2,479,846	12.00%	7.88%	16.11%	18,193,743	88.00%
N/A - Respondent not asked mobility limitation questions	56,071,679	3,122,026	5.57%	4.39%	6.74%	52,949,653	94.43%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,026,447	1,383,609	19.69%	11.53%	27.85%	5,642,838	80.31%
No	90,511,916	7,740,209	8.55%	6.97%	10.13%	82,771,708	91.45%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	472,607	133,835	28.32%	0.26%	56.38%	338,771	71.68%
2 or more times a day	9,005,246	1,517,780	16.85%	10.43%	23.28%	7,487,466	83.15%
Once a day	8,275,588	754,464	9.12%	5.21%	13.03%	7,521,124	90.88%
2 or more time a week	8,103,086	1,035,574	12.78%	8.09%	17.47%	7,067,512	87.22%
Once a week	6,115,640	349,564	5.72%	1.95%	9.48%	5,766,076	94.28%
1-3 times a month	13,225,964	1,599,420	12.09%	7.74%	16.45%	11,626,543	87.91%
Never	52,342,337	3,802,206	7.26%	5.60%	8.93%	48,540,131	92.74%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	472,607	133,835	28.32%	( 0.72%)	51.26%	338,771	71.68%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	8,103,086	1,035,574	12.78%	(11.40%)	55.34%	7,067,512	87.22%
Once a week	6,115,640	349,564	5.72%	(11.24%)	66.58%	5,766,076	94.28%
1-3 times a month	13,225,964	1,599,420	12.09%	1.73%	23.22%	11,626,543	87.91%
Never	52,342,337	3,802,206	7.26%	7.65%	10.65%	48,540,131	92.74%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	97,614,302	9,192,844	9.42%	7.77%	11.07%	88,421,458	90.58%
<b>Age of patient at last menstruation period</b>	-	-	-	-	-	-	-
Missing	624,425	0	0.00%	0.00%	0.00%	624,425	100.00%
Younger than 30	2,882,766	617,157	21.41%	11.92%	30.90%	2,265,609	78.59%
30 to 34	3,241,045	527,249	16.27%	6.14%	26.39%	2,713,796	83.73%
35 to 39	4,541,323	538,743	11.86%	4.56%	19.17%	4,002,580	88.14%
40 to 44	7,445,396	1,088,292	14.62%	7.53%	21.71%	6,357,104	85.38%
45 to 49	9,772,191	1,330,479	13.61%	9.59%	17.64%	8,441,712	86.39%
50 to 54	13,734,798	1,339,020	9.75%	6.32%	13.18%	12,395,778	90.25%
55 or older	5,095,598	886,363	17.39%	11.10%	23.69%	4,209,234	82.61%
Don't know	*	*	*	*	*	*	*
N/A - Had period in last 12 months	50,018,311	2,865,540	5.73%	4.02%	7.43%	47,152,772	94.27%
<b>Total number of pregnancies</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
None	17,526,617	789,721	4.51%	2.15%	6.86%	16,736,896	95.49%
1	12,038,159	889,117	7.39%	3.39%	11.38%	11,149,042	92.61%
2	23,689,401	2,162,913	9.13%	5.96%	12.30%	21,526,488	90.87%
More than 2	44,255,771	5,351,092	12.09%	10.21%	13.97%	38,904,679	87.91%
<b>Number of vaginal deliveries</b>	-	-	-	-	-	-	-
Missing	310,780	29,321	9.43%	( 7.43%)	26.30%	281,459	90.57%
None	32,847,722	2,223,632	6.77%	5.02%	8.52%	30,624,091	93.23%
1	14,225,353	1,038,297	7.30%	2.55%	12.05%	13,187,056	92.70%
2	23,367,578	2,330,719	9.97%	6.21%	13.73%	21,036,858	90.03%
More than 2	26,862,869	3,570,874	13.29%	10.69%	15.89%	23,291,994	86.71%
<b>Number of cesarean deliveries</b>	-	-	-	-	-	-	-
Missing	779,924	29,321	3.76%	( 3.74%)	11.26%	750,603	96.24%
None	25,073,953	2,556,782	10.20%	6.33%	14.06%	22,517,172	89.80%
1	9,753,274	778,210	7.98%	3.81%	12.15%	8,975,064	92.02%
2	5,288,336	522,394	9.88%	4.16%	15.59%	4,765,942	90.12%
More than 2	2,314,074	365,035	15.77%	1.07%	30.48%	1,949,039	84.23%
N/A - Vaginal delivery/deliveries only	54,404,741	4,941,102	9.08%	6.46%	11.71%	49,463,638	90.92%
<b>Gave birth to any babies 9lb or more</b>	-	-	-	-	-	-	-
Missing	22,928,465	1,293,065	5.64%	2.78%	8.50%	21,635,401	94.36%
Yes	14,458,970	1,482,274	10.25%	6.47%	14.03%	12,976,696	89.75%
No	59,898,920	6,371,929	10.64%	8.64%	12.64%	53,526,991	89.36%
Don't know	327,946	45,576	13.90%	( 1.66%)	29.45%	282,370	86.10%
<b>Age of patient when she had 9 lb or more baby</b>	-	-	-	-	-	-	-
Missing	23,256,411	1,338,640	5.76%	2.90%	8.61%	21,917,771	94.24%
35 years old or younger	13,118,409	1,241,612	9.46%	5.83%	13.10%	11,876,797	90.54%
36 years or older	1,284,225	222,381	17.32%	3.84%	30.79%	1,061,844	82.68%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	59,898,920	6,371,929	10.64%	8.64%	12.64%	53,526,991	89.36%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.



Table F05 - Prevalence of insensate urine loss among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	97,614,302	9,192,844	9.42%	7.77%	11.07%	88,421,458	90.58%
<b>Age of patient at 1st live birth</b>	-	-	-	-	-	-	-
Missing	5,401,849	503,344	9.32%	0.83%	17.80%	4,898,505	90.68%
29 years old or younger	53,902,824	6,119,339	11.35%	8.84%	13.86%	47,783,485	88.65%
30 - 40 years old	5,231,391	778,628	14.88%	9.04%	20.72%	4,452,763	85.12%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Never pregnant	17,526,617	789,721	4.51%	2.15%	6.86%	16,736,896	95.49%
Only 1 live birth - see age at last live birth	15,122,928	1,001,812	6.62%	3.03%	10.22%	14,121,115	93.38%
No pregnancies resulting in live births	*	*	*	*	*	*	*
<b>Age of patient at last live birth</b>	-	-	-	-	-	-	-
Missing	6,887,500	586,694	8.52%	1.16%	15.88%	6,300,806	91.48%
29 years old or younger	41,061,749	4,052,100	9.87%	7.12%	12.62%	37,009,650	90.13%
30 - 40 years old	29,435,328	3,470,428	11.79%	9.59%	13.99%	25,964,900	88.21%
41 years or older	2,166,264	211,824	9.78%	1.93%	17.63%	1,954,440	90.22%
Refused/Don't know	207,290	82,077	39.60%	2.11%	77.08%	125,213	60.40%
N/A - Never pregnant	17,526,617	789,721	4.51%	2.15%	6.86%	16,736,896	95.49%
No pregnancies resulting in live births	*	*	*	*	*	*	*
<b>**Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,381,540	3,258,347	15.99%	13.56%	18.42%	17,123,193	84.01%
No	77,182,512	5,934,497	7.69%	6.05%	9.33%	71,248,016	92.31%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Age at time of hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
35 years old or younger	5,352,637	1,006,766	18.81%	11.51%	26.11%	4,345,871	81.19%
36 - 50 years old	10,930,520	1,716,547	15.70%	11.52%	19.89%	9,213,974	84.30%
51 years or older	3,681,428	399,246	10.84%	4.23%	17.46%	3,282,182	89.16%
Refused/Don't know	416,955	135,788	32.57%	11.70%	53.43%	281,167	67.43%
N/A - Did not have procedure	77,182,512	5,934,497	7.69%	6.05%	9.33%	71,248,016	92.31%
<b>Had ovaries removed</b>	-	-	-	-	-	-	-
Missing	1,954,499	134,418	6.88%	0.32%	13.43%	1,820,081	93.12%
Yes	10,081,522	1,476,026	14.64%	9.88%	19.40%	8,605,496	85.36%
No	84,842,800	7,445,076	8.78%	7.25%	10.30%	77,397,724	91.22%
Refused	*	*	*	*	*	*	*
Don't know	723,930	137,324	18.97%	( 6.21%)	44.15%	586,606	81.03%
<b>Age when ovaries were removed</b>	-	-	-	-	-	-	-
Missing	2,689,979	271,742	10.10%	2.15%	18.05%	2,418,238	89.90%
35 years old or younger	2,196,357	387,825	17.66%	6.26%	29.06%	1,808,531	82.34%
36 - 50 years old	5,209,369	805,858	15.47%	7.25%	23.69%	4,403,511	84.53%
51 years or older	2,452,490	204,982	8.36%	3.95%	12.76%	2,247,508	91.64%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Did not have procedure	84,842,800	7,445,076	8.78%	7.25%	10.30%	77,397,724	91.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05 - Prevalence of insensate urine loss among women, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	97,614,302	9,192,844	9.42%	7.77%	11.07%	88,421,458	90.58%
<b>***Had hysterectomy</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	20,181,982	3,242,970	16.07%	13.63%	18.51%	16,939,011	83.93%
No	77,382,071	5,949,873	7.69%	6.06%	9.32%	71,432,198	92.31%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.046: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine without an activity like coughing, lifting, or exercise, or an urge to urinate?

NOTE: \*\*The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: \*\*\* Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.