

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	415,755	63,956	15.38%	147,235	35.41%	56,268	13.53%	148,296	35.67%	0	0.00%	0	0.00%
30-39	1,128,840	202,822	17.97%	241,396	21.38%	11,158	0.99%	673,464	59.66%	0	0.00%	0	0.00%
40-49	1,842,607	262,603	14.25%	189,056	10.26%	475,717	25.82%	915,230	49.67%	0	0.00%	0	0.00%
50-59	1,854,103	604,715	32.61%	520,927	28.10%	310,563	16.75%	417,898	22.54%	0	0.00%	0	0.00%
60-64	613,297	159,828	26.06%	229,283	37.39%	183,334	29.89%	40,852	6.66%	0	0.00%	0	0.00%
65-69	1,211,096	301,525	24.90%	367,913	30.38%	210,017	17.34%	331,641	27.38%	0	0.00%	0	0.00%
70-74	633,916	305,669	48.22%	153,195	24.17%	78,780	12.43%	96,272	15.19%	0	0.00%	0	0.00%
75-79	934,115	452,827	48.48%	295,579	31.64%	95,398	10.21%	49,300	5.28%	41,011	4.39%	0	0.00%
80-84	670,865	330,212	49.22%	110,595	16.49%	172,852	25.77%	27,533	4.10%	29,674	4.42%	0	0.00%
85+	868,482	391,668	45.10%	108,715	12.52%	320,008	36.85%	48,090	5.54%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	466,028	91,717	19.68%	143,888	30.88%	123,109	26.42%	107,314	23.03%	0	0.00%	0	0.00%
Other Hispanic	624,075	261,851	41.96%	247,693	39.69%	0	0.00%	114,530	18.35%	0	0.00%	0	0.00%
Non-Hispanic White	7,728,283	2,319,251	30.01%	1,755,763	22.72%	1,521,591	19.69%	2,060,993	26.67%	70,685	0.91%	0	0.00%
Non-Hispanic Black	817,735	307,673	37.63%	156,268	19.11%	197,587	24.16%	156,207	19.10%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	292,886	40,165	13.71%	143,808	49.10%	31,102	10.62%	77,810	26.57%	0	0.00%	0	0.00%
Less Than 9th Grade	1,103,900	493,268	44.68%	137,409	12.45%	369,961	33.51%	62,251	5.64%	41,011	3.72%	0	0.00%
9-11th Grade	1,441,527	346,405	24.03%	559,364	38.80%	284,508	19.74%	251,250	17.43%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	7,368,596	2,216,308	30.08%	1,523,312	20.67%	1,226,231	16.64%	2,373,071	32.21%	29,674	0.40%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	5,132,815	1,245,622	24.27%	1,309,066	25.50%	1,188,166	23.15%	1,389,960	27.08%	0	0.00%	0	0.00%
Widowed	2,256,948	1,134,510	50.27%	449,856	19.93%	386,079	17.11%	215,818	9.56%	70,685	3.13%	0	0.00%
Divorced	1,223,384	414,906	33.91%	272,586	22.28%	150,821	12.33%	385,071	31.48%	0	0.00%	0	0.00%
Separated	515,018	70,908	13.77%	165,871	32.21%	68,758	13.35%	209,481	40.67%	0	0.00%	0	0.00%
Never Married	810,829	205,407	25.33%	142,379	17.56%	74,169	9.15%	388,873	47.96%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	571,455	175,886	30.78%	195,788	34.26%	47,838	8.37%	110,932	19.41%	41,011	7.18%	0	0.00%
Poverty income ratio less than 1	1,806,760	685,767	37.96%	368,129	20.38%	419,824	23.24%	333,040	18.43%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	7,830,987	2,234,493	28.53%	1,799,976	22.99%	1,446,434	18.47%	2,320,410	29.63%	29,674	0.38%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,357,937	743,121	54.72%	307,604	22.65%	91,601	6.75%	185,937	13.69%	29,674	2.19%	0	0.00%
Normal: less than 25	2,397,610	417,064	17.39%	290,816	12.13%	658,446	27.46%	1,031,285	43.01%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	2,411,701	594,292	24.64%	666,287	27.63%	435,432	18.05%	674,678	27.98%	41,011	1.70%	0	0.00%
Obese: 30 or higher	4,041,955	1,341,668	33.19%	1,099,187	27.19%	728,618	18.03%	872,482	21.59%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	544,549	179,627	32.99%	95,219	17.49%	109,639	20.13%	160,064	29.39%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	6,417,332	1,668,720	26.00%	1,439,040	22.42%	1,184,952	18.46%	2,094,946	32.65%	29,674	0.46%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	2,414,705	1,052,762	43.60%	564,290	23.37%	451,899	18.71%	345,754	14.32%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	832,617	195,037	23.42%	265,345	31.87%	167,606	20.13%	163,618	19.65%	41,011	4.93%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,368,017	524,132	38.31%	320,790	23.45%	276,494	20.21%	205,590	15.03%	41,011	3.00%	0	0.00%
No	8,841,185	2,572,014	29.09%	2,043,103	23.11%	1,637,602	18.52%	2,558,792	28.94%	29,674	0.34%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,045,434	1,499,347	29.72%	1,362,937	27.01%	827,598	16.40%	1,355,552	26.87%	0	0.00%	0	0.00%
No	5,163,768	1,596,798	30.92%	1,000,957	19.38%	1,086,498	21.04%	1,408,830	27.28%	70,685	1.37%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	4,981,130	1,928,145	38.71%	1,033,703	20.75%	1,116,565	22.42%	902,717	18.12%	0	0.00%	0	0.00%
No	5,137,322	1,168,001	22.74%	1,330,191	25.89%	797,532	15.52%	1,770,915	34.47%	70,685	1.38%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,900,937	760,880	40.03%	409,065	21.52%	432,108	22.73%	257,873	13.57%	41,011	2.16%	0	0.00%
Mobility limitations	3,395,578	1,321,997	38.93%	792,813	23.35%	755,335	22.24%	495,759	14.60%	29,674	0.87%	0	0.00%
No mobility limitations	1,691,514	340,111	20.11%	571,912	33.81%	269,018	15.90%	510,474	30.18%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	3,221,173	673,158	20.90%	590,105	18.32%	457,635	14.21%	1,500,276	46.58%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	558,722	93,080	16.66%	94,437	16.90%	144,494	25.86%	226,711	40.58%	0	0.00%	0	0.00%
30 to 34	892,121	289,129	32.41%	75,972	8.52%	277,596	31.12%	249,423	27.96%	0	0.00%	0	0.00%
35 to 39	1,082,167	452,425	41.81%	306,041	28.28%	222,666	20.58%	71,361	6.59%	29,674	2.74%	0	0.00%
40 to 44	946,416	287,468	30.37%	177,270	18.73%	140,776	14.87%	299,891	31.69%	41,011	4.33%	0	0.00%
45 to 49	1,540,105	561,163	36.44%	399,556	25.94%	365,467	23.73%	213,919	13.89%	0	0.00%	0	0.00%
50 to 54	1,443,291	673,050	46.63%	293,470	20.33%	175,762	12.18%	301,008	20.86%	0	0.00%	0	0.00%
55 or older	505,816	141,045	27.88%	206,160	40.76%	135,508	26.79%	23,103	4.57%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	3,110,803	574,901	18.48%	795,057	25.56%	451,826	14.52%	1,289,019	41.44%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	225,704	55,867	24.75%	25,620	11.35%	45,496	20.16%	98,720	43.74%	0	0.00%	0	0.00%
None	785,906	95,723	12.18%	89,447	11.38%	171,518	21.82%	429,218	54.61%	0	0.00%	0	0.00%
1	873,893	151,035	17.28%	298,541	34.16%	243,846	27.90%	180,471	20.65%	0	0.00%	0	0.00%
2	1,699,897	436,368	25.67%	445,749	26.22%	365,842	21.52%	422,264	24.84%	29,674	1.75%	0	0.00%
More than 2	6,623,803	2,357,153	35.59%	1,504,536	22.71%	1,087,394	16.42%	1,633,709	24.66%	41,011	0.62%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	373,716	63,493	16.99%	25,620	6.86%	77,461	20.73%	207,141	55.43%	0	0.00%	0	0.00%
29 years old or younger	7,153,482	2,518,720	35.21%	1,654,965	23.14%	1,295,247	18.11%	1,643,538	22.98%	41,011	0.57%	0	0.00%
30 - 40 years old	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	785,906	95,723	12.18%	89,447	11.38%	171,518	21.82%	429,218	54.61%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	1,229,049	226,683	18.44%	360,675	29.35%	240,682	19.58%	371,335	30.21%	29,674	2.41%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,076,321	356,231	33.10%	268,138	24.91%	108,240	10.06%	314,038	29.18%	29,674	2.76%	0	0.00%
29 years old or younger	4,171,000	1,195,757	28.67%	1,075,265	25.78%	733,870	17.59%	1,166,108	27.96%	0	0.00%	0	0.00%
30 - 40 years old	3,289,445	1,167,706	35.50%	712,048	21.65%	749,036	22.77%	619,644	18.84%	41,011	1.25%	0	0.00%
41 years or older	617,481	185,867	30.10%	69,661	11.28%	133,964	21.70%	227,988	36.92%	0	0.00%	0	0.00%
N/A - Never pregnant	785,906	95,723	12.18%	89,447	11.38%	171,518	21.82%	429,218	54.61%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,347,291	501,968	21.38%	540,186	23.01%	412,089	17.56%	893,048	38.05%	0	0.00%	0	0.00%
Yes	4,499,930	1,718,500	38.19%	929,911	20.67%	934,061	20.76%	887,784	19.73%	29,674	0.66%	0	0.00%
No	3,361,982	875,678	26.05%	893,797	26.59%	567,946	16.89%	983,550	29.26%	41,011	1.22%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,347,291	501,968	21.38%	540,186	23.01%	412,089	17.56%	893,048	38.05%	0	0.00%	0	0.00%
35 years old or younger	1,573,100	534,559	33.98%	279,499	17.77%	260,623	16.57%	498,419	31.68%	0	0.00%	0	0.00%
36 - 50 years old	2,194,473	919,761	41.91%	538,552	24.54%	382,682	17.44%	323,803	14.76%	29,674	1.35%	0	0.00%
51 years or older	645,998	264,179	40.89%	111,860	17.32%	231,930	35.90%	38,029	5.89%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	3,361,982	875,678	26.05%	893,797	26.59%	567,946	16.89%	983,550	29.26%	41,011	1.22%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	901,320	267,097	29.63%	178,442	19.80%	90,808	10.08%	364,974	40.49%	0	0.00%	0	0.00%
Yes	4,499,930	1,718,500	38.19%	929,911	20.67%	934,061	20.76%	887,784	19.73%	29,674	0.66%	0	0.00%
No	4,807,952	1,110,549	23.10%	1,255,541	26.11%	889,227	18.49%	1,511,624	31.44%	41,011	0.85%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	1,017,807	262,971	25.84%	201,815	19.83%	132,733	13.04%	420,287	41.29%	0	0.00%	0	0.00%
40-49	1,115,444	170,188	15.26%	169,673	15.21%	192,380	17.25%	583,205	52.28%	0	0.00%	0	0.00%
50-59	2,203,978	436,239	19.79%	829,578	37.64%	327,398	14.85%	610,763	27.71%	0	0.00%	0	0.00%
60-64	557,625	175,339	31.44%	140,083	25.12%	126,432	22.67%	115,771	20.76%	0	0.00%	0	0.00%
65-69	899,721	327,693	36.42%	374,488	41.62%	46,260	5.14%	151,280	16.81%	0	0.00%	0	0.00%
70-74	944,503	336,867	35.67%	271,753	28.77%	274,286	29.04%	61,597	6.52%	0	0.00%	0	0.00%
75-79	1,580,493	658,139	41.64%	492,471	31.16%	202,790	12.83%	227,092	14.37%	0	0.00%	0	0.00%
80-84	572,824	240,790	42.04%	263,704	46.04%	64,004	11.17%	4,327	0.76%	0	0.00%	0	0.00%
85+	639,643	366,456	57.29%	153,169	23.95%	26,655	4.17%	93,363	14.60%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	493,109	210,116	42.61%	168,297	34.13%	56,258	11.41%	58,439	11.85%	0	0.00%	0	0.00%
Other Hispanic	*	*		*		*		*		*		*	
Non-Hispanic White	7,792,030	2,365,889	30.36%	2,335,602	29.97%	1,083,583	13.91%	2,006,956	25.76%	0	0.00%	0	0.00%
Non-Hispanic Black	1,033,942	271,260	26.24%	285,095	27.57%	225,558	21.82%	252,029	24.38%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	211,763	143,381	67.71%	24,311	11.48%	0	0.00%	44,070	20.81%	0	0.00%	0	0.00%
Less Than 9th Grade	1,333,289	366,687	27.50%	617,824	46.34%	185,838	13.94%	162,940	12.22%	0	0.00%	0	0.00%
9-11th Grade	1,388,920	609,445	43.88%	257,586	18.55%	171,443	12.34%	350,447	25.23%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	6,835,984	1,927,320	28.19%	2,030,847	29.71%	1,042,063	15.24%	1,835,753	26.85%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	5,049,762	1,413,176	27.99%	1,428,565	28.29%	933,965	18.50%	1,274,056	25.23%	0	0.00%	0	0.00%
Widowed	2,163,670	898,700	41.54%	831,409	38.43%	204,462	9.45%	229,099	10.59%	0	0.00%	0	0.00%
Divorced	1,261,828	356,173	28.23%	412,830	32.72%	93,459	7.41%	399,366	31.65%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	740,154	175,123	23.66%	115,121	15.55%	83,025	11.22%	366,885	49.57%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	488,212	250,781	51.37%	165,778	33.96%	0	0.00%	71,653	14.68%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,301,227	401,050	30.82%	430,101	33.05%	185,207	14.23%	284,869	21.89%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	8,005,627	2,413,064	30.14%	2,338,047	29.21%	1,217,827	15.21%	2,036,689	25.44%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	330,560	197,747	59.82%	87,555	26.49%	27,180	8.22%	18,078	5.47%	0	0.00%	0	0.00%
Normal: less than 25	3,003,922	944,723	31.45%	787,122	26.20%	432,896	14.41%	839,181	27.94%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	2,588,365	631,289	24.39%	1,024,706	39.59%	354,171	13.68%	578,198	22.34%	0	0.00%	0	0.00%
Obese: 30 or higher	3,872,219	1,291,136	33.34%	1,034,543	26.72%	588,786	15.21%	957,754	24.73%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	324,055	105,115	32.44%	112,984	34.87%	3,690	1.14%	102,266	31.56%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	6,051,022	1,842,118	30.44%	1,680,496	27.77%	986,199	16.30%	1,542,209	25.49%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,994,999	683,175	34.24%	605,196	30.34%	265,540	13.31%	441,088	22.11%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,424,989	434,487	30.49%	535,250	37.56%	147,604	10.36%	307,648	21.59%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,873,337	756,535	40.38%	603,482	32.21%	215,695	11.51%	297,626	15.89%	0	0.00%	0	0.00%
No	7,921,728	2,308,360	29.14%	2,330,445	29.42%	1,187,339	14.99%	2,095,585	26.45%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,168,551	1,422,965	34.14%	1,050,041	25.19%	684,080	16.41%	1,011,465	24.26%	0	0.00%	0	0.00%
No	5,621,812	1,637,227	29.12%	1,883,885	33.51%	718,954	12.79%	1,381,745	24.58%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,702,215	1,677,474	35.67%	1,604,730	34.13%	754,203	16.04%	665,807	14.16%	0	0.00%	0	0.00%
No	5,014,484	1,387,421	27.67%	1,299,367	25.91%	600,293	11.97%	1,727,403	34.45%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	3,026,711	1,376,814	45.49%	694,650	22.95%	337,631	11.16%	617,616	20.41%	0	0.00%	0	0.00%
Mobility limitations	2,119,706	625,529	29.51%	801,756	37.82%	377,516	17.81%	314,905	14.86%	0	0.00%	0	0.00%
No mobility limitations	2,270,954	721,761	31.78%	831,872	36.63%	299,972	13.21%	417,349	18.38%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	2,377,695	340,791	14.33%	605,648	25.47%	387,915	16.31%	1,043,341	43.88%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	560,933	337,336	60.14%	101,004	18.01%	22,746	4.06%	99,847	17.80%	0	0.00%	0	0.00%
Younger than 30	*	*	*	*	*	*	*	*	*	*	*	*	*
30 to 34	917,553	340,593	37.12%	259,912	28.33%	80,527	8.78%	236,522	25.78%	0	0.00%	0	0.00%
35 to 39	1,025,173	519,395	50.66%	356,782	34.80%	123,508	12.05%	25,488	2.49%	0	0.00%	0	0.00%
40 to 44	981,417	416,225	42.41%	364,719	37.16%	92,555	9.43%	107,918	11.00%	0	0.00%	0	0.00%
45 to 49	1,526,930	308,769	20.22%	489,370	32.05%	143,541	9.40%	585,250	38.33%	0	0.00%	0	0.00%
50 to 54	1,433,048	553,489	38.62%	332,783	23.22%	315,448	22.01%	231,328	16.14%	0	0.00%	0	0.00%
55 or older	749,382	128,290	17.12%	227,639	30.38%	223,933	29.88%	169,519	22.62%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	2,234,157	338,781	15.16%	610,659	27.33%	400,775	17.94%	883,942	39.56%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	481,048	337,336	70.13%	101,004	21.00%	22,746	4.73%	19,962	4.15%	0	0.00%	0	0.00%
None	1,126,090	275,125	24.43%	266,717	23.69%	133,709	11.87%	450,538	40.01%	0	0.00%	0	0.00%
1	1,003,647	175,033	17.44%	419,009	41.75%	271,785	27.08%	137,819	13.73%	0	0.00%	0	0.00%
2	1,488,337	308,122	20.70%	539,218	36.23%	382,291	25.69%	258,705	17.38%	0	0.00%	0	0.00%
More than 2	5,695,944	1,969,279	34.57%	1,607,978	28.23%	592,501	10.40%	1,526,186	26.79%	0	0.00%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	488,637	337,336	69.04%	101,004	20.67%	22,746	4.65%	27,551	5.64%	0	0.00%	0	0.00%
29 years old or younger	6,268,550	2,095,778	33.43%	1,843,265	29.40%	817,184	13.04%	1,512,323	24.13%	0	0.00%	0	0.00%
30 - 40 years old	268,754	48,811	18.16%	64,727	24.08%	48,537	18.06%	106,678	39.69%	0	0.00%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,126,090	275,125	24.43%	266,717	23.69%	133,709	11.87%	450,538	40.01%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	1,442,083	284,273	19.71%	489,069	33.91%	372,621	25.84%	296,120	20.53%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	820,318	531,245	64.76%	130,721	15.94%	87,577	10.68%	70,774	8.63%	0	0.00%	0	0.00%
29 years old or younger	4,553,522	1,332,670	29.27%	1,662,867	36.52%	731,727	16.07%	826,258	18.15%	0	0.00%	0	0.00%
30 - 40 years old	2,887,669	835,130	28.92%	671,902	23.27%	391,799	13.57%	988,838	34.24%	0	0.00%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,126,090	275,125	24.43%	266,717	23.69%	133,709	11.87%	450,538	40.01%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,209,202	3,096,146	30.33%	2,363,894	23.15%	1,914,096	18.75%	2,764,382	27.08%	70,685	0.69%	0	0.00%
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,311,924	550,562	23.81%	514,694	22.26%	416,918	18.03%	829,750	35.89%	0	0.00%	0	0.00%
Yes	4,083,201	1,631,109	39.95%	1,266,698	31.02%	539,605	13.22%	645,789	15.82%	0	0.00%	0	0.00%
No	3,331,945	878,522	26.37%	1,152,534	34.59%	383,217	11.50%	917,672	27.54%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,379,920	555,264	23.33%	514,694	21.63%	480,212	20.18%	829,750	34.86%	0	0.00%	0	0.00%
35 years old or younger	1,632,885	636,750	39.00%	572,918	35.09%	133,297	8.16%	289,919	17.76%	0	0.00%	0	0.00%
36 - 50 years old	1,484,894	641,526	43.20%	425,012	28.62%	239,492	16.13%	178,865	12.05%	0	0.00%	0	0.00%
51 years or older	941,864	329,275	34.96%	268,768	28.54%	166,816	17.71%	177,005	18.79%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	3,331,945	878,522	26.37%	1,152,534	34.59%	383,217	11.50%	917,672	27.54%	0	0.00%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,795,065	3,064,895	31.29%	2,933,926	29.95%	1,403,033	14.32%	2,393,210	24.43%	0	0.00%	0	0.00%
***Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	481,048	337,336	70.13%	101,004	21.00%	22,746	4.73%	19,962	4.15%	0	0.00%	0	0.00%
Yes	4,083,201	1,631,109	39.95%	1,266,698	31.02%	539,605	13.22%	645,789	15.82%	0	0.00%	0	0.00%
No	5,162,821	1,091,748	21.15%	1,566,225	30.34%	777,389	15.06%	1,727,460	33.46%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	331,541	0	0.00%	59,169	17.85%	85,303	25.73%	187,069	56.42%	0	0.00%	0	0.00%
30-39	701,154	57,050	8.14%	112,821	16.09%	441,007	62.90%	90,276	12.88%	0	0.00%	0	0.00%
40-49	1,977,424	539,152	27.27%	288,352	14.58%	454,400	22.98%	695,520	35.17%	0	0.00%	0	0.00%
50-59	1,795,103	124,630	6.94%	169,839	9.46%	796,016	44.34%	685,875	38.21%	18,743	1.04%	0	0.00%
60-64	749,267	290,528	38.77%	112,463	15.01%	99,930	13.34%	246,347	32.88%	0	0.00%	0	0.00%
65-69	819,993	165,343	20.16%	126,091	15.38%	505,585	61.66%	22,975	2.80%	0	0.00%	0	0.00%
70-74	823,366	235,912	28.65%	135,561	16.46%	202,422	24.58%	249,470	30.30%	0	0.00%	0	0.00%
75-79	396,041	58,390	14.74%	91,248	23.04%	159,700	40.32%	34,980	8.83%	51,724	13.06%	0	0.00%
80-84	646,336	230,835	35.71%	219,327	33.93%	196,174	30.35%	0	0.00%	0	0.00%	0	0.00%
85+	350,980	156,663	44.64%	63,833	18.19%	103,893	29.60%	26,591	7.58%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	418,689	7,343	1.75%	100,142	23.92%	135,078	32.26%	157,382	37.59%	18,743	4.48%	0	0.00%
Other Hispanic	*	*		*		*		*		*		*	
Non-Hispanic White	6,598,771	1,553,717	23.55%	817,492	12.39%	2,299,343	34.85%	1,876,495	28.44%	51,724	0.78%	0	0.00%
Non-Hispanic Black	859,810	117,287	13.64%	349,167	40.61%	263,480	30.64%	129,875	15.11%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	855,100	65,539	7.66%	259,204	30.31%	322,587	37.73%	189,026	22.11%	18,743	2.19%	0	0.00%
9-11th Grade	1,365,111	498,955	36.55%	143,538	10.51%	590,491	43.26%	80,403	5.89%	51,724	3.79%	0	0.00%
High School Grad/GED or Equivalent	6,242,983	1,189,285	19.05%	969,381	15.53%	2,114,643	33.87%	1,969,673	31.55%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Married	4,356,667	803,426	18.44%	550,039	12.63%	1,928,370	44.26%	1,023,108	23.48%	51,724	1.19%	0	0.00%
Widowed	1,284,310	462,499	36.01%	373,588	29.09%	338,742	26.38%	109,481	8.52%	0	0.00%	0	0.00%
Divorced	1,274,206	373,840	29.34%	180,377	14.16%	275,727	21.64%	444,263	34.87%	0	0.00%	0	0.00%
Separated	436,218	30,079	6.90%	97,294	22.30%	54,180	12.42%	235,921	54.08%	18,743	4.30%	0	0.00%
Never Married	777,819	36,585	4.70%	109,859	14.12%	276,315	35.52%	355,060	45.65%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	1,120,508	261,253	23.32%	275,848	24.62%	425,872	38.01%	138,792	12.39%	18,743	1.67%	0	0.00%
Poverty income ratio 1 or greater	7,201,925	1,597,250	22.18%	960,792	13.34%	2,504,642	34.78%	2,087,518	28.99%	51,724	0.72%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,662,769	343,816	20.68%	84,493	5.08%	662,926	39.87%	519,810	31.26%	51,724	3.11%	0	0.00%
Overweight: 25.0 - 29.9	2,518,496	481,711	19.13%	388,016	15.41%	815,451	32.38%	833,318	33.09%	0	0.00%	0	0.00%
Obese: 30 or higher	4,202,102	895,940	21.32%	884,570	21.05%	1,523,498	36.26%	879,351	20.93%	18,743	0.45%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	5,515,343	1,330,106	24.12%	561,030	10.17%	2,002,578	36.31%	1,621,628	29.40%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	2,325,504	528,397	22.72%	647,112	27.83%	742,893	31.95%	336,635	14.48%	70,467	3.03%	0	0.00%
Glycohemoglobin 6.5% or higher	514,470	0	0.00%	146,461	28.47%	186,941	36.34%	181,068	35.20%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,461,519	311,016	21.28%	472,147	32.31%	497,288	34.03%	181,068	12.39%	0	0.00%	0	0.00%
No	7,129,686	1,547,487	21.70%	906,556	12.72%	2,547,142	35.73%	2,058,034	28.87%	70,467	0.99%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,904,173	652,929	16.72%	634,246	16.25%	1,403,035	35.94%	1,213,962	31.09%	0	0.00%	0	0.00%
No	4,687,032	1,205,574	25.72%	744,457	15.88%	1,641,395	35.02%	1,025,140	21.87%	70,467	1.50%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,391,598	1,265,889	28.83%	824,906	18.78%	1,191,890	27.14%	1,108,914	25.25%	0	0.00%	0	0.00%
No	4,199,606	592,614	14.11%	553,797	13.19%	1,852,540	44.11%	1,130,189	26.91%	70,467	1.68%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,797,364	665,103	37.00%	430,355	23.94%	406,757	22.63%	295,150	16.42%	0	0.00%	0	0.00%
Mobility limitations	1,947,722	485,158	24.91%	421,488	21.64%	551,841	28.33%	470,492	24.16%	18,743	0.96%	0	0.00%
No mobility limitations	1,997,206	340,708	17.06%	207,106	10.37%	1,073,602	53.76%	324,066	16.23%	51,724	2.59%	0	0.00%
N/A - Respondent not asked mobility limitation questions	2,848,912	367,533	12.90%	319,755	11.22%	1,012,230	35.53%	1,149,394	40.35%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	994,987	234,491	23.57%	412,155	41.42%	236,418	23.76%	111,923	11.25%	0	0.00%	0	0.00%
No	7,583,603	1,611,397	21.25%	966,548	12.75%	2,808,012	37.03%	2,127,179	28.05%	70,467	0.93%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	1,624,460	294,885	18.15%	302,707	18.63%	632,978	38.97%	393,890	24.25%	0	0.00%	0	0.00%
Once a day	740,751	38,305	5.17%	151,285	20.42%	400,458	54.06%	150,703	20.34%	0	0.00%	0	0.00%
2 or more time a week	1,127,433	172,823	15.33%	113,920	10.10%	383,716	34.03%	438,231	38.87%	18,743	1.66%	0	0.00%
Once a week	509,854	114,730	22.50%	52,108	10.22%	235,545	46.20%	107,472	21.08%	0	0.00%	0	0.00%
1-3 times a month	1,585,059	267,350	16.87%	181,552	11.45%	668,465	42.17%	467,692	29.51%	0	0.00%	0	0.00%
Never	2,924,548	924,939	31.63%	543,501	18.58%	723,268	24.73%	681,115	23.29%	51,724	1.77%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	8,059,551	1,622,224	20.13%	1,272,655	15.79%	2,855,103	35.43%	2,239,103	27.78%	70,467	0.87%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Younger than 30	608,691	316,257	51.96%	108,031	17.75%	20,405	3.35%	163,999	26.94%	0	0.00%	0	0.00%
30 to 34	*	*	*	*	*	*	*	*	*	*	*	*	*
35 to 39	845,331	90,960	10.76%	218,664	25.87%	365,965	43.29%	169,741	20.08%	0	0.00%	0	0.00%
40 to 44	1,166,113	306,121	26.25%	132,736	11.38%	358,001	30.70%	369,256	31.67%	0	0.00%	0	0.00%
45 to 49	1,220,611	163,196	13.37%	239,512	19.62%	633,838	51.93%	132,341	10.84%	51,724	4.24%	0	0.00%
50 to 54	1,218,090	190,631	15.65%	259,036	21.27%	463,486	38.05%	304,937	25.03%	0	0.00%	0	0.00%
55 or older	409,726	102,556	25.03%	48,241	11.77%	148,140	36.16%	110,789	27.04%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Had period in last 12 months	2,553,814	388,783	15.22%	369,857	14.48%	951,851	37.27%	843,322	33.02%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
None	796,270	0	0.00%	93,123	11.69%	213,371	26.80%	489,776	61.51%	0	0.00%	0	0.00%
1	1,143,023	223,918	19.59%	259,298	22.69%	364,904	31.92%	294,904	25.80%	0	0.00%	0	0.00%
2	1,377,198	235,221	17.08%	179,370	13.02%	698,410	50.71%	245,454	17.82%	18,743	1.36%	0	0.00%
More than 2	5,274,713	1,399,364	26.53%	846,911	16.06%	1,767,746	33.51%	1,208,968	22.92%	51,724	0.98%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	1,830,851	141,095	7.71%	355,388	19.41%	589,242	32.18%	745,126	40.70%	0	0.00%	0	0.00%
1	1,101,117	108,111	9.82%	118,422	10.75%	504,889	45.85%	369,694	33.57%	0	0.00%	0	0.00%
2	1,912,599	375,478	19.63%	387,450	20.26%	793,781	41.50%	337,147	17.63%	18,743	0.98%	0	0.00%
More than 2	3,710,217	1,233,818	33.25%	511,899	13.80%	1,148,724	30.96%	764,052	20.59%	51,724	1.39%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	3,211,569	635,357	19.78%	692,177	21.55%	1,203,015	37.46%	681,020	21.21%	0	0.00%	0	0.00%
1	635,809	176,831	27.81%	146,525	23.05%	111,268	17.50%	201,185	31.64%	0	0.00%	0	0.00%
2	*	*	*	*	*	*	*	*	*	*	*	*	*
More than 2	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	4,284,887	1,021,026	23.83%	527,448	12.31%	1,357,140	31.67%	1,308,806	30.54%	70,467	1.64%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
29 years old or younger	5,817,235	1,500,758	25.80%	872,729	15.00%	2,266,267	38.96%	1,107,015	19.03%	70,467	1.21%	0	0.00%
30 - 40 years old	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	796,270	0	0.00%	93,123	11.69%	213,371	26.80%	489,776	61.51%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	946,721	229,752	24.27%	177,182	18.72%	301,499	31.85%	238,288	25.17%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
29 years old or younger	3,852,200	1,080,204	28.04%	483,459	12.55%	1,577,246	40.94%	692,549	17.98%	18,743	0.49%	0	0.00%
30 - 40 years old	3,133,784	633,961	20.23%	576,167	18.39%	1,119,669	35.73%	752,264	24.00%	51,724	1.65%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	796,270	0	0.00%	93,123	11.69%	213,371	26.80%	489,776	61.51%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,214,911	388,783	17.55%	305,675	13.80%	915,052	41.31%	605,401	27.33%	0	0.00%	0	0.00%
Yes	3,111,137	1,060,652	34.09%	468,163	15.05%	880,784	28.31%	701,538	22.55%	0	0.00%	0	0.00%
No	3,265,157	409,068	12.53%	604,865	18.52%	1,248,594	38.24%	932,163	28.55%	70,467	2.16%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,214,911	388,783	17.55%	305,675	13.80%	915,052	41.31%	605,401	27.33%	0	0.00%	0	0.00%
35 years old or younger	1,256,247	584,251	46.51%	129,654	10.32%	166,504	13.25%	375,837	29.92%	0	0.00%	0	0.00%
36 - 50 years old	1,506,680	392,625	26.06%	227,105	15.07%	597,249	39.64%	289,701	19.23%	0	0.00%	0	0.00%
51 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	3,265,157	409,068	12.53%	604,865	18.52%	1,248,594	38.24%	932,163	28.55%	70,467	2.16%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	8,591,205	1,858,503	21.63%	1,378,703	16.05%	3,044,430	35.44%	2,239,103	26.06%	70,467	0.82%	0	0.00%
***Had hysterectomy	-	-		-		-		-		-		-	
Yes	3,111,137	1,060,652	34.09%	468,163	15.05%	880,784	28.31%	701,538	22.55%	0	0.00%	0	0.00%
No	5,480,068	797,851	14.56%	910,540	16.62%	2,163,646	39.48%	1,537,565	28.06%	70,467	1.29%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	668,375	7,761	1.16%	66,968	10.02%	141,211	21.13%	429,920	64.32%	22,516	3.37%	0	0.00%
30-39	785,340	199,787	25.44%	168,234	21.42%	179,506	22.86%	237,812	30.28%	0	0.00%	0	0.00%
40-49	1,473,057	191,176	12.98%	229,422	15.57%	538,963	36.59%	513,496	34.86%	0	0.00%	0	0.00%
50-59	1,831,130	285,146	15.57%	433,411	23.67%	630,810	34.45%	481,763	26.31%	0	0.00%	0	0.00%
60-64	531,525	191,488	36.03%	41,282	7.77%	210,249	39.56%	88,506	16.65%	0	0.00%	0	0.00%
65-69	1,142,855	232,103	20.31%	159,134	13.92%	273,864	23.96%	477,754	41.80%	0	0.00%	0	0.00%
70-74	796,825	95,361	11.97%	113,105	14.19%	325,654	40.87%	191,475	24.03%	71,231	8.94%	0	0.00%
75-79	798,084	102,002	12.78%	247,614	31.03%	305,589	38.29%	142,879	17.90%	0	0.00%	0	0.00%
80+	1,090,291	394,389	36.17%	201,997	18.53%	256,607	23.54%	237,299	21.76%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	507,189	47,119	9.29%	103,825	20.47%	278,570	54.92%	77,675	15.31%	0	0.00%	0	0.00%
Other Hispanic	400,231	82,130	20.52%	70,641	17.65%	117,936	29.47%	107,009	26.74%	22,516	5.63%	0	0.00%
Non-Hispanic White	7,117,719	1,422,142	19.98%	1,170,448	16.44%	2,252,447	31.65%	2,201,451	30.93%	71,231	1.00%	0	0.00%
Non-Hispanic Black	905,114	125,465	13.86%	266,529	29.45%	213,500	23.59%	299,621	33.10%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	1,060,094	322,406	30.41%	245,115	23.12%	263,141	24.82%	229,433	21.64%	0	0.00%	0	0.00%
9-11th Grade	1,091,691	224,712	20.58%	190,808	17.48%	334,889	30.68%	318,766	29.20%	22,516	2.06%	0	0.00%
High School Grad/GED or Equivalent	6,703,785	1,106,990	16.51%	1,139,947	17.00%	2,214,573	33.03%	2,171,045	32.39%	71,231	1.06%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	5,012,841	777,400	15.51%	962,023	19.19%	1,949,627	38.89%	1,252,559	24.99%	71,231	1.42%	0	0.00%
Widowed	1,467,915	405,325	27.61%	252,456	17.20%	365,138	24.87%	444,996	30.31%	0	0.00%	0	0.00%
Divorced	1,160,194	346,590	29.87%	94,362	8.13%	214,581	18.50%	504,661	43.50%	0	0.00%	0	0.00%
Separated	261,394	73,304	28.04%	78,090	29.87%	85,878	32.85%	24,122	9.23%	0	0.00%	0	0.00%
Never Married	872,982	96,593	11.06%	198,445	22.73%	230,349	26.39%	325,079	37.24%	22,516	2.58%	0	0.00%
Living with Partner	337,055	0	0.00%	70,690	20.97%	16,879	5.01%	249,486	74.02%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	691,218	115,022	16.64%	108,927	15.76%	314,650	45.52%	152,619	22.08%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,309,416	157,190	12.00%	316,379	24.16%	320,075	24.44%	515,771	39.39%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	7,116,849	1,427,000	20.05%	1,235,862	17.37%	2,227,727	31.30%	2,132,513	29.96%	93,747	1.32%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,988,235	419,494	21.10%	350,450	17.63%	412,806	20.76%	782,969	39.38%	22,516	1.13%	0	0.00%
Overweight: 25.0 - 29.9	2,599,311	413,268	15.90%	404,926	15.58%	905,675	34.84%	875,442	33.68%	0	0.00%	0	0.00%
Obese: 30 or higher	4,385,952	772,439	17.61%	905,792	20.65%	1,521,135	34.68%	1,115,356	25.43%	71,231	1.62%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	496,522	13,028	2.62%	109,693	22.09%	169,710	34.18%	204,091	41.10%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	4,357,411	654,103	15.01%	695,579	15.96%	1,558,706	35.77%	1,426,507	32.74%	22,516	0.52%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	2,927,731	881,258	30.10%	558,960	19.09%	653,267	22.31%	834,245	28.49%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,335,819	150,823	11.29%	296,936	22.23%	480,770	35.99%	336,060	25.16%	71,231	5.33%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,751,046	280,660	16.03%	567,524	32.41%	389,077	22.22%	442,554	25.27%	71,231	4.07%	0	0.00%
No	7,366,436	1,418,552	19.26%	1,093,644	14.85%	2,473,375	33.58%	2,358,349	32.01%	22,516	0.31%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,955,072	840,150	21.24%	798,959	20.20%	1,094,391	27.67%	1,127,825	28.52%	93,747	2.37%	0	0.00%
No	5,162,411	859,062	16.64%	862,209	16.70%	1,768,062	34.25%	1,673,078	32.41%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,939,492	846,712	17.14%	1,065,749	21.58%	1,457,771	29.51%	1,498,028	30.33%	71,231	1.44%	0	0.00%
No	4,177,990	852,500	20.40%	595,419	14.25%	1,404,681	33.62%	1,302,875	31.18%	22,516	0.54%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,628,578	458,392	28.15%	252,595	15.51%	299,901	18.41%	617,689	37.93%	0	0.00%	0	0.00%
Mobility limitations	2,081,480	404,091	19.41%	348,538	16.74%	824,742	39.62%	432,878	20.80%	71,231	3.42%	0	0.00%
No mobility limitations	2,426,943	419,624	17.29%	522,767	21.54%	770,213	31.74%	714,339	29.43%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	2,980,481	417,105	13.99%	537,268	18.03%	967,595	32.46%	1,035,997	34.76%	22,516	0.76%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,662,416	331,071	19.92%	438,629	26.39%	568,516	34.20%	324,199	19.50%	0	0.00%	0	0.00%
No	7,455,067	1,368,141	18.35%	1,222,539	16.40%	2,293,937	30.77%	2,476,704	33.22%	93,747	1.26%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	1,117,761	350,101	31.32%	186,425	16.68%	297,101	26.58%	284,135	25.42%	0	0.00%	0	0.00%
Once a day	677,767	56,601	8.35%	61,212	9.03%	415,397	61.29%	144,558	21.33%	0	0.00%	0	0.00%
2 or more time a week	949,718	154,896	16.31%	161,040	16.96%	395,431	41.64%	238,350	25.10%	0	0.00%	0	0.00%
Once a week	978,226	58,039	5.93%	150,459	15.38%	180,511	18.45%	589,217	60.23%	0	0.00%	0	0.00%
1-3 times a month	1,580,930	331,258	20.95%	370,873	23.46%	335,925	21.25%	471,643	29.83%	71,231	4.51%	0	0.00%
Never	3,727,438	735,289	19.73%	731,159	19.62%	1,197,859	32.14%	1,040,615	27.92%	22,516	0.60%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	601,033	202,251	33.65%	58,125	9.67%	48,309	8.04%	221,116	36.79%	71,231	11.85%	0	0.00%
Never	8,258,897	1,361,928	16.49%	1,531,846	18.55%	2,762,820	33.45%	2,579,787	31.24%	22,516	0.27%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Younger than 30	423,228	72,635	17.16%	141,261	33.38%	95,062	22.46%	43,040	10.17%	71,231	16.83%	0	0.00%
30 to 34	383,865	21,727	5.66%	12,093	3.15%	36,144	9.42%	313,901	81.77%	0	0.00%	0	0.00%
35 to 39	525,354	92,593	17.62%	35,257	6.71%	201,336	38.32%	196,168	37.34%	0	0.00%	0	0.00%
40 to 44	920,906	273,434	29.69%	213,446	23.18%	239,827	26.04%	194,199	21.09%	0	0.00%	0	0.00%
45 to 49	1,593,743	275,329	17.28%	338,044	21.21%	533,212	33.46%	447,157	28.06%	0	0.00%	0	0.00%
50 to 54	1,513,834	549,579	36.30%	171,338	11.32%	524,343	34.64%	268,575	17.74%	0	0.00%	0	0.00%
55 or older	763,634	106,692	13.97%	254,176	33.29%	208,976	27.37%	193,790	25.38%	0	0.00%	0	0.00%
N/A - Had period in last 12 months	2,992,920	307,224	10.27%	495,554	16.56%	1,023,554	34.20%	1,144,073	38.23%	22,516	0.75%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
None	1,207,501	282,409	23.39%	205,792	17.04%	293,002	24.27%	426,298	35.30%	0	0.00%	0	0.00%
1	1,087,943	297,045	27.30%	211,972	19.48%	291,977	26.84%	286,948	26.38%	0	0.00%	0	0.00%
2	1,423,890	166,118	11.67%	394,832	27.73%	521,092	36.60%	341,848	24.01%	0	0.00%	0	0.00%
More than 2	5,398,149	953,640	17.67%	848,572	15.72%	1,756,381	32.54%	1,745,809	32.34%	93,747	1.74%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	2,265,188	368,437	16.27%	335,731	14.82%	753,323	33.26%	736,466	32.51%	71,231	3.14%	0	0.00%
1	1,170,086	244,515	20.90%	311,045	26.58%	436,783	37.33%	177,743	15.19%	0	0.00%	0	0.00%
2	1,685,041	232,386	13.79%	450,601	26.74%	565,202	33.54%	436,852	25.93%	0	0.00%	0	0.00%
More than 2	3,949,583	853,874	21.62%	536,889	13.59%	1,107,144	28.03%	1,429,159	36.19%	22,516	0.57%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	2,594,369	371,968	14.34%	447,753	17.26%	902,090	34.77%	850,042	32.76%	22,516	0.87%	0	0.00%
1	840,156	229,877	27.36%	195,874	23.31%	287,270	34.19%	127,134	15.13%	0	0.00%	0	0.00%
2	*	*	*	*	*	*	*	*	*	*	*	*	*
More than 2	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	5,089,680	1,072,706	21.08%	863,557	16.97%	1,416,258	27.83%	1,737,159	34.13%	0	0.00%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,628,516	304,135	18.68%	232,693	14.29%	527,557	32.39%	564,130	34.64%	0	0.00%	0	0.00%
Yes	1,383,386	183,340	13.25%	99,803	7.21%	638,455	46.15%	461,789	33.38%	0	0.00%	0	0.00%
No	6,088,701	1,211,737	19.90%	1,328,672	21.82%	1,679,561	27.58%	1,774,985	29.15%	93,747	1.54%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,645,395	304,135	18.48%	232,693	14.14%	544,437	33.09%	564,130	34.29%	0	0.00%	0	0.00%
35 years old or younger	1,359,049	183,340	13.49%	99,803	7.34%	638,455	46.98%	437,452	32.19%	0	0.00%	0	0.00%
36 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	6,088,701	1,211,737	19.90%	1,328,672	21.82%	1,679,561	27.58%	1,774,985	29.15%	93,747	1.54%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
29 years old or younger	5,727,469	1,048,107	18.30%	913,162	15.94%	1,777,685	31.04%	1,894,769	33.08%	93,747	1.64%	0	0.00%
30 - 40 years old	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,207,501	282,409	23.39%	205,792	17.04%	293,002	24.27%	426,298	35.30%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	1,394,439	308,816	22.15%	402,042	28.83%	405,716	29.10%	277,865	19.93%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	497,540	98,251	19.75%	26,902	5.41%	234,555	47.14%	137,831	27.70%	0	0.00%	0	0.00%
29 years old or younger	3,900,864	632,383	16.21%	702,720	18.01%	1,014,992	26.02%	1,457,022	37.35%	93,747	2.40%	0	0.00%
30 - 40 years old	3,248,452	665,071	20.47%	654,241	20.14%	1,279,882	39.40%	649,258	19.99%	0	0.00%	0	0.00%
41 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	1,207,501	282,409	23.39%	205,792	17.04%	293,002	24.27%	426,298	35.30%	0	0.00%	0	0.00%
No pregnancies resulting in live births	*	*	*	*	*	*	*	*	*	*	*	*	*
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,164,286	635,572	20.09%	715,659	22.62%	892,347	28.20%	849,477	26.85%	71,231	2.25%	0	0.00%
No	5,953,196	1,063,640	17.87%	945,509	15.88%	1,970,106	33.09%	1,951,426	32.78%	22,516	0.38%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
35 years old or younger	537,281	40,868	7.61%	153,354	28.54%	83,909	15.62%	187,919	34.98%	71,231	13.26%	0	0.00%
36 - 50 years old	2,064,040	452,606	21.93%	339,937	16.47%	633,367	30.69%	638,130	30.92%	0	0.00%	0	0.00%
51 years or older	539,538	142,098	26.34%	222,368	41.21%	175,072	32.45%	0	0.00%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	5,953,196	1,063,640	17.87%	945,509	15.88%	1,970,106	33.09%	1,951,426	32.78%	22,516	0.38%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	2,094,025	514,613	24.58%	377,748	18.04%	504,385	24.09%	626,048	29.90%	71,231	3.40%	0	0.00%
No	6,913,187	1,184,599	17.14%	1,256,518	18.18%	2,295,382	33.20%	2,154,172	31.16%	22,516	0.33%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
35 years old or younger	*	*	*	*	*	*	*	*	*	*	*	*	*
36 - 50 years old	1,112,276	297,733	26.77%	132,703	11.93%	313,036	28.14%	368,804	33.16%	0	0.00%	0	0.00%
51 years or older	678,554	194,524	28.67%	210,032	30.95%	131,040	19.31%	142,957	21.07%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	6,913,187	1,184,599	17.14%	1,256,518	18.18%	2,295,382	33.20%	2,154,172	31.16%	22,516	0.33%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,117,482	1,699,212	18.64%	1,661,168	18.22%	2,862,453	31.40%	2,800,903	30.72%	93,747	1.03%	0	0.00%
***Had hysterectomy	-	-		-		-		-		-		-	
Yes	3,134,924	635,572	20.27%	715,659	22.83%	862,984	27.53%	849,477	27.10%	71,231	2.27%	0	0.00%
No	5,982,559	1,063,640	17.78%	945,509	15.80%	1,999,468	33.42%	1,951,426	32.62%	22,516	0.38%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	519,643	132,636	25.52%	45,597	8.77%	26,984	5.19%	314,426	60.51%	0	0.00%	0	0.00%
30-39	752,154	0	0.00%	43,063	5.73%	412,626	54.86%	296,464	39.42%	0	0.00%	0	0.00%
40-49	1,842,589	216,471	11.75%	344,809	18.71%	703,126	38.16%	578,183	31.38%	0	0.00%	0	0.00%
50-59	1,536,416	176,441	11.48%	433,870	28.24%	476,503	31.01%	449,602	29.26%	0	0.00%	0	0.00%
60-64	928,762	169,049	18.20%	111,880	12.05%	268,244	28.88%	379,589	40.87%	0	0.00%	0	0.00%
65-69	1,052,909	332,311	31.56%	120,286	11.42%	361,865	34.37%	238,447	22.65%	0	0.00%	0	0.00%
70-74	749,231	267,248	35.67%	163,303	21.80%	233,535	31.17%	85,146	11.36%	0	0.00%	0	0.00%
75-79	743,102	72,435	9.75%	154,099	20.74%	327,354	44.05%	189,214	25.46%	0	0.00%	0	0.00%
80+	1,068,038	448,470	41.99%	211,139	19.77%	253,279	23.71%	117,676	11.02%	37,475	3.51%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	564,696	121,368	21.49%	151,342	26.80%	187,740	33.25%	104,247	18.46%	0	0.00%	0	0.00%
Other Hispanic	181,284	42,268	23.32%	28,534	15.74%	50,576	27.90%	59,905	33.04%	0	0.00%	0	0.00%
Non-Hispanic White	7,326,980	1,616,807	22.07%	1,231,677	16.81%	2,450,205	33.44%	1,990,816	27.17%	37,475	0.51%	0	0.00%
Non-Hispanic Black	865,608	34,617	4.00%	193,540	22.36%	241,676	27.92%	395,775	45.72%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	598,173	92,406	15.45%	117,900	19.71%	307,840	51.46%	80,026	13.38%	0	0.00%	0	0.00%
9-11th Grade	966,081	196,028	20.29%	217,345	22.50%	341,316	35.33%	211,393	21.88%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	7,274,457	1,444,670	19.86%	1,101,000	15.14%	2,340,427	32.17%	2,350,885	32.32%	37,475	0.52%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Married	5,484,160	1,081,096	19.71%	949,171	17.31%	1,847,419	33.69%	1,606,474	29.29%	0	0.00%	0	0.00%
Widowed	1,492,588	344,318	23.07%	311,838	20.89%	551,014	36.92%	247,944	16.61%	37,475	2.51%	0	0.00%
Divorced	910,403	199,761	21.94%	80,670	8.86%	358,245	39.35%	271,727	29.85%	0	0.00%	0	0.00%
Separated	225,238	45,514	20.21%	26,422	11.73%	50,013	22.20%	103,289	45.86%	0	0.00%	0	0.00%
Never Married	804,039	144,371	17.96%	223,157	27.75%	172,577	21.46%	263,934	32.83%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	597,493	156,204	26.14%	125,679	21.03%	249,261	41.72%	66,349	11.10%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,339,274	271,612	20.28%	316,301	23.62%	472,641	35.29%	241,246	18.01%	37,475	2.80%	0	0.00%
Poverty income ratio 1 or greater	7,256,077	1,387,244	19.12%	1,186,066	16.35%	2,341,613	32.27%	2,341,154	32.26%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	2,314,853	189,942	8.21%	158,737	6.86%	909,211	39.28%	1,056,963	45.66%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	2,461,339	554,271	22.52%	489,888	19.90%	854,000	34.70%	525,705	21.36%	37,475	1.52%	0	0.00%
Obese: 30 or higher	4,346,697	1,025,056	23.58%	979,421	22.53%	1,276,140	29.36%	1,066,080	24.53%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	325,483	45,236	13.90%	48,864	15.01%	122,272	37.57%	109,111	33.52%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	4,793,325	930,836	19.42%	934,356	19.49%	1,599,743	33.37%	1,328,391	27.71%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	3,294,171	724,834	22.00%	452,147	13.73%	1,002,752	30.44%	1,076,962	32.69%	37,475	1.14%	0	0.00%
Glycohemoglobin 6.5% or higher	779,864	114,154	14.64%	192,680	24.71%	338,748	43.44%	134,283	17.22%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,378,672	316,935	22.99%	274,323	19.90%	507,180	36.79%	280,235	20.33%	0	0.00%	0	0.00%
No	7,814,171	1,498,125	19.17%	1,353,723	17.32%	2,556,335	32.71%	2,368,513	30.31%	37,475	0.48%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,035,294	740,140	18.34%	756,274	18.74%	1,509,328	37.40%	1,029,553	25.51%	0	0.00%	0	0.00%
No	5,157,549	1,074,920	20.84%	871,772	16.90%	1,554,187	30.13%	1,619,195	31.39%	37,475	0.73%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	4,334,527	1,146,693	26.45%	964,065	22.24%	1,567,447	36.16%	656,321	15.14%	0	0.00%	0	0.00%
No	4,858,317	668,366	13.76%	663,981	13.67%	1,496,068	30.79%	1,992,427	41.01%	37,475	0.77%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,817,520	586,780	32.28%	345,759	19.02%	462,721	25.46%	422,261	23.23%	0	0.00%	0	0.00%
Mobility limitations	1,773,451	540,514	30.48%	364,495	20.55%	738,860	41.66%	129,582	7.31%	0	0.00%	0	0.00%
No mobility limitations	2,479,846	436,493	17.60%	445,565	17.97%	880,334	35.50%	679,979	27.42%	37,475	1.51%	0	0.00%
N/A - Respondent not asked mobility limitation questions	3,122,026	251,273	8.05%	472,226	15.13%	981,600	31.44%	1,416,926	45.38%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,383,609	282,828	20.44%	227,826	16.47%	579,448	41.88%	293,507	21.21%	0	0.00%	0	0.00%
No	7,740,209	1,532,231	19.80%	1,331,194	17.20%	2,484,067	32.09%	2,355,241	30.43%	37,475	0.48%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	1,517,780	488,166	32.16%	294,884	19.43%	456,691	30.09%	278,040	18.32%	0	0.00%	0	0.00%
Once a day	754,464	118,784	15.74%	192,084	25.46%	186,112	24.67%	257,484	34.13%	0	0.00%	0	0.00%
2 or more time a week	1,035,574	261,066	25.21%	134,441	12.98%	379,244	36.62%	260,823	25.19%	0	0.00%	0	0.00%
Once a week	349,564	26,491	7.58%	72,477	20.73%	155,916	44.60%	94,680	27.09%	0	0.00%	0	0.00%
1-3 times a month	1,599,420	363,200	22.71%	379,333	23.72%	456,464	28.54%	362,949	22.69%	37,475	2.34%	0	0.00%
Never	3,802,206	469,838	12.36%	554,828	14.59%	1,429,088	37.59%	1,348,453	35.47%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	8,656,354	1,688,341	19.50%	1,533,342	17.71%	2,837,068	32.77%	2,560,127	29.58%	37,475	0.43%	0	0.00%
Age of patient at last menstruation period	-	-	-	-	-	-	-	-	-	-	-	-	-
Younger than 30	617,157	120,590	19.54%	127,898	20.72%	279,614	45.31%	89,055	14.43%	0	0.00%	0	0.00%
30 to 34	527,249	134,577	25.52%	67,834	12.87%	239,423	45.41%	85,416	16.20%	0	0.00%	0	0.00%
35 to 39	538,743	157,594	29.25%	112,263	20.84%	268,886	49.91%	0	0.00%	0	0.00%	0	0.00%
40 to 44	1,088,292	298,463	27.42%	272,354	25.03%	203,559	18.70%	313,916	28.84%	0	0.00%	0	0.00%
45 to 49	1,330,479	257,429	19.35%	225,544	16.95%	373,745	28.09%	436,285	32.79%	37,475	2.82%	0	0.00%
50 to 54	1,339,020	307,001	22.93%	270,416	20.20%	510,813	38.15%	250,791	18.73%	0	0.00%	0	0.00%
55 or older	886,363	336,071	37.92%	77,512	8.74%	165,842	18.71%	306,938	34.63%	0	0.00%	0	0.00%
N/A - Had period in last 12 months	2,865,540	203,335	7.10%	474,225	16.55%	1,021,633	35.65%	1,166,347	40.70%	0	0.00%	0	0.00%
Total number of pregnancies	-	-	-	-	-	-	-	-	-	-	-	-	-
None	789,721	176,999	22.41%	58,911	7.46%	193,208	24.47%	360,602	45.66%	0	0.00%	0	0.00%
1	889,117	127,731	14.37%	101,574	11.42%	368,966	41.50%	290,846	32.71%	0	0.00%	0	0.00%
2	2,162,913	364,027	16.83%	260,660	12.05%	964,793	44.61%	573,433	26.51%	0	0.00%	0	0.00%
More than 2	5,351,092	1,146,302	21.42%	1,206,901	22.55%	1,536,547	28.71%	1,423,866	26.61%	37,475	0.70%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Number of vaginal deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	2,223,632	187,371	8.43%	232,158	10.44%	728,035	32.74%	1,076,068	48.39%	0	0.00%	0	0.00%
1	1,038,297	278,268	26.80%	144,150	13.88%	284,260	27.38%	331,619	31.94%	0	0.00%	0	0.00%
2	2,330,719	366,778	15.74%	472,479	20.27%	1,097,901	47.11%	393,562	16.89%	0	0.00%	0	0.00%
More than 2	3,570,874	953,322	26.70%	779,259	21.82%	953,319	26.70%	847,499	23.73%	37,475	1.05%	0	0.00%
Number of cesarean deliveries	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
None	2,556,782	226,794	8.87%	509,627	19.93%	1,287,950	50.37%	532,412	20.82%	0	0.00%	0	0.00%
1	778,210	177,582	22.82%	127,067	16.33%	157,146	20.19%	278,940	35.84%	37,475	4.82%	0	0.00%
2	522,394	85,108	16.29%	35,168	6.73%	62,938	12.05%	339,179	64.93%	0	0.00%	0	0.00%
More than 2	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Vaginal delivery/deliveries only	4,941,102	1,259,519	25.49%	807,102	16.33%	1,548,688	31.34%	1,325,794	26.83%	0	0.00%	0	0.00%
Gave birth to any babies 9lb or more	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,293,065	206,321	15.96%	58,911	4.56%	596,525	46.13%	431,307	33.36%	0	0.00%	0	0.00%
Yes	1,482,274	292,312	19.72%	299,874	20.23%	397,902	26.84%	454,710	30.68%	37,475	2.53%	0	0.00%
No	6,371,929	1,286,280	20.19%	1,269,260	19.92%	2,069,087	32.47%	1,747,301	27.42%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
Age of patient when she had 9 lb or more baby	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,338,640	236,467	17.66%	58,911	4.40%	596,525	44.56%	446,736	33.37%	0	0.00%	0	0.00%
35 years old or younger	1,241,612	292,312	23.54%	240,327	19.36%	265,623	21.39%	405,875	32.69%	37,475	3.02%	0	0.00%
36 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have baby weighing 9 lbs or more	6,371,929	1,286,280	20.19%	1,269,260	19.92%	2,069,087	32.47%	1,747,301	27.42%	0	0.00%	0	0.00%
Age of patient at 1st live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	503,344	29,321	5.83%	0	0.00%	403,317	80.13%	70,705	14.05%	0	0.00%	0	0.00%
29 years old or younger	6,119,339	1,248,379	20.40%	1,346,904	22.01%	1,973,313	32.25%	1,513,268	24.73%	37,475	0.61%	0	0.00%
30 - 40 years old	778,628	223,818	28.75%	69,026	8.87%	147,637	18.96%	338,147	43.43%	0	0.00%	0	0.00%
N/A - Never pregnant	789,721	176,999	22.41%	58,911	7.46%	193,208	24.47%	360,602	45.66%	0	0.00%	0	0.00%
Only 1 live birth - see age at last live birth	1,001,812	136,542	13.63%	153,205	15.29%	346,039	34.54%	366,026	36.54%	0	0.00%	0	0.00%
Age of patient at last live birth	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	586,694	29,321	5.00%	0	0.00%	425,816	72.58%	131,557	22.42%	0	0.00%	0	0.00%
29 years old or younger	4,052,100	764,836	18.88%	768,713	18.97%	1,620,162	39.98%	898,388	22.17%	0	0.00%	0	0.00%
30 - 40 years old	3,470,428	748,632	21.57%	746,249	21.50%	717,347	20.67%	1,258,201	36.25%	0	0.00%	0	0.00%
41 years or older	211,824	95,271	44.98%	35,891	16.94%	80,662	38.08%	0	0.00%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Never pregnant	789,721	176,999	22.41%	58,911	7.46%	193,208	24.47%	360,602	45.66%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
**Had hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,258,347	908,953	27.90%	808,360	24.81%	1,000,081	30.69%	503,478	15.45%	37,475	1.15%	0	0.00%
No	5,934,497	906,107	15.27%	819,686	13.81%	2,063,434	34.77%	2,145,270	36.15%	0	0.00%	0	0.00%
Age at time of hysterectomy	-	-	-	-	-	-	-	-	-	-	-	-	-
35 years old or younger	1,006,766	218,031	21.66%	135,423	13.45%	478,840	47.56%	174,471	17.33%	0	0.00%	0	0.00%
36 - 50 years old	1,716,547	621,664	36.22%	538,026	31.34%	361,976	21.09%	194,881	11.35%	0	0.00%	0	0.00%
51 years or older	399,246	69,258	17.35%	105,185	26.35%	106,107	26.58%	118,697	29.73%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	5,934,497	906,107	15.27%	819,686	13.81%	2,063,434	34.77%	2,145,270	36.15%	0	0.00%	0	0.00%
Had ovaries removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	1,476,026	410,076	27.78%	348,425	23.61%	482,558	32.69%	234,966	15.92%	0	0.00%	0	0.00%
No	7,445,076	1,369,841	18.40%	1,255,457	16.86%	2,475,672	33.25%	2,306,631	30.98%	37,475	0.50%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Age when ovaries were removed	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
35 years old or younger	387,825	125,371	32.33%	115,218	29.71%	96,733	24.94%	50,503	13.02%	0	0.00%	0	0.00%
36 - 50 years old	805,858	221,019	27.43%	149,759	18.58%	310,998	38.59%	124,082	15.40%	0	0.00%	0	0.00%
51 years or older	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Did not have procedure	7,445,076	1,369,841	18.40%	1,255,457	16.86%	2,475,672	33.25%	2,306,631	30.98%	37,475	0.50%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.

Table F05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	9,192,844	1,815,060	19.74%	1,628,046	17.71%	3,063,515	33.32%	2,648,748	28.81%	37,475	0.41%	0	0.00%
***Had hysterectomy	-	-		-		-		-		-		-	
Yes	3,242,970	908,953	28.03%	808,360	24.93%	984,704	30.36%	503,478	15.53%	37,475	1.16%	0	0.00%
No	5,949,873	906,107	15.23%	819,686	13.78%	2,078,811	34.94%	2,145,270	36.06%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

NOTE: **The question asking whether the respondent had a hysterectomy was only asked of women who reported not having at least 1 or a regular period prior to survey administration.

NOTE: *** Those who reported having at least 1 or regular periods in the last 12 months were categorized as not having had a hysterectomy in this table. This table represents an alternative definition for "no" or missing hysterectomy data responses.