

Table M01 -Prevalence of any urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	90,330,740	10,266,409	11.37%	9.70%	13.03%	80,064,331	88.63%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	318,425	7,364	2.31%	( 2.96%)	7.59%	311,061	97.69%
20-29	11,767,942	373,700	3.18%	0.91%	5.44%	11,394,242	96.82%
30-39	18,212,868	1,245,096	6.84%	3.44%	10.24%	16,967,772	93.16%
40-49	22,567,759	1,282,331	5.68%	3.69%	7.67%	21,285,428	94.32%
50-59	18,568,743	2,024,486	10.90%	7.27%	14.54%	16,544,257	89.10%
60-64	4,648,416	929,881	20.00%	11.13%	28.88%	3,718,535	80.00%
65-69	3,884,538	1,080,369	27.81%	19.78%	35.84%	2,804,169	72.19%
70-74	4,039,191	1,373,568	34.01%	20.24%	47.77%	2,665,623	65.99%
75-79	2,934,099	758,372	25.85%	18.09%	33.60%	2,175,726	74.15%
80-84	2,102,740	624,458	29.70%	22.34%	37.06%	1,478,281	70.30%
85+	1,286,018	566,783	44.07%	27.97%	60.18%	719,236	55.93%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,055,890	475,436	6.74%	3.66%	9.81%	6,580,454	93.26%
Other Hispanic	4,966,199	553,379	11.14%	3.30%	18.98%	4,412,820	88.86%
Non-Hispanic White	66,176,833	7,730,812	11.68%	9.29%	14.07%	58,446,021	88.32%
Non-Hispanic Black	9,016,771	1,311,346	14.54%	8.16%	20.93%	7,705,425	85.46%
Other Race - Including Multi-Racial	3,115,046	195,436	6.27%	( 0.20%)	12.75%	2,919,610	93.73%
<b>Education</b>	-	-	-	-	-	-	-
Missing	933,223	86,728	9.29%	( 3.80%)	22.39%	846,494	90.71%
Less Than 9th Grade	6,516,375	1,274,348	19.56%	13.11%	26.01%	5,242,027	80.44%
9-11th Grade	10,170,827	1,393,333	13.70%	7.23%	20.17%	8,777,494	86.30%
High School Grad/GED or Equivalent	72,563,484	7,469,140	10.29%	8.67%	11.92%	65,094,345	89.71%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	909,023	190,990	21.01%	0.45%	41.57%	718,033	78.99%
Married	59,561,278	6,617,196	11.11%	9.08%	13.14%	52,944,082	88.89%
Widowed	4,052,855	861,620	21.26%	13.41%	29.11%	3,191,235	78.74%
Divorced	7,804,451	1,262,656	16.18%	10.33%	22.02%	6,541,795	83.82%
Separated	2,163,795	112,297	5.19%	0.06%	10.32%	2,051,498	94.81%
Never Married	10,212,194	983,139	9.63%	4.70%	14.55%	9,229,055	90.37%
Living with Partner	5,531,980	159,429	2.88%	( 0.17%)	5.94%	5,372,551	97.12%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	4,980,213	593,049	11.91%	6.26%	17.56%	4,387,165	88.09%
Poverty income ratio less than 1	9,683,534	1,516,688	15.66%	9.61%	21.72%	8,166,847	84.34%
Poverty income ratio 1 or greater	75,666,992	8,156,673	10.78%	8.88%	12.68%	67,510,319	89.22%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	3,185,619	874,992	27.47%	16.29%	38.65%	2,310,627	72.53%
Normal: less than 25	27,340,807	2,739,084	10.02%	6.46%	13.58%	24,601,723	89.98%
Overweight: 25.0 - 29.9	36,239,254	3,551,109	9.80%	7.74%	11.86%	32,688,145	90.20%
Obese: 30 or higher	23,565,060	3,101,225	13.16%	10.60%	15.72%	20,463,836	86.84%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	90,330,740	10,266,409	11.37%	9.70%	13.03%	80,064,331	88.63%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,136,613	416,345	13.27%	5.99%	20.56%	2,720,268	86.73%
Glycohemoglobin < 5.7%	69,910,044	6,339,137	9.07%	7.67%	10.46%	63,570,907	90.93%
Glycohemoglobin 5.7% - 6.4%	11,173,113	1,757,161	15.73%	9.88%	21.57%	9,415,952	84.27%
Glycohemoglobin 6.5% or higher	6,110,970	1,753,766	28.70%	20.88%	36.51%	4,357,203	71.30%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,576,525	1,695,100	25.78%	17.49%	34.06%	4,881,424	74.22%
No	83,754,215	8,571,309	10.23%	8.89%	11.58%	75,182,906	89.77%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	51,198,375	7,293,348	14.25%	11.84%	16.65%	43,905,027	85.75%
No	39,045,204	2,973,061	7.61%	6.28%	8.95%	36,072,143	92.39%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	1,046,623	10,036	0.96%	( 1.18%)	3.09%	1,036,586	99.04%
Yes	21,461,970	3,876,071	18.06%	14.29%	21.83%	17,585,899	81.94%
No	67,725,407	6,380,301	9.42%	7.67%	11.17%	61,345,105	90.58%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	5,308,418	1,480,096	27.88%	17.76%	38.01%	3,828,322	72.12%
Mobility limitations	6,868,586	2,270,983	33.06%	24.91%	41.22%	4,597,603	66.94%
No mobility limitations	18,439,144	3,383,951	18.35%	14.15%	22.56%	15,055,193	81.65%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	59,699,976	3,116,764	5.22%	3.77%	6.68%	56,583,213	94.78%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	90,330,740	10,266,409	11.37%	9.70%	13.03%	80,064,331	88.63%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	90,330,740	10,266,409	11.37%	9.70%	13.03%	80,064,331	88.63%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	90,330,740	10,266,409	11.37%	9.70%	13.03%	80,064,331	88.63%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	91,018,484	13,702,607	15.05%	12.65%	17.46%	77,315,876	84.95%
<b>Age at screening</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	538,199	0	0.00%	0.00%	0.00%	538,199	100.00%
20-29	12,110,825	920,444	7.60%	3.15%	12.05%	11,190,381	92.40%
30-39	17,735,503	1,961,991	11.06%	6.51%	15.62%	15,773,512	88.94%
40-49	20,941,149	1,948,077	9.30%	4.91%	13.70%	18,993,072	90.70%
50-59	17,868,567	2,843,199	15.91%	10.37%	21.45%	15,025,368	84.09%
60-64	5,666,205	1,000,574	17.66%	10.28%	25.03%	4,665,631	82.34%
65-69	5,762,216	1,335,420	23.18%	16.19%	30.17%	4,426,796	76.82%
70-74	3,988,818	1,398,131	35.05%	27.45%	42.65%	2,590,687	64.95%
75-79	3,030,438	1,080,370	35.65%	23.67%	47.63%	1,950,068	64.35%
80-84	2,022,770	662,074	32.73%	20.40%	45.06%	1,360,696	67.27%
85+	1,353,795	552,328	40.80%	24.81%	56.79%	801,467	59.20%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,316,409	987,200	13.49%	9.66%	17.33%	6,329,208	86.51%
Other Hispanic	2,977,019	501,938	16.86%	3.99%	29.73%	2,475,080	83.14%
Non-Hispanic White	66,918,680	9,809,316	14.66%	12.16%	17.16%	57,109,364	85.34%
Non-Hispanic Black	9,127,400	2,024,497	22.18%	17.19%	27.17%	7,102,902	77.82%
Other Race - Including Multi-Racial	4,678,976	379,655	8.11%	0.31%	15.92%	4,299,321	91.89%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,796,573	170,614	9.50%	4.67%	14.32%	1,625,959	90.50%
Less Than 9th Grade	6,556,875	1,425,778	21.74%	15.12%	28.37%	5,131,096	78.26%
9-11th Grade	8,965,501	1,149,203	12.82%	7.56%	18.08%	7,816,298	87.18%
High School Grad/GED or Equivalent	73,351,938	10,919,458	14.89%	12.01%	17.76%	62,432,480	85.11%
Refused	*	*	*	*	*	*	*
Don't Know	335,074	37,555	11.21%	( 3.89%)	26.30%	297,519	88.79%
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	1,019,622	215,516	21.14%	( 0.36%)	42.63%	804,106	78.86%
Married	59,788,711	8,855,567	14.81%	12.23%	17.40%	50,933,143	85.19%
Widowed	3,574,691	869,715	24.33%	13.83%	34.83%	2,704,976	75.67%
Divorced	7,570,723	1,217,057	16.08%	10.62%	21.53%	6,353,666	83.92%
Separated	1,505,477	334,445	22.22%	9.29%	35.14%	1,171,032	77.78%
Never Married	11,206,160	1,587,594	14.17%	8.12%	20.21%	9,618,566	85.83%
Living with Partner	5,961,389	622,715	10.45%	3.27%	17.62%	5,338,674	89.55%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	4,481,624	572,832	12.78%	7.45%	18.11%	3,908,792	87.22%
Poverty income ratio less than 1	9,237,240	1,153,158	12.48%	5.80%	19.17%	8,084,081	87.52%
Poverty income ratio 1 or greater	77,299,620	11,976,617	15.49%	12.75%	18.24%	65,323,003	84.51%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	960,921	268,019	27.89%	7.40%	48.39%	692,902	72.11%
Normal: less than 25	27,588,667	3,529,345	12.79%	10.05%	15.54%	24,059,322	87.21%
Overweight: 25.0 - 29.9	34,644,037	4,301,128	12.42%	9.27%	15.56%	30,342,909	87.58%
Obese: 30 or higher	27,824,858	5,604,114	20.14%	16.14%	24.14%	22,220,743	79.86%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	91,018,484	13,702,607	15.05%	12.65%	17.46%	77,315,876	84.95%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	2,412,745	361,588	14.99%	7.62%	22.36%	2,051,156	85.01%
Glycohemoglobin < 5.7%	69,680,688	8,890,742	12.76%	10.31%	15.21%	60,789,946	87.24%
Glycohemoglobin 5.7% - 6.4%	12,825,067	2,728,162	21.27%	16.75%	25.80%	10,096,904	78.73%
Glycohemoglobin 6.5% or higher	6,099,984	1,722,115	28.23%	21.08%	35.38%	4,377,869	71.77%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,922,443	2,199,006	31.77%	25.74%	37.79%	4,723,437	68.23%
No	84,096,041	11,503,602	13.68%	11.15%	16.21%	72,592,439	86.32%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,623,866	8,509,424	16.17%	13.81%	18.53%	44,114,442	83.83%
No	38,356,405	5,181,352	13.51%	9.62%	17.40%	33,175,052	86.49%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	789,967	34,108	4.32%	( 3.97%)	12.61%	755,859	95.68%
Yes	27,630,586	6,058,883	21.93%	17.75%	26.11%	21,571,703	78.07%
No	62,492,258	7,560,067	12.10%	9.76%	14.44%	54,932,190	87.90%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	9,336,477	2,964,807	31.76%	26.42%	37.09%	6,371,670	68.24%
Mobility limitations	7,834,752	1,853,409	23.66%	18.57%	28.74%	5,981,343	76.34%
No mobility limitations	19,576,491	3,470,630	17.73%	14.86%	20.59%	16,105,860	82.27%
N/A - Respondent not asked mobility limitation questions	54,270,763	5,413,761	9.98%	6.91%	13.04%	48,857,002	90.02%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	91,018,484	13,702,607	15.05%	12.65%	17.46%	77,315,876	84.95%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	91,018,484	13,702,607	15.05%	12.65%	17.46%	77,315,876	84.95%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	91,018,484	13,702,607	15.05%	12.65%	17.46%	77,315,876	84.95%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	93,642,892	14,122,748	15.08%	12.19%	17.98%	79,520,144	84.92%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	384,918	22,647	5.88%	( 5.78%)	17.54%	362,271	94.12%
20-29	14,540,015	899,897	6.19%	2.46%	9.92%	13,640,118	93.81%
30-39	17,503,545	1,031,971	5.90%	2.63%	9.17%	16,471,574	94.10%
40-49	21,809,592	2,978,059	13.65%	7.96%	19.35%	18,831,533	86.35%
50-59	18,142,193	3,614,824	19.92%	13.27%	26.58%	14,527,370	80.08%
60-64	6,373,461	1,085,089	17.03%	11.21%	22.84%	5,288,373	82.97%
65-69	5,297,573	1,466,777	27.69%	22.78%	32.59%	3,830,795	72.31%
70-74	4,006,172	1,031,394	25.75%	17.96%	33.53%	2,974,778	74.25%
75-79	2,536,289	959,797	37.84%	26.90%	48.79%	1,576,492	62.16%
80-84	2,002,541	733,159	36.61%	15.68%	57.54%	1,269,382	63.39%
85+	1,046,592	299,135	28.58%	17.79%	39.37%	747,457	71.42%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,979,822	1,116,500	13.99%	10.48%	17.51%	6,863,322	86.01%
Other Hispanic	2,862,199	285,223	9.97%	3.09%	16.85%	2,576,975	90.03%
Non-Hispanic White	69,085,485	10,764,351	15.58%	12.25%	18.91%	58,321,134	84.42%
Non-Hispanic Black	9,553,122	1,520,452	15.92%	12.31%	19.52%	8,032,670	84.08%
Other Race - Including Multi-Racial	4,162,265	436,222	10.48%	1.51%	19.45%	3,726,043	89.52%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,125,917	21,315	1.89%	( 2.21%)	5.99%	1,104,602	98.11%
Less Than 9th Grade	6,333,269	1,110,708	17.54%	12.56%	22.51%	5,222,561	82.46%
9-11th Grade	10,260,776	1,514,077	14.76%	9.80%	19.71%	8,746,699	85.24%
High School Grad/GED or Equivalent	75,757,393	11,429,497	15.09%	11.93%	18.24%	64,327,896	84.91%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	1,042,842	104,477	10.02%	( 6.91%)	26.94%	938,365	89.98%
Married	60,157,438	9,694,781	16.12%	12.94%	19.29%	50,462,657	83.88%
Widowed	2,914,762	577,904	19.83%	8.10%	31.55%	2,336,858	80.17%
Divorced	9,188,198	1,797,233	19.56%	13.22%	25.90%	7,390,965	80.44%
Separated	2,712,975	472,734	17.42%	5.43%	29.42%	2,240,241	82.58%
Never Married	9,472,270	999,423	10.55%	5.39%	15.71%	8,472,847	89.45%
Living with Partner	8,115,581	437,370	5.39%	0.35%	10.43%	7,678,211	94.61%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	2,977,906	811,536	27.25%	15.00%	39.50%	2,166,370	72.75%
Poverty income ratio less than 1	9,079,370	1,447,959	15.95%	10.88%	21.01%	7,631,411	84.05%
Poverty income ratio 1 or greater	81,585,616	11,863,253	14.54%	11.33%	17.75%	69,722,363	85.46%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	1,009,735	126,132	12.49%	2.16%	22.82%	883,603	87.51%
Normal: less than 25	25,820,546	3,052,898	11.82%	8.15%	15.50%	22,767,647	88.18%
Overweight: 25.0 - 29.9	35,914,294	5,408,869	15.06%	11.42%	18.70%	30,505,426	84.94%
Obese: 30 or higher	30,898,317	5,534,849	17.91%	12.91%	22.92%	25,363,468	82.09%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	93,642,892	14,122,748	15.08%	12.19%	17.98%	79,520,144	84.92%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,564,155	300,823	8.44%	1.76%	15.13%	3,263,332	91.56%
Glycohemoglobin < 5.7%	70,254,781	9,038,263	12.86%	10.11%	15.62%	61,216,519	87.14%
Glycohemoglobin 5.7% - 6.4%	14,222,346	3,182,425	22.38%	18.32%	26.43%	11,039,921	77.62%
Glycohemoglobin 6.5% or higher	5,601,610	1,601,238	28.59%	19.70%	37.47%	4,000,372	71.41%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,979,396	1,702,435	24.39%	17.67%	31.12%	5,276,960	75.61%
No	86,663,496	12,420,313	14.33%	11.48%	17.19%	74,243,184	85.67%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,967,879	9,608,565	18.14%	14.32%	21.96%	43,359,313	81.86%
No	40,675,013	4,514,183	11.10%	7.67%	14.52%	36,160,830	88.90%
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	26,812,046	6,581,507	24.55%	19.51%	29.58%	20,230,539	75.45%
No	66,652,383	7,484,663	11.23%	8.40%	14.06%	59,167,720	88.77%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	6,866,793	1,617,259	23.55%	14.72%	32.38%	5,249,535	76.45%
Mobility limitations	5,930,657	2,018,270	34.03%	23.09%	44.97%	3,912,386	65.97%
No mobility limitations	20,291,603	4,434,930	21.86%	19.01%	24.71%	15,856,674	78.14%
N/A - Respondent not asked mobility limitation questions	60,553,839	6,052,290	9.99%	7.60%	12.39%	54,501,549	90.01%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,936,022	2,079,229	26.20%	18.41%	33.99%	5,856,793	73.80%
No	85,590,767	12,043,519	14.07%	11.25%	16.90%	73,547,248	85.93%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	372,256	64,954	17.45%	( 4.15%)	39.05%	307,302	82.55%
2 or more times a day	13,735,930	2,904,617	21.15%	15.12%	27.17%	10,831,314	78.85%
Once a day	8,509,610	1,026,938	12.07%	5.88%	18.26%	7,482,673	87.93%
2 or more time a week	5,956,076	1,451,505	24.37%	13.77%	34.97%	4,504,570	75.63%
Once a week	5,314,628	1,065,210	20.04%	8.40%	31.68%	4,249,417	79.96%
1-3 times a month	9,361,331	2,228,101	23.80%	15.85%	31.75%	7,133,231	76.20%
Never	50,100,244	5,381,424	10.74%	7.87%	13.61%	44,718,821	89.26%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	372,256	64,954	17.45%	( 4.15%)	39.05%	307,302	82.55%
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	749,457	598,247	79.82%	60.18%	99.47%	151,210	20.18%
Never	92,124,364	13,338,270	14.48%	11.54%	17.42%	78,786,094	85.52%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	94,932,386	14,875,469	15.67%	13.41%	17.93%	80,056,917	84.33%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	581,889	15,708	2.70%	( 3.35%)	8.75%	566,180	97.30%
20-29	14,324,402	604,177	4.22%	1.76%	6.68%	13,720,225	95.78%
30-39	17,096,292	1,125,718	6.58%	4.54%	8.63%	15,970,573	93.42%
40-49	21,351,091	2,794,224	13.09%	8.41%	17.77%	18,556,867	86.91%
50-59	18,864,707	3,385,303	17.95%	12.46%	23.43%	15,479,404	82.05%
60-64	7,469,087	1,707,386	22.86%	18.09%	27.63%	5,761,701	77.14%
65-69	5,293,210	1,518,446	28.69%	21.73%	35.65%	3,774,765	71.31%
70-74	3,767,362	1,283,484	34.07%	25.05%	43.09%	2,483,878	65.93%
75-79	2,921,173	1,207,845	41.35%	31.07%	51.63%	1,713,329	58.65%
80+	3,263,173	1,233,178	37.79%	29.11%	46.48%	2,029,995	62.21%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	9,009,819	1,208,868	13.42%	7.69%	19.14%	7,800,950	86.58%
Other Hispanic	4,540,869	624,985	13.76%	9.33%	18.20%	3,915,884	86.24%
Non-Hispanic White	65,896,437	10,135,298	15.38%	12.97%	17.79%	55,761,138	84.62%
Non-Hispanic Black	9,832,078	2,092,450	21.28%	17.10%	25.46%	7,739,628	78.72%
Other Race - Including Multi-Racial	5,653,183	813,867	14.40%	3.37%	25.42%	4,839,316	85.60%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,275,540	181,349	14.22%	( 0.54%)	28.98%	1,094,191	85.78%
Less Than 9th Grade	6,695,497	1,853,886	27.69%	16.91%	38.47%	4,841,611	72.31%
9-11th Grade	11,881,477	2,135,076	17.97%	12.30%	23.64%	9,746,401	82.03%
High School Grad/GED or Equivalent	74,943,444	10,623,592	14.18%	11.68%	16.67%	64,319,852	85.82%
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	1,282,066	49,279	3.84%	( 0.12%)	7.81%	1,232,787	96.16%
Married	60,655,911	9,743,242	16.06%	13.23%	18.90%	50,912,668	83.94%
Widowed	3,208,196	884,467	27.57%	18.31%	36.83%	2,323,728	72.43%
Divorced	8,428,145	1,942,388	23.05%	14.67%	31.43%	6,485,757	76.95%
Separated	2,134,533	423,395	19.84%	12.51%	27.16%	1,711,137	80.16%
Never Married	12,388,270	818,909	6.61%	4.25%	8.97%	11,569,361	93.39%
Living with Partner	6,649,894	1,013,788	15.25%	7.27%	23.22%	5,636,106	84.75%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	6,832,632	1,314,414	19.24%	13.86%	24.62%	5,518,218	80.76%
Poverty income ratio less than 1	11,430,626	1,579,500	13.82%	9.00%	18.64%	9,851,125	86.18%
Poverty income ratio 1 or greater	76,669,128	11,981,554	15.63%	13.26%	18.00%	64,687,574	84.37%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	808,899	211,693	26.17%	11.32%	41.02%	597,206	73.83%
Normal: less than 25	26,748,202	3,280,900	12.27%	9.54%	14.99%	23,467,302	87.73%
Overweight: 25.0 - 29.9	36,919,637	5,264,464	14.26%	10.48%	18.03%	31,655,173	85.74%
Obese: 30 or higher	30,455,646	6,118,411	20.09%	17.10%	23.08%	24,337,235	79.91%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	94,932,386	14,875,469	15.67%	13.41%	17.93%	80,056,917	84.33%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,550,279	182,953	5.15%	2.49%	7.82%	3,367,326	94.85%
Glycohemoglobin < 5.7%	62,899,045	7,603,718	12.09%	9.18%	15.00%	55,295,326	87.91%
Glycohemoglobin 5.7% - 6.4%	21,210,133	5,052,896	23.82%	18.22%	29.42%	16,157,237	76.18%
Glycohemoglobin 6.5% or higher	7,272,929	2,035,901	27.99%	18.96%	37.03%	5,237,028	72.01%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	8,457,156	2,791,313	33.01%	25.48%	40.53%	5,665,843	66.99%
No	86,475,230	12,084,156	13.97%	11.55%	16.40%	74,391,074	86.03%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,071,877	9,678,148	18.59%	15.76%	21.41%	42,393,729	81.41%
No	42,792,655	5,159,491	12.06%	9.06%	15.05%	37,633,163	87.94%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,230,043	7,126,875	25.25%	21.39%	29.10%	21,103,168	74.75%
No	66,599,730	7,748,593	11.63%	9.47%	13.80%	58,851,137	88.37%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	8,986,169	2,826,339	31.45%	25.01%	37.90%	6,159,829	68.55%
Mobility limitations	6,860,598	2,377,127	34.65%	28.36%	40.94%	4,483,471	65.35%
No mobility limitations	18,940,648	3,828,510	20.21%	16.78%	23.64%	15,112,138	79.79%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	60,107,142	5,805,663	9.66%	6.79%	12.53%	54,301,478	90.34%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,814,272	1,782,327	22.81%	12.26%	33.36%	6,031,945	77.19%
No	86,905,951	13,063,306	15.03%	12.67%	17.39%	73,842,645	84.97%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	610,540	117,400	19.23%	0.95%	37.51%	493,141	80.77%
2 or more times a day	10,308,368	1,880,197	18.24%	11.85%	24.63%	8,428,171	81.76%
Once a day	8,264,954	1,620,551	19.61%	12.86%	26.36%	6,644,403	80.39%
2 or more time a week	5,908,062	1,385,683	23.45%	16.93%	29.98%	4,522,379	76.55%
Once a week	4,371,150	844,703	19.32%	10.88%	27.76%	3,526,447	80.68%
1-3 times a month	8,315,400	2,186,284	26.29%	18.29%	34.29%	6,129,116	73.71%
Never	56,772,742	6,823,374	12.02%	9.82%	14.22%	49,949,368	87.98%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	610,540	117,400	19.23%	0.95%	37.51%	493,141	80.77%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	1,037,565	566,107	54.56%	28.99%	80.14%	471,458	45.44%
Never	92,548,070	14,035,507	15.17%	13.11%	17.22%	78,512,563	84.83%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).



Table M01 -Prevalence of any urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	95,585,798	15,475,302	16.19%	14.32%	18.06%	80,110,496	83.81%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	297,636	41,692	14.01%	(10.98%)	38.99%	255,943	85.99%
20-29	12,944,492	827,931	6.40%	3.15%	9.64%	12,116,561	93.60%
30-39	16,576,959	1,431,660	8.64%	4.63%	12.65%	15,145,299	91.36%
40-49	21,324,506	2,979,005	13.97%	9.24%	18.70%	18,345,501	86.03%
50-59	21,456,557	3,262,484	15.21%	11.78%	18.63%	18,194,073	84.79%
60-64	7,051,206	1,986,235	28.17%	18.79%	37.55%	5,064,971	71.83%
65-69	6,200,612	1,488,541	24.01%	19.21%	28.80%	4,712,071	75.99%
70-74	3,763,064	1,114,946	29.63%	22.71%	36.55%	2,648,118	70.37%
75-79	2,936,105	976,453	33.26%	20.41%	46.11%	1,959,652	66.74%
80+	3,034,662	1,366,355	45.02%	38.03%	52.02%	1,668,307	54.98%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	9,079,208	1,243,153	13.69%	10.72%	16.66%	7,836,055	86.31%
Other Hispanic	4,899,222	611,745	12.49%	7.37%	17.60%	4,287,477	87.51%
Non-Hispanic White	66,118,840	10,871,907	16.44%	14.14%	18.75%	55,246,933	83.56%
Non-Hispanic Black	9,655,413	2,146,928	22.24%	14.41%	30.06%	7,508,485	77.76%
Other Race - Including Multi-Racial	5,833,115	601,568	10.31%	2.78%	17.84%	5,231,547	89.69%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,359,443	300,834	22.13%	7.75%	36.51%	1,058,609	77.87%
Less Than 9th Grade	6,477,166	1,222,885	18.88%	14.02%	23.74%	5,254,282	81.12%
9-11th Grade	10,241,048	1,874,404	18.30%	12.96%	23.64%	8,366,643	81.70%
High School Grad/GED or Equivalent	77,361,680	12,077,179	15.61%	13.73%	17.49%	65,284,500	84.39%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	313,184	57,241	18.28%	( 8.62%)	45.17%	255,943	81.72%
Married	61,053,634	10,682,851	17.50%	14.67%	20.33%	50,370,783	82.50%
Widowed	3,241,681	569,758	17.58%	8.97%	26.18%	2,671,923	82.42%
Divorced	8,826,206	1,496,763	16.96%	10.59%	23.32%	7,329,443	83.04%
Separated	2,184,790	210,011	9.61%	1.77%	17.46%	1,974,780	90.39%
Never Married	12,431,390	1,602,517	12.89%	7.49%	18.29%	10,828,872	87.11%
Living with Partner	7,478,020	810,924	10.84%	6.70%	14.99%	6,667,095	89.16%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	6,719,530	1,455,121	21.66%	14.26%	29.05%	5,264,409	78.34%
Poverty income ratio less than 1	10,991,125	1,596,007	14.52%	10.49%	18.55%	9,395,117	85.48%
Poverty income ratio 1 or greater	77,875,143	12,424,174	15.95%	13.97%	17.94%	65,450,969	84.05%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	914,576	292,076	31.94%	5.01%	58.86%	622,500	68.06%
Normal: less than 25	24,656,831	4,066,463	16.49%	12.71%	20.27%	20,590,367	83.51%
Overweight: 25.0 - 29.9	36,228,379	4,788,091	13.22%	10.84%	15.59%	31,440,288	86.78%
Obese: 30 or higher	33,786,012	6,328,671	18.73%	16.57%	20.90%	27,457,341	81.27%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	2,643,943	169,169	6.40%	1.04%	11.76%	2,474,774	93.60%
Glycohemoglobin < 5.7%	61,620,636	8,629,123	14.00%	11.23%	16.78%	52,991,513	86.00%
Glycohemoglobin 5.7% - 6.4%	23,078,335	4,513,678	19.56%	16.23%	22.88%	18,564,657	80.44%
Glycohemoglobin 6.5% or higher	8,242,884	2,163,331	26.24%	21.90%	30.59%	6,079,552	73.76%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).

Table M01 -Prevalence of any urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	94,932,386	14,875,469	15.67%	13.41%	17.93%	80,056,917	84.33%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	8,780,629	2,480,826	28.25%	19.61%	36.90%	6,299,804	71.75%
No	86,805,168	12,994,476	14.97%	12.81%	17.13%	73,810,692	85.03%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	47,479,384	8,875,386	18.69%	16.20%	21.19%	38,603,998	81.31%
No	48,106,414	6,599,916	13.72%	10.52%	16.92%	41,506,498	86.28%
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,857,116	6,397,998	22.17%	18.86%	25.48%	22,459,118	77.83%
No	66,617,365	9,066,420	13.61%	11.83%	15.39%	57,550,945	86.39%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	7,924,746	2,512,390	31.70%	23.56%	39.85%	5,412,356	68.30%
Mobility limitations	6,892,982	2,045,907	29.68%	22.65%	36.71%	4,847,075	70.32%
No mobility limitations	20,034,171	4,531,815	22.62%	18.15%	27.10%	15,502,355	77.38%
N/A - Respondent not asked mobility limitation questions	60,733,899	6,385,189	10.51%	8.16%	12.87%	54,348,710	89.49%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,676,778	1,963,555	25.58%	15.78%	35.37%	5,713,222	74.42%
No	87,705,670	13,504,920	15.40%	13.40%	17.40%	74,200,750	84.60%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	459,339	254,678	55.44%	18.74%	92.15%	204,661	44.56%
2 or more times a day	9,167,832	1,691,877	18.45%	11.78%	25.13%	7,475,956	81.55%
Once a day	8,770,661	1,484,183	16.92%	9.60%	24.24%	7,286,479	83.08%
2 or more time a week	6,773,718	1,294,731	19.11%	11.21%	27.02%	5,478,987	80.89%
Once a week	4,927,442	1,010,064	20.50%	12.89%	28.10%	3,917,378	79.50%
1-3 times a month	10,278,875	2,725,838	26.52%	21.46%	31.58%	7,553,037	73.48%
Never	55,144,254	7,013,931	12.72%	10.67%	14.77%	48,130,323	87.28%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	459,339	254,678	55.44%	18.74%	92.15%	204,661	44.56%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	844,643	466,082	55.18%	33.06%	77.30%	378,562	44.82%
Never	93,745,910	14,575,578	15.55%	13.70%	17.39%	79,170,332	84.45%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042, KIQ.044, and KIQ.046 (stress urinary incontinence, urge incontinence, and insensate urine loss).