

Table M02 - Prevalence of stress urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,358,261	4,130,306	4.57%	3.80%	5.34%	86,227,955	95.43%
Age at screening	-	-	-	-	-	-	-
10-19	318,425	7,364	2.31%	(2.96%)	7.59%	311,061	97.69%
20-29	11,767,942	64,473	0.55%	(0.59%)	1.69%	11,703,469	99.45%
30-39	18,212,868	354,527	1.95%	(0.04%)	3.94%	17,858,341	98.05%
40-49	22,595,280	485,985	2.15%	0.24%	4.06%	22,109,296	97.85%
50-59	18,568,743	791,153	4.26%	1.50%	7.02%	17,777,591	95.74%
60-64	4,648,416	436,506	9.39%	4.31%	14.47%	4,211,911	90.61%
65-69	3,884,538	423,659	10.91%	4.61%	17.20%	3,460,879	89.09%
70-74	4,039,191	677,633	16.78%	7.08%	26.47%	3,361,558	83.22%
75-79	2,934,099	375,114	12.78%	7.26%	18.31%	2,558,984	87.22%
80-84	2,102,740	269,155	12.80%	7.20%	18.40%	1,833,585	87.20%
85+	1,286,018	244,737	19.03%	8.14%	29.92%	1,041,281	80.97%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,055,890	139,968	1.98%	0.65%	3.32%	6,915,922	98.02%
Other Hispanic	4,966,199	241,602	4.86%	(0.14%)	9.87%	4,724,597	95.14%
Non-Hispanic White	66,204,355	3,244,599	4.90%	3.64%	6.16%	62,959,755	95.10%
Non-Hispanic Black	9,016,771	371,718	4.12%	0.94%	7.31%	8,645,053	95.88%
Other Race - Including Multi-Racial	3,115,046	132,419	4.25%	(1.21%)	9.71%	2,982,628	95.75%
Education	-	-	-	-	-	-	-
Missing	933,223	51,502	5.52%	(2.97%)	14.01%	881,721	94.48%
Less Than 9th Grade	6,516,375	685,491	10.52%	7.54%	13.50%	5,830,884	89.48%
9-11th Grade	10,170,827	552,669	5.43%	1.43%	9.43%	9,618,158	94.57%
High School Grad/GED or Equivalent	72,591,006	2,808,297	3.87%	2.93%	4.80%	69,782,709	96.13%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	909,023	141,372	15.55%	(4.08%)	35.18%	767,651	84.45%
Married	59,588,799	2,830,542	4.75%	3.58%	5.92%	56,758,257	95.25%
Widowed	4,052,855	354,058	8.74%	2.46%	15.01%	3,698,796	91.26%
Divorced	7,804,451	508,087	6.51%	3.17%	9.85%	7,296,364	93.49%
Separated	2,163,795	2,737	0.13%	(0.14%)	0.40%	2,161,058	99.87%
Never Married	10,212,194	97,764	0.96%	(0.15%)	2.07%	10,114,430	99.04%
Living with Partner	5,531,980	116,664	2.11%	(0.96%)	5.18%	5,415,316	97.89%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,980,213	230,956	4.64%	1.20%	8.08%	4,749,258	95.36%
Poverty income ratio less than 1	9,683,534	816,740	8.43%	4.05%	12.82%	8,866,794	91.57%
Poverty income ratio 1 or greater	75,694,513	3,082,610	4.07%	3.04%	5.11%	72,611,903	95.93%
Body mass index	-	-	-	-	-	-	-
Missing	3,185,619	509,070	15.98%	7.97%	23.99%	2,676,549	84.02%
Normal: less than 25	27,368,328	992,499	3.63%	2.10%	5.15%	26,375,829	96.37%
Overweight: 25.0 - 29.9	36,239,254	1,338,335	3.69%	2.07%	5.31%	34,900,918	96.31%
Obese: 30 or higher	23,565,060	1,290,401	5.48%	4.04%	6.92%	22,274,659	94.52%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,136,613	111,255	3.55%	1.06%	6.04%	3,025,358	96.45%
Glycohemoglobin < 5.7%	69,910,044	2,604,912	3.73%	2.80%	4.65%	67,305,131	96.27%
Glycohemoglobin 5.7% - 6.4%	11,200,635	566,138	5.05%	3.06%	7.04%	10,634,496	94.95%
Glycohemoglobin 6.5% or higher	6,110,970	848,000	13.88%	9.17%	18.58%	5,262,970	86.12%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,358,261	4,130,306	4.57%	3.80%	5.34%	86,227,955	95.43%
Diabetes	-	-	-	-	-	-	-
Yes	6,576,525	680,985	10.35%	4.13%	16.58%	5,895,540	89.65%
No	83,781,736	3,449,321	4.12%	3.25%	4.98%	80,332,415	95.88%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	51,225,896	3,164,894	6.18%	5.09%	7.27%	48,061,002	93.82%
No	39,045,204	965,412	2.47%	1.57%	3.38%	38,079,792	97.53%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	1,046,623	0	0.00%	0.00%	0.00%	1,046,623	100.00%
Yes	21,489,492	1,862,937	8.67%	6.99%	10.35%	19,626,555	91.33%
No	67,725,407	2,267,369	3.35%	2.23%	4.47%	65,458,038	96.65%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	5,308,418	836,611	15.76%	7.09%	24.43%	4,471,807	84.24%
Mobility limitations	6,896,108	971,264	14.08%	8.37%	19.80%	5,924,844	85.92%
No mobility limitations	18,439,144	1,464,916	7.94%	5.38%	10.51%	16,974,229	92.06%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	59,699,976	842,901	1.41%	0.58%	2.24%	58,857,075	98.59%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	90,358,261	4,130,306	4.57%	3.80%	5.34%	86,227,955	95.43%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	90,358,261	4,130,306	4.57%	3.80%	5.34%	86,227,955	95.43%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	90,358,261	4,130,306	4.57%	3.80%	5.34%	86,227,955	95.43%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,966,201	4,898,189	5.38%	3.85%	6.92%	86,068,011	94.62%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	538,199	0	0.00%	0.00%	0.00%	538,199	100.00%
20-29	12,110,825	118,862	0.98%	(0.20%)	2.17%	11,991,963	99.02%
30-39	17,705,119	591,943	3.34%	0.51%	6.18%	17,113,176	96.66%
40-49	20,941,149	462,569	2.21%	0.34%	4.07%	20,478,580	97.79%
50-59	17,826,163	940,602	5.28%	2.33%	8.22%	16,885,561	94.72%
60-64	5,666,205	379,692	6.70%	2.88%	10.53%	5,286,512	93.30%
65-69	5,762,216	440,514	7.64%	3.41%	11.88%	5,321,702	92.36%
70-74	3,988,818	637,943	15.99%	8.72%	23.26%	3,350,875	84.01%
75-79	3,030,438	597,566	19.72%	10.42%	29.02%	2,432,872	80.28%
80-84	2,043,274	409,854	20.06%	12.38%	27.74%	1,633,420	79.94%
85+	1,353,795	318,644	23.54%	9.20%	37.87%	1,035,151	76.46%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,316,409	200,674	2.74%	1.14%	4.35%	7,115,735	97.26%
Other Hispanic	2,977,019	96,716	3.25%	1.44%	5.06%	2,880,303	96.75%
Non-Hispanic White	66,896,781	3,876,085	5.79%	3.96%	7.63%	63,020,696	94.21%
Non-Hispanic Black	9,097,016	545,633	6.00%	3.74%	8.26%	8,551,383	94.00%
Other Race - Including Multi-Racial	4,678,976	179,081	3.83%	(1.36%)	9.02%	4,499,895	96.17%
Education	-	-	-	-	-	-	-
Missing	1,796,573	64,105	3.57%	(1.94%)	9.08%	1,732,468	96.43%
Less Than 9th Grade	6,526,491	491,040	7.52%	3.25%	11.80%	6,035,451	92.48%
9-11th Grade	8,965,501	640,498	7.14%	2.87%	11.41%	8,325,003	92.86%
High School Grad/GED or Equivalent	73,330,038	3,702,546	5.05%	3.60%	6.50%	69,627,492	94.95%
Refused	*	*	*	*	*	*	*
Don't Know	335,074	0	0.00%	0.00%	0.00%	335,074	100.00%
Marital status	-	-	-	-	-	-	-
Missing	1,019,622	20,487	2.01%	(1.41%)	5.43%	999,134	97.99%
Married	59,788,711	3,586,650	6.00%	4.13%	7.86%	56,202,061	94.00%
Widowed	3,595,195	329,023	9.15%	3.79%	14.52%	3,266,172	90.85%
Divorced	7,528,319	317,782	4.22%	0.14%	8.30%	7,210,538	95.78%
Separated	1,505,477	123,681	8.22%	0.53%	15.90%	1,381,796	91.78%
Never Married	11,175,776	444,983	3.98%	0.88%	7.09%	10,730,793	96.02%
Living with Partner	5,961,389	75,583	1.27%	(0.74%)	3.28%	5,885,805	98.73%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,481,624	389,885	8.70%	5.02%	12.38%	4,091,739	91.30%
Poverty income ratio less than 1	9,206,856	431,761	4.69%	1.88%	7.50%	8,775,095	95.31%
Poverty income ratio 1 or greater	77,277,721	4,076,543	5.28%	3.41%	7.14%	73,201,178	94.72%
Body mass index	-	-	-	-	-	-	-
Missing	960,921	162,865	16.95%	(0.07%)	33.96%	798,056	83.05%
Normal: less than 25	27,546,264	1,448,716	5.26%	3.26%	7.26%	26,097,548	94.74%
Overweight: 25.0 - 29.9	34,664,541	1,530,110	4.41%	2.74%	6.08%	33,134,432	95.59%
Obese: 30 or higher	27,794,474	1,756,498	6.32%	3.40%	9.24%	26,037,976	93.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,966,201	4,898,189	5.38%	3.85%	6.92%	86,068,011	94.62%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,412,745	78,565	3.26%	(0.60%)	7.11%	2,334,179	96.74%
Glycohemoglobin < 5.7%	69,628,405	3,145,109	4.52%	3.04%	6.00%	66,483,296	95.48%
Glycohemoglobin 5.7% - 6.4%	12,825,067	1,243,061	9.69%	6.02%	13.36%	11,582,006	90.31%
Glycohemoglobin 6.5% or higher	6,099,984	431,454	7.07%	3.32%	10.83%	5,668,531	92.93%
Diabetes	-	-	-	-	-	-	-
Yes	6,922,443	809,376	11.69%	6.68%	16.70%	6,113,067	88.31%
No	84,043,758	4,088,813	4.87%	3.45%	6.28%	79,954,945	95.13%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,601,967	3,435,919	6.53%	4.82%	8.24%	49,166,048	93.47%
No	38,326,021	1,450,439	3.78%	1.86%	5.71%	36,875,582	96.22%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	789,967	0	0.00%	0.00%	0.00%	789,967	100.00%
Yes	27,608,687	2,429,534	8.80%	5.77%	11.83%	25,179,153	91.20%
No	62,461,874	2,468,655	3.95%	2.68%	5.22%	59,993,219	96.05%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	9,336,477	1,288,667	13.80%	7.43%	20.18%	8,047,810	86.20%
Mobility limitations	7,792,349	962,469	12.35%	7.63%	17.08%	6,829,880	87.65%
No mobility limitations	19,566,611	1,665,782	8.51%	6.30%	10.73%	17,900,829	91.49%
N/A - Respondent not asked mobility limitation questions	54,270,763	981,271	1.81%	0.56%	3.06%	53,289,492	98.19%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	90,966,201	4,898,189	5.38%	3.85%	6.92%	86,068,011	94.62%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	90,966,201	4,898,189	5.38%	3.85%	6.92%	86,068,011	94.62%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	90,966,201	4,898,189	5.38%	3.85%	6.92%	86,068,011	94.62%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KI.Q.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,713,557	3,268,109	3.49%	2.39%	4.58%	90,445,448	96.51%
Age at screening	-	-	-	-	-	-	-
10-19	384,918	0	0.00%	0.00%	0.00%	384,918	100.00%
20-29	14,540,015	312,026	2.15%	0.20%	4.09%	14,227,990	97.85%
30-39	17,503,545	409,737	2.34%	0.23%	4.45%	17,093,808	97.66%
40-49	21,809,592	670,290	3.07%	0.94%	5.20%	21,139,302	96.93%
50-59	18,142,193	584,702	3.22%	0.99%	5.45%	17,557,492	96.78%
60-64	6,373,461	127,251	2.00%	(0.06%)	4.05%	6,246,211	98.00%
65-69	5,297,573	310,561	5.86%	1.48%	10.24%	4,987,012	94.14%
70-74	4,006,172	245,050	6.12%	2.89%	9.34%	3,761,122	93.88%
75-79	2,536,289	310,634	12.25%	0.35%	24.14%	2,225,656	87.75%
80-84	2,073,206	243,358	11.74%	1.83%	21.65%	1,829,848	88.26%
85+	1,046,592	54,502	5.21%	(0.97%)	11.38%	992,090	94.79%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,979,822	201,354	2.52%	0.59%	4.45%	7,778,468	97.48%
Other Hispanic	2,862,199	176,433	6.16%	0.26%	12.07%	2,685,765	93.84%
Non-Hispanic White	69,156,150	2,571,822	3.72%	2.28%	5.16%	66,584,328	96.28%
Non-Hispanic Black	9,553,122	243,985	2.55%	1.64%	3.47%	9,309,137	97.45%
Other Race - Including Multi-Racial	4,162,265	74,515	1.79%	(2.01%)	5.59%	4,087,750	98.21%
Education	-	-	-	-	-	-	-
Missing	1,125,917	21,315	1.89%	(2.21%)	5.99%	1,104,602	98.11%
Less Than 9th Grade	6,403,935	106,329	1.66%	0.26%	3.06%	6,297,605	98.34%
9-11th Grade	10,260,776	252,603	2.46%	1.02%	3.91%	10,008,173	97.54%
High School Grad/GED or Equivalent	75,757,393	2,874,984	3.79%	2.45%	5.14%	72,882,409	96.21%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,042,842	12,877	1.23%	(0.85%)	3.32%	1,029,964	98.77%
Married	60,157,438	2,203,266	3.66%	2.34%	4.98%	57,954,172	96.34%
Widowed	2,985,428	107,286	3.59%	0.00%	7.18%	2,878,141	96.41%
Divorced	9,188,198	387,918	4.22%	2.61%	5.83%	8,800,280	95.78%
Separated	2,712,975	130,888	4.82%	(3.96%)	13.61%	2,582,086	95.18%
Never Married	9,472,270	298,368	3.15%	0.58%	5.72%	9,173,902	96.85%
Living with Partner	8,115,581	127,504	1.57%	(0.84%)	3.98%	7,988,077	98.43%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	2,977,906	227,868	7.65%	0.66%	14.64%	2,750,038	92.35%
Poverty income ratio less than 1	9,079,370	294,772	3.25%	0.07%	6.42%	8,784,599	96.75%
Poverty income ratio 1 or greater	81,656,281	2,745,470	3.36%	2.13%	4.59%	78,910,811	96.64%
Body mass index	-	-	-	-	-	-	-
Missing	1,009,735	33,677	3.34%	(1.58%)	8.25%	976,058	96.66%
Normal: less than 25	25,820,546	502,632	1.95%	1.30%	2.59%	25,317,914	98.05%
Overweight: 25.0 - 29.9	35,984,960	906,923	2.52%	0.99%	4.05%	35,078,037	97.48%
Obese: 30 or higher	30,898,317	1,824,878	5.91%	3.80%	8.01%	29,073,440	94.09%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,564,155	26,585	0.75%	(0.86%)	2.35%	3,537,570	99.25%
Glycohemoglobin < 5.7%	70,325,447	1,691,844	2.41%	1.47%	3.34%	68,633,602	97.59%
Glycohemoglobin 5.7% - 6.4%	14,222,346	931,035	6.55%	3.62%	9.48%	13,291,311	93.45%
Glycohemoglobin 6.5% or higher	5,601,610	618,645	11.04%	3.64%	18.45%	4,982,964	88.96%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,713,557	3,268,109	3.49%	2.39%	4.58%	90,445,448	96.51%
Diabetes	-	-	-	-	-	-	-
Yes	6,979,396	620,250	8.89%	4.10%	13.68%	6,359,145	91.11%
No	86,734,162	2,647,859	3.05%	1.94%	4.16%	84,086,303	96.95%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	53,038,544	2,228,678	4.20%	2.40%	6.01%	50,809,866	95.80%
No	40,675,013	1,039,431	2.56%	1.47%	3.64%	39,635,582	97.44%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	26,882,711	1,725,907	6.42%	4.55%	8.29%	25,156,804	93.58%
No	66,652,383	1,542,202	2.31%	1.17%	3.45%	65,110,181	97.69%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,937,459	489,894	7.06%	2.53%	11.59%	6,447,564	92.94%
Mobility limitations	5,930,657	498,795	8.41%	4.47%	12.35%	5,431,861	91.59%
No mobility limitations	20,291,603	865,811	4.27%	1.44%	7.10%	19,425,792	95.73%
N/A - Respondent not asked mobility limitation questions	60,553,839	1,413,608	2.33%	1.30%	3.37%	59,140,231	97.67%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,936,022	523,387	6.60%	1.98%	11.21%	7,412,635	93.40%
No	85,661,432	2,744,722	3.20%	2.13%	4.28%	82,916,710	96.80%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	372,256	22,521	6.05%	(3.06%)	15.16%	349,735	93.95%
2 or more times a day	13,806,596	730,645	5.29%	1.96%	8.62%	13,075,951	94.71%
Once a day	8,509,610	297,195	3.49%	(0.24%)	7.23%	8,212,415	96.51%
2 or more time a week	5,956,076	448,774	7.53%	2.96%	12.11%	5,507,302	92.47%
Once a week	5,314,628	260,268	4.90%	(0.29%)	10.09%	5,054,360	95.10%
1-3 times a month	9,361,331	219,140	2.34%	0.94%	3.74%	9,142,192	97.66%
Never	50,100,244	1,289,566	2.57%	1.07%	4.08%	48,810,678	97.43%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	372,256	22,521	6.05%	(3.06%)	15.16%	349,735	93.95%
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	749,457	178,104	23.76%	0.12%	47.41%	571,353	76.24%
Never	92,195,030	3,050,314	3.31%	2.28%	4.34%	89,144,716	96.69%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,990,698	4,090,064	4.31%	3.06%	5.55%	90,900,633	95.69%
Age at screening	-	-	-	-	-	-	-
10-19	581,889	0	0.00%	0.00%	0.00%	581,889	100.00%
20-29	14,324,402	130,419	0.91%	0.15%	1.67%	14,193,983	99.09%
30-39	17,120,661	283,354	1.66%	0.23%	3.08%	16,837,307	98.34%
40-49	21,351,091	722,090	3.38%	0.25%	6.52%	20,629,002	96.62%
50-59	18,864,707	1,056,982	5.60%	3.30%	7.91%	17,807,725	94.40%
60-64	7,469,087	461,531	6.18%	2.31%	10.04%	7,007,555	93.82%
65-69	5,293,210	416,020	7.86%	1.94%	13.78%	4,877,190	92.14%
70-74	3,776,339	404,387	10.71%	5.18%	16.24%	3,371,953	89.29%
75-79	2,931,444	306,493	10.46%	5.60%	15.31%	2,624,951	89.54%
80+	3,277,867	308,789	9.42%	4.44%	14.40%	2,969,079	90.58%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,009,819	180,962	2.01%	0.63%	3.38%	8,828,857	97.99%
Other Hispanic	4,540,869	118,120	2.60%	(0.23%)	5.43%	4,422,750	97.40%
Non-Hispanic White	65,930,379	2,812,432	4.27%	2.89%	5.64%	63,117,947	95.73%
Non-Hispanic Black	9,832,078	463,002	4.71%	2.39%	7.03%	9,369,076	95.29%
Other Race - Including Multi-Racial	5,677,553	515,549	9.08%	(0.85%)	19.01%	5,162,003	90.92%
Education	-	-	-	-	-	-	-
Missing	1,275,540	44,574	3.49%	(0.95%)	7.94%	1,230,967	96.51%
Less Than 9th Grade	6,695,497	454,954	6.79%	3.34%	10.25%	6,240,542	93.21%
9-11th Grade	11,906,442	535,298	4.50%	2.43%	6.56%	11,371,144	95.50%
High School Grad/GED or Equivalent	74,976,791	3,055,238	4.07%	2.78%	5.37%	71,921,554	95.93%
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,282,066	0	0.00%	0.00%	0.00%	1,282,066	100.00%
Married	60,689,258	3,135,187	5.17%	3.23%	7.10%	57,554,071	94.83%
Widowed	3,222,890	162,351	5.04%	1.55%	8.52%	3,060,539	94.96%
Divorced	8,438,415	173,201	2.05%	0.69%	3.42%	8,265,214	97.95%
Separated	2,134,533	212,697	9.96%	2.19%	17.74%	1,921,836	90.04%
Never Married	12,388,270	193,629	1.56%	0.32%	2.80%	12,194,641	98.44%
Living with Partner	6,649,894	213,000	3.20%	0.80%	5.61%	6,436,894	96.80%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,832,632	528,158	7.73%	4.29%	11.17%	6,304,473	92.27%
Poverty income ratio less than 1	11,439,603	602,614	5.27%	2.20%	8.34%	10,836,989	94.73%
Poverty income ratio 1 or greater	76,718,463	2,959,291	3.86%	2.64%	5.07%	73,759,171	96.14%
Body mass index	-	-	-	-	-	-	-
Missing	819,170	50,034	6.11%	(1.61%)	13.83%	769,136	93.89%
Normal: less than 25	26,748,202	896,037	3.35%	1.85%	4.85%	25,852,166	96.65%
Overweight: 25.0 - 29.9	36,934,332	1,505,040	4.07%	1.77%	6.38%	35,429,292	95.93%
Obese: 30 or higher	30,488,993	1,638,954	5.38%	2.97%	7.78%	28,850,040	94.62%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,574,649	93,150	2.61%	1.06%	4.15%	3,481,499	97.39%
Glycohemoglobin < 5.7%	62,913,740	1,957,517	3.11%	1.73%	4.50%	60,956,222	96.89%
Glycohemoglobin 5.7% - 6.4%	21,210,133	1,410,909	6.65%	4.64%	8.66%	19,799,224	93.35%
Glycohemoglobin 6.5% or higher	7,292,177	628,489	8.62%	3.43%	13.81%	6,663,688	91.38%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,990,698	4,090,064	4.31%	3.06%	5.55%	90,900,633	95.69%
Diabetes	-	-	-	-	-	-	-
Yes	8,491,796	573,852	6.76%	3.33%	10.19%	7,917,944	93.24%
No	86,498,902	3,516,212	4.07%	2.79%	5.34%	82,982,690	95.93%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,105,819	2,827,737	5.43%	4.05%	6.80%	49,278,083	94.57%
No	42,817,024	1,262,328	2.95%	1.37%	4.53%	41,554,697	97.05%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,240,314	2,004,288	7.10%	4.56%	9.63%	26,236,025	92.90%
No	66,647,772	2,085,776	3.13%	1.92%	4.34%	64,561,996	96.87%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	8,996,439	1,009,901	11.23%	7.22%	15.23%	7,986,537	88.77%
Mobility limitations	6,884,968	898,681	13.05%	6.67%	19.44%	5,986,286	86.95%
No mobility limitations	18,964,320	826,673	4.36%	2.25%	6.47%	18,137,647	95.64%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	60,107,142	1,354,808	2.25%	0.92%	3.59%	58,752,333	97.75%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,838,641	707,764	9.03%	4.60%	13.46%	7,130,877	90.97%
No	86,939,893	3,352,464	3.86%	2.66%	5.05%	83,587,429	96.14%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	625,235	37,943	6.07%	(2.18%)	14.31%	587,292	93.93%
2 or more times a day	10,308,368	617,089	5.99%	2.10%	9.88%	9,691,279	94.01%
Once a day	8,264,954	202,074	2.44%	0.44%	4.45%	8,062,880	97.56%
2 or more time a week	5,908,062	515,090	8.72%	4.66%	12.77%	5,392,972	91.28%
Once a week	4,371,150	339,144	7.76%	0.97%	14.55%	4,032,006	92.24%
1-3 times a month	8,315,400	613,448	7.38%	2.42%	12.33%	7,701,952	92.62%
Never	56,816,359	1,765,277	3.11%	1.57%	4.64%	55,051,082	96.89%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	625,235	37,943	6.07%	(2.18%)	14.31%	587,292	93.93%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	1,037,565	272,523	26.27%	(3.04%)	55.57%	765,042	73.73%
Never	92,581,417	3,740,622	4.04%	2.74%	5.34%	88,840,796	95.96%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,576,022	3,711,449	3.88%	3.27%	4.49%	91,864,574	96.12%
Age at screening	-	-	-	-	-	-	-
10-19	297,636	0	0.00%	0.00%	0.00%	297,636	100.00%
20-29	12,944,492	135,101	1.04%	0.02%	2.06%	12,809,392	98.96%
30-39	16,576,959	356,927	2.15%	0.35%	3.95%	16,220,032	97.85%
40-49	21,324,506	494,604	2.32%	0.88%	3.76%	20,829,902	97.68%
50-59	21,456,557	986,089	4.60%	2.55%	6.64%	20,470,468	95.40%
60-64	7,041,431	195,866	2.78%	(0.72%)	6.28%	6,845,565	97.22%
65-69	6,200,612	512,329	8.26%	3.86%	12.67%	5,688,283	91.74%
70-74	3,763,064	338,057	8.98%	3.26%	14.71%	3,425,007	91.02%
75-79	2,936,105	258,006	8.79%	2.28%	15.29%	2,678,099	91.21%
80+	3,034,662	434,470	14.32%	7.96%	20.68%	2,600,192	85.68%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,079,208	156,697	1.73%	0.37%	3.09%	8,922,511	98.27%
Other Hispanic	4,899,222	171,577	3.50%	0.94%	6.07%	4,727,645	96.50%
Non-Hispanic White	66,118,840	2,778,872	4.20%	3.13%	5.28%	63,339,967	95.80%
Non-Hispanic Black	9,645,637	359,280	3.72%	2.47%	4.98%	9,286,358	96.28%
Other Race - Including Multi-Racial	5,833,115	245,023	4.20%	(0.01%)	8.41%	5,588,092	95.80%
Education	-	-	-	-	-	-	-
Missing	1,359,443	70,472	5.18%	(3.98%)	14.35%	1,288,971	94.82%
Less Than 9th Grade	6,477,166	288,264	4.45%	2.76%	6.14%	6,188,902	95.55%
9-11th Grade	10,241,048	330,306	3.23%	0.77%	5.68%	9,910,742	96.77%
High School Grad/GED or Equivalent	77,351,904	3,022,407	3.91%	3.02%	4.79%	74,329,497	96.09%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	313,184	15,549	4.96%	(5.58%)	15.51%	297,636	95.04%
Married	61,053,634	2,535,087	4.15%	3.32%	4.98%	58,518,547	95.85%
Widowed	3,241,681	168,108	5.19%	0.66%	9.71%	3,073,573	94.81%
Divorced	8,826,206	415,297	4.71%	0.81%	8.60%	8,410,909	95.29%
Separated	2,184,790	62,760	2.87%	(0.90%)	6.65%	2,122,031	97.13%
Never Married	12,421,614	350,607	2.82%	1.06%	4.59%	12,071,007	97.18%
Living with Partner	7,478,020	135,851	1.82%	0.31%	3.32%	7,342,169	98.18%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,719,530	355,714	5.29%	0.48%	10.11%	6,363,816	94.71%
Poverty income ratio less than 1	10,981,349	368,611	3.36%	1.96%	4.76%	10,612,738	96.64%
Poverty income ratio 1 or greater	77,875,143	2,987,123	3.84%	2.82%	4.85%	74,888,020	96.16%
Body mass index	-	-	-	-	-	-	-
Missing	914,576	58,215	6.37%	1.84%	10.89%	856,361	93.63%
Normal: less than 25	24,656,831	1,260,091	5.11%	3.60%	6.62%	23,396,739	94.89%
Overweight: 25.0 - 29.9	36,218,604	1,075,739	2.97%	1.91%	4.03%	35,142,865	97.03%
Obese: 30 or higher	33,786,012	1,317,404	3.90%	2.08%	5.72%	32,468,608	96.10%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,643,943	94,554	3.58%	(1.65%)	8.80%	2,549,389	96.42%
Glycohemoglobin < 5.7%	61,620,636	1,903,851	3.09%	2.27%	3.91%	59,716,785	96.91%
Glycohemoglobin 5.7% - 6.4%	23,068,560	1,274,404	5.52%	3.49%	7.55%	21,794,156	94.48%
Glycohemoglobin 6.5% or higher	8,242,884	438,640	5.32%	2.18%	8.46%	7,804,243	94.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?

Table M02 - Prevalence of stress urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,576,022	3,711,449	3.88%	3.27%	4.49%	91,864,574	96.12%
Diabetes	-	-	-	-	-	-	-
Yes	8,780,629	554,195	6.31%	1.92%	10.71%	8,226,435	93.69%
No	86,795,393	3,157,254	3.64%	3.04%	4.23%	83,638,139	96.36%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	47,479,384	2,220,224	4.68%	3.35%	6.00%	45,259,160	95.32%
No	48,096,639	1,491,225	3.10%	1.93%	4.27%	46,605,414	96.90%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,847,341	1,523,871	5.28%	3.97%	6.59%	27,323,469	94.72%
No	66,617,365	2,187,578	3.28%	2.77%	3.80%	64,429,787	96.72%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	7,914,971	645,338	8.15%	4.58%	11.72%	7,269,632	91.85%
Mobility limitations	6,892,982	560,493	8.13%	4.93%	11.34%	6,332,489	91.87%
No mobility limitations	20,034,171	1,163,829	5.81%	3.74%	7.88%	18,870,342	94.19%
N/A - Respondent not asked mob. lim. questions	60,733,899	1,341,788	2.21%	1.38%	3.03%	59,392,111	97.79%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,676,778	416,679	5.43%	0.66%	10.19%	7,260,099	94.57%
No	87,695,895	3,294,770	3.76%	3.12%	4.39%	84,401,125	96.24%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	459,339	48,387	10.53%	(1.92%)	22.99%	410,952	89.47%
2 or more times a day	9,167,832	488,526	5.33%	2.18%	8.47%	8,679,307	94.67%
Once a day	8,760,886	411,361	4.70%	1.42%	7.97%	8,349,525	95.30%
2 or more time a week	6,773,718	273,003	4.03%	0.74%	7.33%	6,500,715	95.97%
Once a week	4,927,442	360,744	7.32%	0.07%	14.57%	4,566,698	92.68%
1-3 times a month	10,278,875	625,817	6.09%	2.85%	9.33%	9,653,058	93.91%
Never	55,144,254	1,503,611	2.73%	1.80%	3.65%	53,640,643	97.27%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	459,339	48,387	10.53%	(1.92%)	22.99%	410,952	89.47%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	844,643	188,067	22.27%	7.05%	37.49%	656,577	77.73%
Never	93,736,134	3,444,825	3.68%	3.01%	4.34%	90,291,309	96.32%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.042: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with activity like coughing, lifting, or exercise?