

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	485,985	0	0.00%	21,240	4.37%	221,677	45.61%	243,067	50.02%	0	0.00%	0	0.00%
50-59	791,153	160,286	20.26%	148,038	18.71%	255,014	32.23%	227,815	28.80%	0	0.00%	0	0.00%
60-64	436,506	151,026	34.60%	11,073	2.54%	147,101	33.70%	127,306	29.16%	0	0.00%	0	0.00%
65-69	423,659	106,145	25.05%	165,630	39.10%	74,517	17.59%	77,368	18.26%	0	0.00%	0	0.00%
70-74	677,633	153,174	22.60%	138,483	20.44%	278,840	41.15%	107,136	15.81%	0	0.00%	0	0.00%
75-79	375,114	166,591	44.41%	115,269	30.73%	64,639	17.23%	28,616	7.63%	0	0.00%	0	0.00%
80-84	269,155	87,901	32.66%	109,918	40.84%	71,336	26.50%	0	0.00%	0	0.00%	0	0.00%
85+	244,737	116,670	47.67%	68,160	27.85%	49,454	20.21%	10,453	4.27%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	139,968	52,311	37.37%	7,818	5.59%	42,104	30.08%	37,736	26.96%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	3,244,599	802,955	24.75%	643,250	19.83%	1,069,046	32.95%	729,349	22.48%	0	0.00%	0	0.00%
Non-Hispanic Black	371,718	64,188	17.27%	83,322	22.42%	150,155	40.39%	74,053	19.92%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	685,491	295,875	43.16%	65,633	9.57%	240,079	35.02%	83,904	12.24%	0	0.00%	0	0.00%
9-11th Grade	552,669	96,400	17.44%	105,704	19.13%	230,733	41.75%	119,832	21.68%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	2,808,297	561,634	20.00%	574,126	20.44%	791,946	28.20%	880,591	31.36%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,830,542	778,438	27.50%	642,165	22.69%	708,322	25.02%	701,617	24.79%	0	0.00%	0	0.00%
Widowed	354,058	141,051	39.84%	42,585	12.03%	143,433	40.51%	26,989	7.62%	0	0.00%	0	0.00%
Divorced	508,087	19,566	3.85%	49,736	9.79%	240,380	47.31%	198,404	39.05%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	*	*	*	*	*	*	*	*	*	*	*	*	*
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	816,740	61,547	7.54%	109,803	13.44%	188,914	23.13%	456,476	55.89%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	3,082,610	911,768	29.58%	579,383	18.80%	997,439	32.36%	594,020	19.27%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	509,070	195,346	38.37%	150,181	29.50%	123,653	24.29%	39,890	7.84%	0	0.00%	0	0.00%
Normal: less than 25	992,499	245,975	24.78%	174,702	17.60%	230,849	23.26%	340,974	34.36%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,338,335	311,920	23.31%	355,065	26.53%	490,863	36.68%	180,487	13.49%	0	0.00%	0	0.00%
Obese: 30 or higher	1,290,401	225,121	17.45%	97,862	7.58%	444,443	34.44%	522,976	40.53%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	2,604,912	674,309	25.89%	482,612	18.53%	665,861	25.56%	782,131	30.03%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	566,138	166,766	29.46%	108,181	19.11%	246,025	43.46%	45,167	7.98%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	848,000	88,057	10.38%	153,571	18.11%	377,921	44.57%	228,451	26.94%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	680,985	140,902	20.69%	133,642	19.62%	406,441	59.68%	0	0.00%	0	0.00%	0	0.00%
No	3,449,321	837,461	24.28%	644,167	18.68%	883,366	25.61%	1,084,327	31.44%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,164,894	692,941	21.89%	608,830	19.24%	954,102	30.15%	909,021	28.72%	0	0.00%	0	0.00%
No	965,412	285,422	29.56%	168,979	17.50%	335,705	34.77%	175,305	18.16%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,862,937	537,160	28.83%	246,273	13.22%	634,193	34.04%	445,311	23.90%	0	0.00%	0	0.00%
No	2,267,369	441,203	19.46%	531,537	23.44%	655,614	28.92%	639,016	28.18%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	836,611	156,629	18.72%	104,331	12.47%	338,297	40.44%	237,354	28.37%	0	0.00%	0	0.00%
Mobility limitations	971,264	444,942	45.81%	200,507	20.64%	270,211	27.82%	55,604	5.72%	0	0.00%	0	0.00%
No mobility limitations	1,464,916	261,122	17.83%	400,794	27.36%	391,236	26.71%	411,763	28.11%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	842,901	101,056	11.99%	72,178	8.56%	290,062	34.41%	379,605	45.04%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,130,306	978,363	23.69%	777,810	18.83%	1,289,807	31.23%	1,084,327	26.25%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	591,943	34,732	5.87%	96,763	16.35%	117,980	19.93%	342,469	57.86%	0	0.00%	0	0.00%
40-49	462,569	9,367	2.02%	192,456	41.61%	130,056	28.12%	130,691	28.25%	0	0.00%	0	0.00%
50-59	940,602	70,898	7.54%	122,665	13.04%	75,387	8.01%	671,652	71.41%	0	0.00%	0	0.00%
60-64	379,692	96,573	25.43%	57,519	15.15%	33,643	8.86%	191,958	50.56%	0	0.00%	0	0.00%
65-69	440,514	86,069	19.54%	149,493	33.94%	169,053	38.38%	35,898	8.15%	0	0.00%	0	0.00%
70-74	637,943	120,492	18.89%	355,067	55.66%	113,765	17.83%	48,619	7.62%	0	0.00%	0	0.00%
75-79	597,566	101,210	16.94%	121,778	20.38%	308,233	51.58%	66,345	11.10%	0	0.00%	0	0.00%
80-84	409,854	214,883	52.43%	0	0.00%	106,351	25.95%	88,620	21.62%	0	0.00%	0	0.00%
85+	318,644	116,085	36.43%	63,286	19.86%	77,094	24.19%	40,712	12.78%	21,468	6.74%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	200,674	42,681	21.27%	89,551	44.63%	59,812	29.81%	8,629	4.30%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	3,876,085	722,655	18.64%	946,949	24.43%	829,907	21.41%	1,355,106	34.96%	21,468	0.55%	0	0.00%
Non-Hispanic Black	545,633	57,964	10.62%	125,528	23.01%	206,091	37.77%	156,051	28.60%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	491,040	169,791	34.58%	225,896	46.00%	63,207	12.87%	32,146	6.55%	0	0.00%	0	0.00%
9-11th Grade	640,498	121,967	19.04%	172,295	26.90%	181,322	28.31%	164,915	25.75%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,702,546	540,343	14.59%	792,805	21.41%	928,027	25.06%	1,419,903	38.35%	21,468	0.58%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	3,586,650	561,959	15.67%	823,763	22.97%	992,986	27.69%	1,186,474	33.08%	21,468	0.60%	0	0.00%
Widowed	329,023	234,450	71.26%	38,754	11.78%	35,042	10.65%	20,777	6.31%	0	0.00%	0	0.00%
Divorced	317,782	0	0.00%	179,312	56.43%	2,145	0.67%	136,325	42.90%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	444,983	50,615	11.37%	68,871	15.48%	121,896	27.39%	203,601	45.75%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	389,885	68,725	17.63%	80,555	20.66%	154,791	39.70%	64,345	16.50%	21,468	5.51%	0	0.00%
Poverty income ratio less than 1	431,761	85,631	19.83%	169,055	39.15%	119,798	27.75%	57,277	13.27%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	4,076,543	720,244	17.67%	962,991	23.62%	897,967	22.03%	1,495,341	36.68%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,448,716	266,371	18.39%	254,712	17.58%	536,197	37.01%	391,436	27.02%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,530,110	225,838	14.76%	535,680	35.01%	270,106	17.65%	498,487	32.58%	0	0.00%	0	0.00%
Obese: 30 or higher	1,756,498	279,444	15.91%	387,875	22.08%	353,384	20.12%	714,327	40.67%	21,468	1.22%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	3,145,109	578,523	18.39%	782,311	24.87%	601,394	19.12%	1,182,881	37.61%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,243,061	199,116	16.02%	275,359	22.15%	411,154	33.08%	335,964	27.03%	21,468	1.73%	0	0.00%
Glycohemoglobin 6.5% or higher	431,454	62,228	14.42%	152,471	35.34%	118,635	27.50%	98,119	22.74%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	809,376	166,115	20.52%	232,813	28.76%	230,039	28.42%	180,409	22.29%	0	0.00%	0	0.00%
No	4,088,813	708,484	17.33%	979,788	23.96%	942,517	23.05%	1,436,555	35.13%	21,468	0.53%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	3,435,919	671,274	19.54%	802,009	23.34%	798,058	23.23%	1,164,578	33.89%	0	0.00%	0	0.00%
No	1,450,439	191,495	13.20%	410,593	28.31%	374,498	25.82%	452,385	31.19%	21,468	1.48%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,429,534	410,179	16.88%	736,946	30.33%	605,208	24.91%	677,201	27.87%	0	0.00%	0	0.00%
No	2,468,655	464,420	18.81%	475,656	19.27%	567,349	22.98%	939,762	38.07%	21,468	0.87%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,288,667	487,951	37.86%	353,479	27.43%	192,449	14.93%	233,320	18.11%	21,468	1.67%	0	0.00%
Mobility limitations	962,469	174,151	18.09%	139,461	14.49%	394,848	41.02%	254,010	26.39%	0	0.00%	0	0.00%
No mobility limitations	1,665,782	177,766	10.67%	559,811	33.61%	393,106	23.60%	535,100	32.12%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	981,271	34,732	3.54%	159,851	16.29%	192,154	19.58%	594,534	60.59%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,898,189	874,600	17.86%	1,212,602	24.76%	1,172,556	23.94%	1,616,963	33.01%	21,468	0.44%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,268,109	116,336	3.56%	198,729	6.08%	923,970	28.27%	2,029,073	62.09%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	409,737	0	0.00%	32,002	7.81%	104,299	25.46%	273,435	66.73%	0	0.00%	0	0.00%
40-49	670,290	41,182	6.14%	0	0.00%	184,495	27.52%	444,613	66.33%	0	0.00%	0	0.00%
50-59	584,702	0	0.00%	0	0.00%	129,801	22.20%	454,901	77.80%	0	0.00%	0	0.00%
60-64	*	*	*	*	*	*	*	*	*	*	*	*	*
65-69	*	*	*	*	*	*	*	*	*	*	*	*	*
70-74	245,050	21,315	8.70%	35,316	14.41%	67,609	27.59%	120,811	49.30%	0	0.00%	0	0.00%
75-79	*	*	*	*	*	*	*	*	*	*	*	*	*
80-84	*	*	*	*	*	*	*	*	*	*	*	*	*
85+	*	*	*	*	*	*	*	*	*	*	*	*	*
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	201,354	3,907	1.94%	23,439	11.64%	46,283	22.99%	127,725	63.43%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	2,571,822	103,459	4.02%	122,579	4.77%	620,730	24.14%	1,725,054	67.08%	0	0.00%	0	0.00%
Non-Hispanic Black	243,985	8,970	3.68%	0	0.00%	71,568	29.33%	163,446	66.99%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*
9-11th Grade	252,603	0	0.00%	32,002	12.67%	116,073	45.95%	104,528	41.38%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	2,874,984	82,144	2.86%	144,393	5.02%	795,332	27.66%	1,853,116	64.46%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,268,109	116,336	3.56%	198,729	6.08%	923,970	28.27%	2,029,073	62.09%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,203,266	40,962	1.86%	78,701	3.57%	651,666	29.58%	1,431,938	64.99%	0	0.00%	0	0.00%
Widowed	*	*	*	*	*	*	*	*	*	*	*	*	*
Divorced	387,918	41,182	10.62%	0	0.00%	111,085	28.64%	235,652	60.75%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	*	*	*	*	*	*	*	*	*	*	*	*	*
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio 1 or greater	2,745,470	116,336	4.24%	90,522	3.30%	770,775	28.07%	1,767,837	64.39%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	502,632	8,970	1.78%	23,341	4.64%	137,831	27.42%	332,490	66.15%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	906,923	0	0.00%	79,024	8.71%	404,652	44.62%	423,247	46.67%	0	0.00%	0	0.00%
Obese: 30 or higher	1,824,878	107,366	5.88%	96,365	5.28%	347,811	19.06%	1,273,336	69.78%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	1,691,844	50,152	2.96%	132,741	7.85%	644,038	38.07%	864,913	51.12%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	931,035	62,277	6.69%	22,335	2.40%	160,159	17.20%	686,264	73.71%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	618,645	3,907	0.63%	43,654	7.06%	119,773	19.36%	451,312	72.95%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	620,250	0	0.00%	43,654	7.04%	17,170	2.77%	559,427	90.19%	0	0.00%	0	0.00%
No	2,647,859	116,336	4.39%	155,076	5.86%	906,801	34.25%	1,469,646	55.50%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,268,109	116,336	3.56%	198,729	6.08%	923,970	28.27%	2,029,073	62.09%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,228,678	112,429	5.04%	99,921	4.48%	627,807	28.17%	1,388,521	62.30%	0	0.00%	0	0.00%
No	1,039,431	3,907	0.38%	98,808	9.51%	296,163	28.49%	640,552	61.63%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,725,907	66,184	3.83%	128,422	7.44%	447,906	25.95%	1,083,395	62.77%	0	0.00%	0	0.00%
No	1,542,202	50,152	3.25%	70,308	4.56%	476,064	30.87%	945,678	61.32%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	489,894	40,962	8.36%	32,002	6.53%	219,428	44.79%	197,503	40.32%	0	0.00%	0	0.00%
Mobility limitations	498,795	21,315	4.27%	43,709	8.76%	125,964	25.25%	307,808	61.71%	0	0.00%	0	0.00%
No mobility limitations	865,811	54,059	6.24%	58,520	6.76%	182,551	21.08%	570,681	65.91%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,413,608	0	0.00%	64,499	4.56%	396,028	28.02%	953,081	67.42%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	523,387	0	0.00%	32,927	6.29%	234,677	44.84%	255,783	48.87%	0	0.00%	0	0.00%
No	2,744,722	116,336	4.24%	165,803	6.04%	689,293	25.11%	1,773,290	64.61%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	730,645	0	0.00%	96,365	13.19%	170,618	23.35%	463,662	63.46%	0	0.00%	0	0.00%
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	448,774	0	0.00%	32,927	7.34%	164,471	36.65%	251,377	56.01%	0	0.00%	0	0.00%
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	1,289,566	66,184	5.13%	58,656	4.55%	346,792	26.89%	817,935	63.43%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,268,109	116,336	3.56%	198,729	6.08%	923,970	28.27%	2,029,073	62.09%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-		-		-		-		-		-	
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	3,050,314	116,336	3.81%	198,729	6.52%	852,270	27.94%	1,882,978	61.73%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,090,064	379,103	9.27%	573,732	14.03%	1,048,936	25.65%	2,055,051	50.24%	33,242	0.81%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	722,090	27,326	3.78%	155,643	21.55%	332,255	46.01%	206,865	28.65%	0	0.00%	0	0.00%
50-59	1,056,982	147,541	13.96%	119,156	11.27%	47,869	4.53%	742,416	70.24%	0	0.00%	0	0.00%
60-64	461,531	0	0.00%	49,589	10.74%	256,489	55.57%	155,453	33.68%	0	0.00%	0	0.00%
65-69	416,020	0	0.00%	59,951	14.41%	125,321	30.12%	230,748	55.47%	0	0.00%	0	0.00%
70-74	404,387	92,535	22.88%	34,654	8.57%	169,088	41.81%	98,784	24.43%	9,325	2.31%	0	0.00%
75-79	306,493	89,695	29.26%	45,406	14.81%	46,623	15.21%	124,769	40.71%	0	0.00%	0	0.00%
80+	308,789	22,006	7.13%	87,653	28.39%	23,147	7.50%	152,065	49.25%	23,917	7.75%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	180,962	19,199	10.61%	33,801	18.68%	21,069	11.64%	106,893	59.07%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	2,812,432	329,149	11.70%	305,074	10.85%	539,862	19.20%	1,605,105	57.07%	33,242	1.18%	0	0.00%
Non-Hispanic Black	463,002	30,755	6.64%	60,382	13.04%	96,766	20.90%	275,099	59.42%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	454,954	60,571	13.31%	57,716	12.69%	144,914	31.85%	182,429	40.10%	9,325	2.05%	0	0.00%
9-11th Grade	535,298	69,479	12.98%	105,784	19.76%	143,374	26.78%	216,661	40.47%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,055,238	249,053	8.15%	392,088	12.83%	734,218	24.03%	1,655,961	54.20%	23,917	0.78%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,090,064	379,103	9.27%	573,732	14.03%	1,048,936	25.65%	2,055,051	50.24%	33,242	0.81%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Married	3,135,187	340,636	10.86%	475,312	15.16%	842,808	26.88%	1,443,189	46.03%	33,242	1.06%	0	0.00%
Widowed	*	*	*	*	*	*	*	*	*	*	*	*	*
Divorced	173,201	0	0.00%	25,365	14.64%	29,104	16.80%	118,731	68.55%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	*	*	*	*	*	*	*	*	*	*	*	*	*
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	528,158	57,776	10.94%	72,808	13.79%	73,480	13.91%	324,094	61.36%	0	0.00%	0	0.00%
Poverty income ratio less than 1	602,614	11,399	1.89%	65,472	10.86%	229,180	38.03%	296,565	49.21%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	2,959,291	309,928	10.47%	435,453	14.71%	746,276	25.22%	1,434,392	48.47%	33,242	1.12%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	896,037	61,842	6.90%	220,156	24.57%	232,120	25.91%	381,919	42.62%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,505,040	85,845	5.70%	128,294	8.52%	473,772	31.48%	807,803	53.67%	9,325	0.62%	0	0.00%
Obese: 30 or higher	1,638,954	221,932	13.54%	225,282	13.75%	316,614	19.32%	851,209	51.94%	23,917	1.46%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	1,957,517	59,700	3.05%	295,395	15.09%	461,717	23.59%	1,116,788	57.05%	23,917	1.22%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,410,909	209,613	14.86%	141,079	10.00%	496,021	35.16%	564,195	39.99%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	628,489	89,900	14.30%	119,114	18.95%	84,308	13.41%	335,166	53.33%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	573,852	65,461	11.41%	129,101	22.50%	45,155	7.87%	334,135	58.23%	0	0.00%	0	0.00%
No	3,516,212	313,642	8.92%	444,631	12.65%	1,003,781	28.55%	1,720,916	48.94%	33,242	0.95%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,090,064	379,103	9.27%	573,732	14.03%	1,048,936	25.65%	2,055,051	50.24%	33,242	0.81%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,827,737	279,655	9.89%	343,108	12.13%	719,344	25.44%	1,476,304	52.21%	9,325	0.33%	0	0.00%
No	1,262,328	99,448	7.88%	230,624	18.27%	329,592	26.11%	578,747	45.85%	23,917	1.89%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,004,288	308,319	15.38%	333,872	16.66%	357,850	17.85%	1,004,248	50.10%	0	0.00%	0	0.00%
No	2,085,776	70,784	3.39%	239,860	11.50%	691,086	33.13%	1,050,803	50.38%	33,242	1.59%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,009,901	199,113	19.72%	17,525	1.74%	191,728	18.98%	577,619	57.20%	23,917	2.37%	0	0.00%
Mobility limitations	898,681	88,813	9.88%	234,993	26.15%	167,534	18.64%	407,341	45.33%	0	0.00%	0	0.00%
No mobility limitations	826,673	11,399	1.38%	84,744	10.25%	342,725	41.46%	378,480	45.78%	9,325	1.13%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,354,808	79,778	5.89%	236,470	17.45%	346,949	25.61%	691,611	51.05%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	707,764	56,644	8.00%	84,863	11.99%	125,720	17.76%	440,538	62.24%	0	0.00%	0	0.00%
No	3,352,464	322,459	9.62%	481,648	14.37%	923,216	27.54%	1,591,899	47.48%	33,242	0.99%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	617,089	30,811	4.99%	65,630	10.64%	134,601	21.81%	386,047	62.56%	0	0.00%	0	0.00%
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	515,090	160,217	31.10%	51,795	10.06%	63,339	12.30%	239,739	46.54%	0	0.00%	0	0.00%
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	613,448	0	0.00%	230,299	37.54%	95,958	15.64%	263,275	42.92%	23,917	3.90%	0	0.00%
Never	1,765,277	158,701	8.99%	198,844	11.26%	493,864	27.98%	904,542	51.24%	9,325	0.53%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,090,064	379,103	9.27%	573,732	14.03%	1,048,936	25.65%	2,055,051	50.24%	33,242	0.81%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	3,740,622	349,729	9.35%	555,588	14.85%	849,700	22.72%	1,952,362	52.19%	33,242	0.89%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,711,449	531,048	14.31%	386,748	10.42%	557,304	15.02%	2,236,350	60.26%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	494,604	38,917	7.87%)	35,267	7.13%)	19,567	3.96%	400,853	81.05%	0	0.00%	0	0.00%
50-59	986,089	269,421	27.32%)	60,005	6.09%)	116,242	11.79%	540,422	54.80%	0	0.00%	0	0.00%
60-64	195,866	54,785	27.97%)	24,660	12.59%)	72,542	37.04%	43,879	22.40%	0	0.00%	0	0.00%
65-69	512,329	28,488	5.56%)	133,558	26.07%)	68,312	13.33%	281,971	55.04%	0	0.00%	0	0.00%
70-74	338,057	32,931	9.74%)	29,609	8.76%)	71,768	21.23%	203,749	60.27%	0	0.00%	0	0.00%
75-79	258,006	61,665	23.90%)	10,144	3.93%)	15,134	5.87%	171,064	66.30%	0	0.00%	0	0.00%
80+	434,470	44,842	10.32%)	75,125	17.29%)	169,375	38.98%	145,129	33.40%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	*	*	*	*	*	*	*	*	*	*	*	*	*
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	2,778,872	462,288	16.64%)	271,791	9.78%)	292,704	10.53%	1,752,089	63.05%	0	0.00%	0	0.00%
Non-Hispanic Black	359,280	15,549	4.33%)	99,385	27.66%)	84,747	23.59%	159,599	44.42%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	288,264	49,450	17.15%)	32,683	11.34%)	54,905	19.05%	151,226	52.46%	0	0.00%	0	0.00%
9-11th Grade	330,306	0	0.00%)	46,956	14.22%)	154,614	46.81%	128,736	38.97%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,022,407	481,598	15.93%)	271,842	8.99%)	347,784	11.51%	1,921,183	63.56%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,711,449	531,048	14.31%	386,748	10.42%	557,304	15.02%	2,236,350	60.26%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,535,087	433,045	17.08%)	233,343	9.20%)	409,094	16.14%	1,459,605	57.58%	0	0.00%	0	0.00%
Widowed	*	*	*	*	*	*	*	*	*	*	*	*	*
Divorced	415,297	0	0.00%)	58,622	14.12%)	12,676	3.05%	343,998	82.83%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	350,607	41,348	11.79%)	39,753	11.34%)	47,833	13.64%	221,674	63.23%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	368,611	21,723	5.89%)	54,577	14.81%)	28,146	7.64%	264,165	71.66%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	2,987,123	509,324	17.05%)	269,173	9.01%)	474,950	15.90%	1,733,676	58.04%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,260,091	96,877	7.69%)	192,856	15.30%)	146,713	11.64%	823,646	65.36%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,075,739	264,528	24.59%)	104,688	9.73%)	158,585	14.74%	547,938	50.94%	0	0.00%	0	0.00%
Obese: 30 or higher	1,317,404	157,025	11.92%)	89,204	6.77%)	206,409	15.67%	864,766	65.64%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	1,903,851	186,003	9.77%)	226,359	11.89%)	261,686	13.75%	1,229,803	64.60%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,274,404	316,066	24.80%)	86,132	6.76%)	209,918	16.47%	662,288	51.97%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	438,640	28,978	6.61%)	44,647	10.18%)	54,269	12.37%	310,746	70.84%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,711,449	531,048	14.31%	386,748	10.42%	557,304	15.02%	2,236,350	60.26%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	554,195	6,764	1.22%)	74,256	13.40%)	123,142	22.22%	350,032	63.16%	0	0.00%	0	0.00%
No	3,157,254	524,283	16.61%)	312,491	9.90%)	434,162	13.75%	1,886,318	59.75%	0	0.00%	0	0.00%
Smoked 100 cig in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,220,224	242,446	10.92%)	144,572	6.51%)	329,190	14.83%	1,504,016	67.74%	0	0.00%	0	0.00%
No	1,491,225	288,602	19.35%)	242,176	16.24%)	228,114	15.30%	732,333	49.11%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,523,871	201,395	13.22%)	205,143	13.46%)	260,804	17.11%	856,529	56.21%	0	0.00%	0	0.00%
No	2,187,578	329,653	15.07%)	181,605	8.30%)	296,499	13.55%	1,379,821	63.08%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	645,338	40,443	6.27%)	77,753	12.05%)	158,851	24.62%	368,291	57.07%	0	0.00%	0	0.00%
Mobility limitations	560,493	76,372	13.63%)	90,814	16.20%)	118,620	21.16%	274,687	49.01%	0	0.00%	0	0.00%
No mobility limitations	1,163,829	172,979	14.86%)	218,181	18.75%)	146,251	12.57%	626,419	53.82%	0	0.00%	0	0.00%
N/A - Respondent not asked mob. lim. questions	1,341,788	241,253	17.98%)	0	0.00%)	133,582	9.96%	966,953	72.06%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	416,679	181,324	43.52%)	54,860	13.17%)	12,579	3.02%	167,917	40.30%	0	0.00%	0	0.00%
No	3,294,770	349,724	10.61%)	331,888	10.07%)	544,724	16.53%	2,068,433	62.78%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	488,526	59,900	12.26%)	61,997	12.69%)	22,696	4.65%	343,933	70.40%	0	0.00%	0	0.00%
Once a day	411,361	164,194	39.91%)	12,081	2.94%)	46,159	11.22%	188,928	45.93%	0	0.00%	0	0.00%
2 or more time a week	273,003	31,431	11.51%)	50,003	18.32%)	76,410	27.99%	115,158	42.18%	0	0.00%	0	0.00%
Once a week	360,744	154,991	42.96%)	81,550	22.61%)	0	0.00%	124,203	34.43%	0	0.00%	0	0.00%
1-3 times a month	625,817	0	0.00%)	79,193	12.65%)	108,647	17.36%	437,977	69.98%	0	0.00%	0	0.00%
Never	1,503,611	87,716	5.83%)	86,353	5.74%)	303,391	20.18%	1,026,151	68.25%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M02a - Frequency of bladder control problems among those who responded 'yes' to stress urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,711,449	531,048	14.31%	386,748	10.42%	557,304	15.02%	2,236,350	60.26%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	188,067	0	0.00%)	28,523	15.17%)	82,826	44.04%	76,717	40.79%	0	0.00%	0	0.00%
Never	3,444,825	498,232	14.46%)	322,258	9.35%)	474,477	13.77%	2,149,858	62.41%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.043: How frequently does this (referring to KIQ.042) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?