

Table M03 - Prevalence of urge urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,412,139	7,948,694	8.79%	7.24%	10.34%	82,463,445	91.21%
Age at screening	-	-	-	-	-	-	-
10-19	318,425	0	0.00%	0.00%	0.00%	318,425	100.00%
20-29	11,767,942	278,488	2.37%	0.65%	4.08%	11,489,454	97.63%
30-39	18,212,868	1,008,989	5.54%	2.75%	8.33%	17,203,879	94.46%
40-49	22,567,759	922,372	4.09%	2.38%	5.79%	21,645,387	95.91%
50-59	18,568,743	1,480,063	7.97%	4.91%	11.03%	17,088,680	92.03%
60-64	4,648,416	653,662	14.06%	6.47%	21.65%	3,994,755	85.94%
65-69	3,936,729	858,062	21.80%	15.03%	28.56%	3,078,668	78.20%
70-74	4,053,453	1,060,126	26.15%	15.00%	37.31%	2,993,327	73.85%
75-79	2,934,099	645,737	22.01%	12.87%	31.15%	2,288,361	77.99%
80-84	2,102,740	546,404	25.99%	19.64%	32.33%	1,556,336	74.01%
85+	1,300,965	494,792	38.03%	21.00%	55.07%	806,173	61.97%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,055,890	401,432	5.69%	3.16%	8.22%	6,654,458	94.31%
Other Hispanic	4,966,199	408,927	8.23%	2.25%	14.22%	4,557,272	91.77%
Non-Hispanic White	66,243,971	5,942,960	8.97%	6.93%	11.01%	60,301,012	91.03%
Non-Hispanic Black	9,016,771	1,132,358	12.56%	7.36%	17.75%	7,884,413	87.44%
Other Race - Including Multi-Racial	3,129,308	63,018	2.01%	(2.30%)	6.33%	3,066,291	97.99%
Education	-	-	-	-	-	-	-
Missing	933,223	54,931	5.89%	(2.51%)	14.28%	878,292	94.11%
Less Than 9th Grade	6,516,375	1,126,857	17.29%	8.53%	26.06%	5,389,518	82.71%
9-11th Grade	10,170,827	1,026,877	10.10%	5.55%	14.64%	9,143,950	89.90%
High School Grad/GED or Equivalent	72,644,884	5,729,516	7.89%	6.34%	9.44%	66,915,368	92.11%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	909,023	33,084	3.64%	(2.44%)	9.72%	875,939	96.36%
Married	59,642,677	5,112,045	8.57%	6.97%	10.17%	54,530,632	91.43%
Widowed	4,052,855	800,360	19.75%	11.24%	28.26%	3,252,495	80.25%
Divorced	7,804,451	935,753	11.99%	5.62%	18.36%	6,868,698	88.01%
Separated	2,163,795	112,297	5.19%	0.06%	10.32%	2,051,498	94.81%
Never Married	10,212,194	860,199	8.42%	4.01%	12.84%	9,351,995	91.58%
Living with Partner	5,531,980	94,956	1.72%	(0.28%)	3.71%	5,437,024	98.28%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,980,213	472,926	9.50%	3.78%	15.21%	4,507,287	90.50%
Poverty income ratio less than 1	9,683,534	898,930	9.28%	5.69%	12.88%	8,784,604	90.72%
Poverty income ratio 1 or greater	75,748,392	6,576,838	8.68%	7.08%	10.29%	69,171,554	91.32%
Body mass index	-	-	-	-	-	-	-
Missing	3,214,827	804,935	25.04%	13.33%	36.75%	2,409,892	74.96%
Normal: less than 25	27,340,807	2,007,883	7.34%	4.44%	10.25%	25,332,924	92.66%
Overweight: 25.0 - 29.9	36,239,254	2,673,520	7.38%	5.59%	9.17%	33,565,734	92.62%
Obese: 30 or higher	23,617,252	2,462,356	10.43%	8.33%	12.52%	21,154,895	89.57%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,165,821	348,421	11.01%	4.25%	17.77%	2,817,400	88.99%
Glycohemoglobin < 5.7%	69,910,044	4,604,975	6.59%	5.31%	7.86%	65,305,069	93.41%
Glycohemoglobin 5.7% - 6.4%	11,173,113	1,617,092	14.47%	9.15%	19.80%	9,556,021	85.53%
Glycohemoglobin 6.5% or higher	6,163,161	1,378,206	22.36%	14.54%	30.18%	4,784,955	77.64%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,412,139	7,948,694	8.79%	7.24%	10.34%	82,463,445	91.21%
Diabetes	-	-	-	-	-	-	-
Yes	6,628,716	1,482,276	22.36%	15.48%	29.24%	5,146,440	77.64%
No	83,783,423	6,466,419	7.72%	6.50%	8.93%	77,317,005	92.28%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	51,264,828	5,342,556	10.42%	8.29%	12.55%	45,922,272	89.58%
No	39,060,150	2,606,138	6.67%	5.26%	8.08%	36,454,012	93.33%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	1,046,623	10,036	0.96%	(1.18%)	3.09%	1,036,586	99.04%
Yes	21,514,162	3,050,605	14.18%	10.41%	17.95%	18,463,557	85.82%
No	67,754,615	4,888,054	7.21%	5.88%	8.55%	62,866,562	92.79%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	5,360,610	1,073,254	20.02%	14.14%	25.90%	4,287,356	79.98%
Mobility limitations	6,883,533	1,950,957	28.34%	20.44%	36.25%	4,932,575	71.66%
No mobility limitations	18,453,406	2,592,143	14.05%	9.99%	18.10%	15,861,264	85.95%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	59,699,976	2,317,726	3.88%	2.61%	5.16%	57,382,250	96.12%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	90,412,139	7,948,694	8.79%	7.24%	10.34%	82,463,445	91.21%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	90,412,139	7,948,694	8.79%	7.24%	10.34%	82,463,445	91.21%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	90,412,139	7,948,694	8.79%	7.24%	10.34%	82,463,445	91.21%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	91,101,576	10,604,628	11.64%	9.65%	13.63%	80,496,947	88.36%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	538,199	0	0.00%	0.00%	0.00%	538,199	100.00%
20-29	12,110,825	537,441	4.44%	1.19%	7.68%	11,573,384	95.56%
30-39	17,822,488	1,420,809	7.97%	4.23%	11.72%	16,401,679	92.03%
40-49	20,941,149	1,653,168	7.89%	4.81%	10.98%	19,287,981	92.11%
50-59	17,868,567	1,878,583	10.51%	6.12%	14.91%	15,989,983	89.49%
60-64	5,666,205	834,614	14.73%	7.45%	22.01%	4,831,591	85.27%
65-69	5,762,216	1,184,079	20.55%	13.29%	27.81%	4,578,136	79.45%
70-74	3,988,818	1,168,904	29.30%	22.46%	36.15%	2,819,914	70.70%
75-79	3,030,438	974,996	32.17%	20.69%	43.65%	2,055,442	67.83%
80-84	2,043,274	551,596	27.00%	15.35%	38.64%	1,491,678	73.00%
85+	1,329,398	400,438	30.12%	16.78%	43.46%	928,960	69.88%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,316,409	825,063	11.28%	8.20%	14.35%	6,491,345	88.72%
Other Hispanic	2,977,019	472,305	15.87%	2.62%	29.11%	2,504,713	84.13%
Non-Hispanic White	66,914,787	7,231,566	10.81%	9.07%	12.54%	59,683,221	89.19%
Non-Hispanic Black	9,127,400	1,793,215	19.65%	14.82%	24.48%	7,334,184	80.35%
Other Race - Including Multi-Racial	4,765,962	282,478	5.93%	(0.25%)	12.10%	4,483,484	94.07%
Education	-	-	-	-	-	-	-
Missing	1,796,573	152,405	8.48%	2.67%	14.30%	1,644,168	91.52%
Less Than 9th Grade	6,556,875	1,246,338	19.01%	12.44%	25.58%	5,310,536	80.99%
9-11th Grade	9,028,089	799,754	8.86%	4.31%	13.41%	8,228,334	91.14%
High School Grad/GED or Equivalent	73,372,442	8,368,576	11.41%	8.94%	13.87%	65,003,865	88.59%
Refused	*	*	*	*	*	*	*
Don't Know	335,074	37,555	11.21%	(3.89%)	26.30%	297,519	88.79%
Marital status	-	-	-	-	-	-	-
Missing	1,019,622	201,216	19.73%	(1.67%)	41.14%	818,405	80.27%
Married	59,764,313	6,613,288	11.07%	8.80%	13.34%	53,151,025	88.93%
Widowed	3,595,195	794,181	22.09%	11.90%	32.28%	2,801,014	77.91%
Divorced	7,657,708	871,936	11.39%	7.83%	14.95%	6,785,772	88.61%
Separated	1,505,477	316,235	21.01%	7.95%	34.06%	1,189,241	78.99%
Never Married	11,206,160	1,285,902	11.47%	5.52%	17.43%	9,920,258	88.53%
Living with Partner	5,961,389	521,869	8.75%	2.76%	14.75%	5,439,519	91.25%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,457,226	484,090	10.86%	6.34%	15.38%	3,973,136	89.14%
Poverty income ratio less than 1	9,324,225	887,662	9.52%	3.73%	15.31%	8,436,563	90.48%
Poverty income ratio 1 or greater	77,320,124	9,232,876	11.94%	9.62%	14.26%	68,087,248	88.06%
Body mass index	-	-	-	-	-	-	-
Missing	960,921	208,102	21.66%	4.58%	38.74%	752,819	78.34%
Normal: less than 25	27,564,270	2,610,499	9.47%	7.05%	11.89%	24,953,771	90.53%
Overweight: 25.0 - 29.9	34,664,541	3,291,627	9.50%	6.96%	12.03%	31,372,914	90.50%
Obese: 30 or higher	27,911,843	4,494,400	16.10%	12.28%	19.93%	23,417,443	83.90%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	91,101,576	10,604,628	11.64%	9.65%	13.63%	80,496,947	88.36%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,412,745	331,531	13.74%	7.10%	20.38%	2,081,214	86.26%
Glycohemoglobin < 5.7%	69,788,177	6,723,250	9.63%	7.64%	11.63%	63,064,927	90.37%
Glycohemoglobin 5.7% - 6.4%	12,800,669	2,145,989	16.76%	12.87%	20.66%	10,654,681	83.24%
Glycohemoglobin 6.5% or higher	6,099,984	1,403,858	23.01%	16.53%	29.50%	4,696,126	76.99%
Diabetes	-	-	-	-	-	-	-
Yes	6,985,031	1,871,723	26.80%	20.49%	33.11%	5,113,308	73.20%
No	84,116,545	8,732,906	10.38%	8.40%	12.37%	75,383,639	89.62%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,706,958	6,850,536	13.00%	11.04%	14.95%	45,856,422	87.00%
No	38,356,405	3,742,261	9.76%	7.10%	12.42%	34,614,143	90.24%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	789,967	34,108	4.32%	(3.97%)	12.61%	755,859	95.68%
Yes	27,626,693	4,853,262	17.57%	14.05%	21.09%	22,773,431	82.43%
No	62,579,243	5,667,710	9.06%	7.00%	11.11%	56,911,533	90.94%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	9,312,080	2,466,516	26.49%	21.37%	31.61%	6,845,564	73.51%
Mobility limitations	7,921,738	1,520,395	19.19%	15.14%	23.25%	6,401,342	80.81%
No mobility limitations	19,596,995	2,676,649	13.66%	9.80%	17.52%	16,920,346	86.34%
N/A - Respondent not asked mobility limitation questions	54,270,763	3,941,068	7.26%	4.94%	9.59%	50,329,695	92.74%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	91,101,576	10,604,628	11.64%	9.65%	13.63%	80,496,947	88.36%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	91,101,576	10,604,628	11.64%	9.65%	13.63%	80,496,947	88.36%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	91,101,576	10,604,628	11.64%	9.65%	13.63%	80,496,947	88.36%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,713,557	10,687,174	11.40%	9.28%	13.53%	83,026,383	88.60%
Age at screening	-	-	-	-	-	-	-
10-19	384,918	22,647	5.88%	(5.78%)	17.54%	362,271	94.12%
20-29	14,540,015	516,711	3.55%	0.98%	6.13%	14,023,304	96.45%
30-39	17,503,545	538,575	3.08%	0.96%	5.20%	16,964,970	96.92%
40-49	21,809,592	2,264,358	10.38%	5.84%	14.93%	19,545,234	89.62%
50-59	18,142,193	3,056,180	16.85%	11.01%	22.68%	15,086,013	83.15%
60-64	6,373,461	805,814	12.64%	6.12%	19.16%	5,567,647	87.36%
65-69	5,297,573	1,194,963	22.56%	16.29%	28.82%	4,102,610	77.44%
70-74	4,006,172	870,715	21.73%	14.91%	28.56%	3,135,457	78.27%
75-79	2,536,289	627,670	24.75%	14.98%	34.51%	1,908,619	75.25%
80-84	2,073,206	534,126	25.76%	11.02%	40.50%	1,539,080	74.24%
85+	1,046,592	255,415	24.40%	14.45%	34.35%	791,177	75.60%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,979,822	900,870	11.29%	8.20%	14.38%	7,078,952	88.71%
Other Hispanic	2,862,199	179,655	6.28%	(0.53%)	13.08%	2,682,543	93.72%
Non-Hispanic White	69,156,150	7,936,741	11.48%	9.09%	13.86%	61,219,409	88.52%
Non-Hispanic Black	9,553,122	1,308,202	13.69%	10.60%	16.79%	8,244,920	86.31%
Other Race - Including Multi-Racial	4,162,265	361,706	8.69%	0.41%	16.97%	3,800,559	91.31%
Education	-	-	-	-	-	-	-
Missing	1,125,917	0	0.00%	0.00%	0.00%	1,125,917	100.00%
Less Than 9th Grade	6,403,935	911,096	14.23%	9.88%	18.58%	5,492,839	85.77%
9-11th Grade	10,260,776	1,345,677	13.11%	8.67%	17.56%	8,915,100	86.89%
High School Grad/GED or Equivalent	75,757,393	8,383,251	11.07%	8.88%	13.25%	67,374,142	88.93%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,042,842	104,477	10.02%	(6.91%)	26.94%	938,365	89.98%
Married	60,157,438	7,034,216	11.69%	9.27%	14.12%	53,123,221	88.31%
Widowed	2,985,428	413,290	13.84%	5.24%	22.45%	2,572,137	86.16%
Divorced	9,188,198	1,469,022	15.99%	10.02%	21.96%	7,719,176	84.01%
Separated	2,712,975	450,125	16.59%	5.14%	28.04%	2,262,849	83.41%
Never Married	9,472,270	874,712	9.23%	4.40%	14.07%	8,597,558	90.77%
Living with Partner	8,115,581	302,506	3.73%	0.66%	6.80%	7,813,076	96.27%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	2,977,906	549,522	18.45%	9.02%	27.89%	2,428,383	81.55%
Poverty income ratio less than 1	9,079,370	1,322,067	14.56%	9.52%	19.60%	7,757,303	85.44%
Poverty income ratio 1 or greater	81,656,281	8,815,585	10.80%	8.71%	12.88%	72,840,696	89.20%
Body mass index	-	-	-	-	-	-	-
Missing	1,009,735	60,390	5.98%	(0.40%)	12.37%	949,345	94.02%
Normal: less than 25	25,820,546	2,404,747	9.31%	6.33%	12.30%	23,415,798	90.69%
Overweight: 25.0 - 29.9	35,984,960	4,139,323	11.50%	8.68%	14.32%	31,845,637	88.50%
Obese: 30 or higher	30,898,317	4,082,714	13.21%	9.01%	17.42%	26,815,603	86.79%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,564,155	229,758	6.45%	0.87%	12.02%	3,334,397	93.55%
Glycohemoglobin < 5.7%	70,325,447	6,824,439	9.70%	7.67%	11.74%	63,501,008	90.30%
Glycohemoglobin 5.7% - 6.4%	14,222,346	2,372,343	16.68%	12.57%	20.79%	11,850,003	83.32%
Glycohemoglobin 6.5% or higher	5,601,610	1,260,634	22.50%	13.47%	31.54%	4,340,976	77.50%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,713,557	10,687,174	11.40%	9.28%	13.53%	83,026,383	88.60%
Diabetes	-	-	-	-	-	-	-
Yes	6,979,396	1,332,328	19.09%	14.14%	24.04%	5,647,067	80.91%
No	86,734,162	9,354,846	10.79%	8.71%	12.86%	77,379,315	89.21%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	53,038,544	7,035,598	13.27%	10.93%	15.60%	46,002,946	86.73%
No	40,675,013	3,651,577	8.98%	5.98%	11.98%	37,023,437	91.02%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	26,882,711	5,314,677	19.77%	14.98%	24.56%	21,568,034	80.23%
No	66,652,383	5,315,919	7.98%	6.04%	9.91%	61,336,464	92.02%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,937,459	1,338,183	19.29%	11.57%	27.01%	5,599,275	80.71%
Mobility limitations	5,930,657	1,753,157	29.56%	19.99%	39.13%	4,177,499	70.44%
No mobility limitations	20,291,603	3,424,288	16.88%	13.16%	20.59%	16,867,316	83.12%
N/A - Respondent not asked mobility limitation questions	60,553,839	4,171,546	6.89%	5.54%	8.23%	56,382,293	93.11%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,936,022	1,600,880	20.17%	14.26%	26.09%	6,335,142	79.83%
No	85,661,432	9,086,294	10.61%	8.51%	12.71%	76,575,138	89.39%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	372,256	52,389	14.07%	(3.44%)	31.59%	319,867	85.93%
2 or more times a day	13,806,596	2,370,770	17.17%	13.53%	20.82%	11,435,826	82.83%
Once a day	8,509,610	790,356	9.29%	4.16%	14.41%	7,719,255	90.71%
2 or more time a week	5,956,076	1,063,506	17.86%	7.67%	28.04%	4,892,569	82.14%
Once a week	5,314,628	657,234	12.37%	5.06%	19.67%	4,657,394	87.63%
1-3 times a month	9,361,331	1,777,533	18.99%	11.39%	26.59%	7,583,799	81.01%
Never	50,100,244	3,975,387	7.93%	5.46%	10.41%	46,124,858	92.07%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	372,256	52,389	14.07%	(3.44%)	31.59%	319,867	85.93%
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	749,457	483,850	64.56%	48.92%	80.20%	265,607	35.44%
Never	92,195,030	10,029,659	10.88%	8.72%	13.04%	82,165,371	89.12%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,932,386	11,272,360	11.87%	9.97%	13.77%	83,660,025	88.13%
Age at screening	-	-	-	-	-	-	-
10-19	581,889	15,708	2.70%	(3.35%)	8.75%	566,180	97.30%
20-29	14,324,402	479,511	3.35%	0.94%	5.75%	13,844,891	96.65%
30-39	17,096,292	945,757	5.53%	3.26%	7.81%	16,150,535	94.47%
40-49	21,351,091	1,992,599	9.33%	5.40%	13.26%	19,358,492	90.67%
50-59	18,864,707	2,400,891	12.73%	9.72%	15.74%	16,463,816	87.27%
60-64	7,469,087	1,268,582	16.98%	11.51%	22.46%	6,200,505	83.02%
65-69	5,293,210	1,336,379	25.25%	17.60%	32.89%	3,956,831	74.75%
70-74	3,767,362	916,570	24.33%	17.13%	31.53%	2,850,792	75.67%
75-79	2,921,173	832,791	28.51%	19.77%	37.24%	2,088,383	71.49%
80+	3,263,173	1,083,571	33.21%	24.56%	41.86%	2,179,601	66.79%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,009,819	1,076,308	11.95%	6.70%	17.19%	7,933,511	88.05%
Other Hispanic	4,540,869	565,954	12.46%	8.50%	16.42%	3,974,916	87.54%
Non-Hispanic White	65,896,437	7,368,578	11.18%	8.94%	13.42%	58,527,858	88.82%
Non-Hispanic Black	9,832,078	1,759,216	17.89%	14.50%	21.29%	8,072,862	82.11%
Other Race - Including Multi-Racial	5,653,183	502,304	8.89%	0.21%	17.56%	5,150,879	91.11%
Education	-	-	-	-	-	-	-
Missing	1,275,540	172,283	13.51%	(1.50%)	28.51%	1,103,257	86.49%
Less Than 9th Grade	6,695,497	1,544,257	23.06%	13.40%	32.73%	5,151,240	76.94%
9-11th Grade	11,881,477	1,924,983	16.20%	10.38%	22.02%	9,956,494	83.80%
High School Grad/GED or Equivalent	74,943,444	7,549,271	10.07%	7.71%	12.44%	67,394,173	89.93%
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,282,066	49,279	3.84%	(0.12%)	7.81%	1,232,787	96.16%
Married	60,655,911	7,134,742	11.76%	9.54%	13.99%	53,521,169	88.24%
Widowed	3,208,196	739,099	23.04%	15.43%	30.65%	2,469,096	76.96%
Divorced	8,428,145	1,624,983	19.28%	11.00%	27.56%	6,803,162	80.72%
Separated	2,134,533	353,160	16.55%	7.15%	25.94%	1,781,373	83.45%
Never Married	12,388,270	692,371	5.59%	3.42%	7.76%	11,695,899	94.41%
Living with Partner	6,649,894	678,726	10.21%	5.36%	15.05%	5,971,168	89.79%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,832,632	1,157,599	16.94%	11.53%	22.35%	5,675,033	83.06%
Poverty income ratio less than 1	11,430,626	1,277,055	11.17%	6.34%	16.01%	10,153,571	88.83%
Poverty income ratio 1 or greater	76,669,128	8,837,707	11.53%	9.55%	13.50%	67,831,421	88.47%
Body mass index	-	-	-	-	-	-	-
Missing	808,899	197,574	24.43%	8.62%	40.23%	611,326	75.58%
Normal: less than 25	26,748,202	2,464,262	9.21%	7.27%	11.16%	24,283,940	90.79%
Overweight: 25.0 - 29.9	36,919,637	3,755,813	10.17%	6.91%	13.43%	33,163,825	89.83%
Obese: 30 or higher	30,455,646	4,854,712	15.94%	12.42%	19.46%	25,600,934	84.06%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,550,279	164,562	4.64%	1.91%	7.36%	3,385,717	95.36%
Glycohemoglobin < 5.7%	62,899,045	5,700,739	9.06%	6.84%	11.29%	57,198,306	90.94%
Glycohemoglobin 5.7% - 6.4%	21,210,133	3,774,940	17.80%	13.99%	21.61%	17,435,193	82.20%
Glycohemoglobin 6.5% or higher	7,272,929	1,632,120	22.44%	13.93%	30.95%	5,640,809	77.56%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,932,386	11,272,360	11.87%	9.97%	13.77%	83,660,025	88.13%
Diabetes	-	-	-	-	-	-	-
Yes	8,457,156	2,511,112	29.69%	21.19%	38.19%	5,946,044	70.31%
No	86,475,230	8,761,248	10.13%	8.28%	11.99%	77,713,981	89.87%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,071,877	7,327,311	14.07%	11.23%	16.91%	44,744,566	85.93%
No	42,792,655	3,907,220	9.13%	6.61%	11.65%	38,885,434	90.87%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,230,043	5,194,367	18.40%	14.95%	21.85%	23,035,677	81.60%
No	66,599,730	6,077,994	9.13%	7.42%	10.83%	60,521,736	90.87%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	8,986,169	2,293,992	25.53%	18.72%	32.34%	6,692,176	74.47%
Mobility limitations	6,860,598	1,878,864	27.39%	19.21%	35.56%	4,981,733	72.61%
No mobility limitations	18,940,648	2,913,602	15.38%	12.05%	18.72%	16,027,047	84.62%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	60,107,142	4,148,073	6.90%	4.52%	9.28%	55,959,069	93.10%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,814,272	1,229,491	15.73%	6.95%	24.51%	6,584,781	84.27%
No	86,905,951	10,013,034	11.52%	9.67%	13.37%	76,892,917	88.48%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	610,540	117,400	19.23%	0.95%	37.51%	493,141	80.77%
2 or more times a day	10,308,368	1,378,698	13.37%	7.68%	19.07%	8,929,670	86.63%
Once a day	8,264,954	1,429,554	17.30%	11.67%	22.93%	6,835,400	82.70%
2 or more time a week	5,908,062	1,103,780	18.68%	13.25%	24.11%	4,804,282	81.32%
Once a week	4,371,150	763,501	17.47%	8.60%	26.33%	3,607,649	82.53%
1-3 times a month	8,315,400	1,361,707	16.38%	12.22%	20.53%	6,953,693	83.62%
Never	56,772,742	5,100,445	8.98%	7.16%	10.80%	51,672,297	91.02%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	610,540	117,400	19.23%	0.95%	37.51%	493,141	80.77%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	1,037,565	463,418	44.66%	14.56%	74.77%	574,147	55.34%
Never	92,548,070	10,535,088	11.38%	9.72%	13.04%	82,012,982	88.62%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,585,798	11,866,036	12.41%	10.58%	14.25%	83,719,761	87.59%
Age at screening	-	-	-	-	-	-	-
10-19	297,636	41,692	14.01%	(10.98%)	38.99%	255,943	85.99%
20-29	12,944,492	533,407	4.12%	1.90%	6.34%	12,411,085	95.88%
30-39	16,576,959	1,171,423	7.07%	3.26%	10.87%	15,405,536	92.93%
40-49	21,324,506	1,952,867	9.16%	5.52%	12.80%	19,371,639	90.84%
50-59	21,456,557	2,571,971	11.99%	8.70%	15.28%	18,884,585	88.01%
60-64	7,051,206	1,611,652	22.86%	16.69%	29.02%	5,439,554	77.14%
65-69	6,200,612	1,077,976	17.38%	10.53%	24.24%	5,122,636	82.62%
70-74	3,763,064	881,903	23.44%	18.02%	28.85%	2,881,161	76.56%
75-79	2,936,105	898,836	30.61%	18.79%	42.44%	2,037,269	69.39%
80+	3,034,662	1,124,309	37.05%	28.69%	45.41%	1,910,352	62.95%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,079,208	1,095,519	12.07%	8.93%	15.20%	7,983,689	87.93%
Other Hispanic	4,899,222	505,517	10.32%	4.98%	15.66%	4,393,705	89.68%
Non-Hispanic White	66,118,840	7,963,037	12.04%	9.71%	14.38%	58,155,802	87.96%
Non-Hispanic Black	9,655,413	2,016,334	20.88%	13.53%	28.23%	7,639,079	79.12%
Other Race - Including Multi-Racial	5,833,115	285,629	4.90%	0.56%	9.23%	5,547,486	95.10%
Education	-	-	-	-	-	-	-
Missing	1,359,443	300,834	22.13%	7.75%	36.51%	1,058,609	77.87%
Less Than 9th Grade	6,477,166	1,128,418	17.42%	12.80%	22.04%	5,348,748	82.58%
9-11th Grade	10,241,048	1,543,160	15.07%	9.63%	20.51%	8,697,888	84.93%
High School Grad/GED or Equivalent	77,361,680	8,893,625	11.50%	9.45%	13.54%	68,468,055	88.50%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	313,184	57,241	18.28%	(8.62%)	45.17%	255,943	81.72%
Married	61,053,634	7,967,730	13.05%	10.57%	15.53%	53,085,904	86.95%
Widowed	3,241,681	394,066	12.16%	3.33%	20.98%	2,847,616	87.84%
Divorced	8,826,206	1,191,296	13.50%	8.46%	18.54%	7,634,910	86.50%
Separated	2,184,790	210,011	9.61%	1.77%	17.46%	1,974,780	90.39%
Never Married	12,431,390	1,253,248	10.08%	5.54%	14.62%	11,178,141	89.92%
Living with Partner	7,478,020	747,208	9.99%	5.90%	14.09%	6,730,812	90.01%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,719,530	1,150,775	17.13%	10.93%	23.32%	5,568,755	82.87%
Poverty income ratio less than 1	10,991,125	1,301,649	11.84%	8.23%	15.45%	9,689,475	88.16%
Poverty income ratio 1 or greater	77,875,143	9,413,612	12.09%	10.18%	13.99%	68,461,531	87.91%
Body mass index	-	-	-	-	-	-	-
Missing	914,576	269,949	29.52%	2.78%	56.25%	644,627	70.48%
Normal: less than 25	24,656,831	2,669,139	10.83%	8.30%	13.35%	21,987,691	89.17%
Overweight: 25.0 - 29.9	36,228,379	3,558,307	9.82%	7.24%	12.41%	32,670,073	90.18%
Obese: 30 or higher	33,786,012	5,368,642	15.89%	13.92%	17.87%	28,417,370	84.11%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,643,943	91,470	3.46%	1.60%	5.32%	2,552,472	96.54%
Glycohemoglobin < 5.7%	61,620,636	6,431,032	10.44%	7.99%	12.89%	55,189,604	89.56%
Glycohemoglobin 5.7% - 6.4%	23,078,335	3,324,840	14.41%	11.04%	17.77%	19,753,495	85.59%
Glycohemoglobin 6.5% or higher	8,242,884	2,018,694	24.49%	20.18%	28.80%	6,224,190	75.51%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?

Table M03 - Prevalence of urge urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,585,798	11,866,036	12.41%	10.58%	14.25%	83,719,761	87.59%
Diabetes	-	-	-	-	-	-	-
Yes	8,780,629	2,236,721	25.47%	18.27%	32.67%	6,543,908	74.53%
No	86,805,168	9,629,315	11.09%	9.06%	13.13%	77,175,853	88.91%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	47,479,384	7,157,497	15.07%	12.88%	17.27%	40,321,887	84.93%
No	48,106,414	4,708,540	9.79%	7.41%	12.17%	43,397,874	90.21%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,857,116	5,361,569	18.58%	15.44%	21.72%	23,495,546	81.42%
No	66,617,365	6,493,583	9.75%	7.97%	11.53%	60,123,782	90.25%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	7,924,746	2,015,691	25.44%	17.87%	33.00%	5,909,055	74.56%
Mobility limitations	6,892,982	1,776,064	25.77%	19.87%	31.66%	5,116,918	74.23%
No mobility limitations	20,034,171	3,469,531	17.32%	13.61%	21.02%	16,564,640	82.68%
N/A - Respondent not asked mobility limitation questions	60,733,899	4,604,751	7.58%	5.15%	10.01%	56,129,148	92.42%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,676,778	1,762,717	22.96%	14.27%	31.65%	5,914,061	77.04%
No	87,705,670	10,096,494	11.51%	9.48%	13.54%	77,609,176	88.49%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	459,339	254,678	55.44%	18.74%	92.15%	204,661	44.56%
2 or more times a day	9,167,832	1,426,665	15.56%	9.70%	21.43%	7,741,168	84.44%
Once a day	8,770,661	830,369	9.47%	5.68%	13.26%	7,940,293	90.53%
2 or more time a week	6,773,718	992,469	14.65%	8.19%	21.11%	5,781,249	85.35%
Once a week	4,927,442	829,676	16.84%	9.28%	24.39%	4,097,766	83.16%
1-3 times a month	10,278,875	2,165,283	21.07%	16.14%	25.99%	8,113,592	78.93%
Never	55,144,254	5,366,897	9.73%	8.07%	11.39%	49,777,356	90.27%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	459,339	254,678	55.44%	18.74%	92.15%	204,661	44.56%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	844,643	392,062	46.42%	31.10%	61.73%	452,582	53.58%
Never	93,745,910	11,040,333	11.78%	9.98%	13.58%	82,705,577	88.22%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.044: During the past 12 months, have you/has specific person leaked or lost control of even a small amount of urine with an urge or pressure to urinate and you/he/she could not get to the toilet fast enough?