

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	1,008,989	17,431	1.73%	149,655	14.83%	301,486	29.88%	540,416	53.56%	0	0.00%	0	0.00%
40-49	922,372	0	0.00%	73,197	7.94%	223,097	24.19%	626,078	67.88%	0	0.00%	0	0.00%
50-59	1,480,063	93,114	6.29%	352,041	23.79%	370,833	25.06%	664,076	44.87%	0	0.00%	0	0.00%
60-64	653,662	144,147	22.05%	84,184	12.88%	110,933	16.97%	314,398	48.10%	0	0.00%	0	0.00%
65-69	858,062	73,522	8.57%	142,309	16.58%	136,626	15.92%	505,605	58.92%	0	0.00%	0	0.00%
70-74	1,060,126	159,728	15.07%	215,898	20.37%	357,958	33.77%	301,259	28.42%	25,283	2.38%	0	0.00%
75-79	645,737	148,021	22.92%	134,584	20.84%	181,620	28.13%	181,512	28.11%	0	0.00%	0	0.00%
80-84	546,404	73,271	13.41%	208,923	38.24%	179,083	32.77%	85,127	15.58%	0	0.00%	0	0.00%
85+	494,792	118,409	23.93%	163,146	32.97%	85,720	17.32%	127,516	25.77%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	401,432	76,035	18.94%	86,605	21.57%	95,167	23.71%	143,625	35.78%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	5,942,960	662,296	11.14%	1,081,830	18.20%	1,548,733	26.06%	2,624,818	44.17%	25,283	0.43%	0	0.00%
Non-Hispanic Black	1,132,358	67,247	5.94%	319,674	28.23%	357,101	31.54%	388,337	34.29%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	1,126,857	280,693	24.91%	213,809	18.97%	218,960	19.43%	413,395	36.69%	0	0.00%	0	0.00%
9-11th Grade	1,026,877	130,631	12.72%	235,725	22.96%	419,354	40.84%	241,168	23.49%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	5,729,516	385,398	6.73%	1,101,713	19.23%	1,380,138	24.09%	2,836,986	49.52%	25,283	0.44%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	5,112,045	655,465	12.82%	941,962	18.43%	1,279,860	25.04%	2,234,758	43.72%	0	0.00%	0	0.00%
Widowed	800,360	100,172	12.52%	183,573	22.94%	272,473	34.04%	244,143	30.50%	0	0.00%	0	0.00%
Divorced	935,753	14,352	1.53%	198,680	21.23%	278,583	29.77%	418,856	44.76%	25,283	2.70%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	860,199	49,628	5.77%	212,662	24.72%	172,268	20.03%	425,641	49.48%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	472,926	6,038	1.28%	35,933	7.60%	198,153	41.90%	232,803	49.23%	0	0.00%	0	0.00%
Poverty income ratio less than 1	898,930	89,612	9.97%	212,026	23.59%	199,332	22.17%	397,961	44.27%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	6,576,838	731,992	11.13%	1,327,299	20.18%	1,631,480	24.81%	2,860,784	43.50%	25,283	0.38%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	804,935	210,478	26.15%	177,768	22.08%	317,493	39.44%	99,196	12.32%	0	0.00%	0	0.00%
Normal: less than 25	2,007,883	194,589	9.69%	451,672	22.49%	479,037	23.86%	857,304	42.70%	25,283	1.26%	0	0.00%
Overweight: 25.0 - 29.9	2,673,520	156,392	5.85%	603,451	22.57%	819,578	30.66%	1,094,099	40.92%	0	0.00%	0	0.00%
Obese: 30 or higher	2,462,356	266,183	10.81%	342,368	13.90%	412,856	16.77%	1,440,950	58.52%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	348,421	41,633	11.95%	34,360	9.86%	182,302	52.32%	90,127	25.87%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	4,604,975	361,328	7.85%	963,431	20.92%	920,242	19.98%	2,334,692	50.70%	25,283	0.55%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,617,092	218,198	13.49%	376,375	23.27%	427,255	26.42%	595,264	36.81%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,378,206	206,483	14.98%	201,092	14.59%	499,166	36.22%	471,464	34.21%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,482,276	231,722	15.63%	321,929	21.72%	544,958	36.76%	383,666	25.88%	0	0.00%	0	0.00%
No	6,466,419	595,920	9.22%	1,253,329	19.38%	1,484,006	22.95%	3,107,882	48.06%	25,283	0.39%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,342,556	505,286	9.46%	1,096,328	20.52%	1,445,068	27.05%	2,270,591	42.50%	25,283	0.47%	0	0.00%
No	2,606,138	322,356	12.37%	478,930	18.38%	583,896	22.40%	1,220,957	46.85%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	3,050,605	564,800	18.51%	666,661	21.85%	867,882	28.45%	951,261	31.18%	0	0.00%	0	0.00%
No	4,888,054	262,842	5.38%	908,597	18.59%	1,151,046	23.55%	2,540,287	51.97%	25,283	0.52%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,073,254	203,303	18.94%	264,824	24.67%	256,374	23.89%	348,753	32.49%	0	0.00%	0	0.00%
Mobility limitations	1,950,957	393,936	20.19%	319,818	16.39%	483,589	24.79%	753,615	38.63%	0	0.00%	0	0.00%
No mobility limitations	2,592,143	203,232	7.84%	563,004	21.72%	620,955	23.96%	1,179,669	45.51%	25,283	0.98%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	2,317,726	12,556	0.54%	427,612	18.45%	668,046	28.82%	1,209,512	52.19%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	7,948,694	827,642	10.41%	1,575,258	19.82%	2,028,964	25.53%	3,491,548	43.93%	25,283	0.32%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	537,441	24,290	4.52%	0	0.00%	100,178	18.64%	412,973	76.84%	0	0.00%	0	0.00%
30-39	1,420,809	57,839	4.07%	272,214	19.16%	238,145	16.76%	852,611	60.01%	0	0.00%	0	0.00%
40-49	1,653,168	12,450	0.75%	136,161	8.24%	262,666	15.89%	1,241,891	75.12%	0	0.00%	0	0.00%
50-59	1,878,583	106,965	5.69%	261,656	13.93%	683,372	36.38%	826,591	44.00%	0	0.00%	0	0.00%
60-64	834,614	23,132	2.77%	191,867	22.99%	255,034	30.56%	364,580	43.68%	0	0.00%	0	0.00%
65-69	1,184,079	144,407	12.20%	89,270	7.54%	330,664	27.93%	619,738	52.34%	0	0.00%	0	0.00%
70-74	1,168,904	134,521	11.51%	222,099	19.00%	427,429	36.57%	384,856	32.92%	0	0.00%	0	0.00%
75-79	974,996	51,992	5.33%	290,042	29.75%	265,340	27.21%	367,621	37.70%	0	0.00%	0	0.00%
80-84	551,596	200,732	36.39%	32,239	5.84%	61,362	11.12%	257,264	46.64%	0	0.00%	0	0.00%
85+	400,438	183,962	45.94%	20,504	5.12%	111,586	27.87%	62,918	15.71%	21,468	5.36%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	825,063	75,626	9.17%	88,928	10.78%	153,877	18.65%	506,632	61.41%	0	0.00%	0	0.00%
Other Hispanic	472,305	104,632	22.15%	0	0.00%	49,890	10.56%	317,783	67.28%	0	0.00%	0	0.00%
Non-Hispanic White	7,231,566	721,676	9.98%	1,069,306	14.79%	1,839,785	25.44%	3,579,331	49.50%	21,468	0.30%	0	0.00%
Non-Hispanic Black	1,793,215	38,356	2.14%	328,043	18.29%	594,596	33.16%	832,220	46.41%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	1,246,338	284,077	22.79%	215,652	17.30%	328,030	26.32%	418,579	33.58%	0	0.00%	0	0.00%
9-11th Grade	799,754	99,982	12.50%	128,806	16.11%	190,022	23.76%	380,945	47.63%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	8,368,576	531,941	6.36%	1,094,444	13.08%	2,166,759	25.89%	4,553,964	54.42%	21,468	0.26%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	6,613,288	676,999	10.24%	736,977	11.14%	1,685,896	25.49%	3,491,948	52.80%	21,468	0.32%	0	0.00%
Widowed	794,181	202,314	25.47%	145,052	18.26%	147,472	18.57%	299,343	37.69%	0	0.00%	0	0.00%
Divorced	871,936	42,853	4.91%	99,184	11.38%	234,113	26.85%	495,786	56.86%	0	0.00%	0	0.00%
Separated	316,235	9,367	2.96%	48,827	15.44%	129,189	40.85%	128,853	40.75%	0	0.00%	0	0.00%
Never Married	1,285,902	8,757	0.68%	238,837	18.57%	393,360	30.59%	644,948	50.16%	0	0.00%	0	0.00%
Living with Partner	521,869	0	0.00%	247,175	47.36%	118,420	22.69%	156,274	29.95%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	484,090	124,761	25.77%	98,459	20.34%	179,368	37.05%	60,034	12.40%	21,468	4.43%	0	0.00%
Poverty income ratio less than 1	887,662	62,082	6.99%	91,275	10.28%	252,134	28.40%	482,171	54.32%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	9,232,876	753,447	8.16%	1,326,317	14.37%	2,304,274	24.96%	4,848,838	52.52%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	2,610,499	238,498	9.14%	595,561	22.81%	644,692	24.70%	1,131,749	43.35%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	3,291,627	247,761	7.53%	393,512	11.95%	889,939	27.04%	1,760,415	53.48%	0	0.00%	0	0.00%
Obese: 30 or higher	4,494,400	325,791	7.25%	468,584	10.43%	1,179,677	26.25%	2,498,880	55.60%	21,468	0.48%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	331,531	0	0.00%	53,454	16.12%	157,840	47.61%	120,237	36.27%	0	0.00%	0	0.00%
Glycohemoglobin < 5.7%	6,723,250	468,726	6.97%	972,386	14.46%	1,622,301	24.13%	3,659,836	54.44%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	2,145,989	316,597	14.75%	251,150	11.70%	621,748	28.97%	935,026	43.57%	21,468	1.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,403,858	154,968	11.04%	239,060	17.03%	333,887	23.78%	675,944	48.15%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,871,723	320,691	17.13%	301,602	16.11%	420,002	22.44%	829,428	44.31%	0	0.00%	0	0.00%
No	8,732,906	619,599	7.09%	1,214,449	13.91%	2,315,774	26.52%	4,561,616	52.23%	21,468	0.25%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%
Smoked at least 100 cigarettes in life	-	-		-		-		-		-		-	
Yes	6,850,536	609,713	8.90%	1,143,795	16.70%	1,741,464	25.42%	3,355,564	48.98%	0	0.00%	0	0.00%
No	3,742,261	330,577	8.83%	360,425	9.63%	994,312	26.57%	2,035,479	54.39%	21,468	0.57%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-		-		-		-		-		-	
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Yes	4,853,262	562,943	11.60%	717,344	14.78%	1,437,137	29.61%	2,135,839	44.01%	0	0.00%	0	0.00%
No	5,667,710	356,843	6.30%	798,707	14.09%	1,264,532	22.31%	3,226,160	56.92%	21,468	0.38%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-		-		-		-		-		-	
Missing	2,466,516	489,748	19.86%	396,332	16.07%	606,013	24.57%	952,955	38.64%	21,468	0.87%	0	0.00%
Mobility limitations	1,520,395	182,822	12.02%	258,357	16.99%	600,478	39.49%	478,739	31.49%	0	0.00%	0	0.00%
No mobility limitations	2,676,649	221,653	8.28%	446,513	16.68%	772,868	28.87%	1,235,614	46.16%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	3,941,068	46,067	1.17%	414,849	10.53%	756,418	19.19%	2,723,735	69.11%	0	0.00%	0	0.00%
Sleep disorder	-	-		-		-		-		-		-	
N/A for these years of data	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%
Bowel leakage - gas	-	-		-		-		-		-		-	
N/A for these years of data	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%
Bowel leakage - stool	-	-		-		-		-		-		-	
N/A for these years of data	10,604,628	940,290	8.87%	1,516,051	14.30%	2,735,776	25.80%	5,391,043	50.84%	21,468	0.20%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	516,711	39,090	7.57%	0	0.00%	102,069	19.75%	375,553	72.68%	0	0.00%	0	0.00%
30-39	538,575	0	0.00%	58,075	10.78%	79,129	14.69%	401,372	74.52%	0	0.00%	0	0.00%
40-49	2,264,358	300,348	13.26%	305,259	13.48%	439,916	19.43%	1,218,835	53.83%	0	0.00%	0	0.00%
50-59	3,056,180	143,074	4.68%	329,848	10.79%	459,725	15.04%	2,123,533	69.48%	0	0.00%	0	0.00%
60-64	805,814	39,500	4.90%	0	0.00%	440,192	54.63%	326,122	40.47%	0	0.00%	0	0.00%
65-69	1,194,963	70,999	5.94%	369,121	30.89%	266,427	22.30%	488,415	40.87%	0	0.00%	0	0.00%
70-74	870,715	39,717	4.56%	92,184	10.59%	376,105	43.19%	362,709	41.66%	0	0.00%	0	0.00%
75-79	627,670	40,962	6.53%	64,915	10.34%	248,743	39.63%	273,050	43.50%	0	0.00%	0	0.00%
80-84	534,126	69,042	12.93%	34,457	6.45%	107,514	20.13%	323,113	60.49%	0	0.00%	0	0.00%
85+	255,415	36,311	14.22%	45,300	17.74%	103,558	40.54%	70,246	27.50%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	900,870	35,559	3.95%	57,922	6.43%	325,938	36.18%	481,451	53.44%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	7,936,741	724,557	9.13%	887,093	11.18%	1,859,978	23.44%	4,465,113	56.26%	0	0.00%	0	0.00%
Non-Hispanic Black	1,308,202	18,927	1.45%	179,971	13.76%	272,652	20.84%	836,652	63.95%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Less Than 9th Grade	911,096	50,032	5.49%	107,114	11.76%	311,965	34.24%	441,985	48.51%	0	0.00%	0	0.00%
9-11th Grade	1,345,677	91,775	6.82%	73,789	5.48%	560,085	41.62%	620,027	46.08%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	8,383,251	624,358	7.45%	1,118,256	13.34%	1,751,327	20.89%	4,889,311	58.32%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	7,034,216	415,157	5.90%	952,732	13.54%	1,906,476	27.10%	3,759,851	53.45%	0	0.00%	0	0.00%
Widowed	413,290	11,167	2.70%	154,988	37.50%	129,421	31.31%	117,715	28.48%	0	0.00%	0	0.00%
Divorced	1,469,022	261,947	17.83%	53,823	3.66%	255,192	17.37%	898,059	61.13%	0	0.00%	0	0.00%
Separated	450,125	21,351	4.74%	44,954	9.99%	40,379	8.97%	343,442	76.30%	0	0.00%	0	0.00%
Never Married	874,712	37,842	4.33%	92,662	10.59%	225,382	25.77%	518,825	59.31%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	549,522	0	0.00%	99,765	18.15%	219,950	40.03%	229,808	41.82%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,322,067	3,914	0.30%	201,230	15.22%	312,822	23.66%	804,101	60.82%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	8,815,585	775,129	8.79%	998,164	11.32%	2,090,605	23.71%	4,951,687	56.17%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	2,404,747	188,390	7.83%	322,024	13.39%	461,113	19.18%	1,433,221	59.60%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	4,139,323	275,586	6.66%	699,319	16.89%	1,231,051	29.74%	1,933,367	46.71%	0	0.00%	0	0.00%
Obese: 30 or higher	4,082,714	289,923	7.10%	268,846	6.58%	924,053	22.63%	2,599,892	63.68%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	6,824,439	491,212	7.20%	873,169	12.79%	1,632,090	23.92%	3,827,968	56.09%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	2,372,343	233,164	9.83%	315,252	13.29%	545,185	22.98%	1,278,742	53.90%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	1,260,634	54,667	4.34%	89,371	7.09%	370,004	29.35%	746,591	59.22%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,332,328	186,208	13.98%	84,666	6.35%	438,153	32.89%	623,301	46.78%	0	0.00%	0	0.00%
No	9,354,846	592,834	6.34%	1,214,493	12.98%	2,185,223	23.36%	5,362,295	57.32%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	7,035,598	627,393	8.92%	968,787	13.77%	1,848,106	26.27%	3,591,311	51.04%	0	0.00%	0	0.00%
No	3,651,577	151,649	4.15%	330,372	9.05%	775,271	21.23%	2,394,284	65.57%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,314,677	555,364	10.45%	610,920	11.49%	1,428,160	26.87%	2,720,232	51.18%	0	0.00%	0	0.00%
No	5,315,919	223,679	4.21%	688,239	12.95%	1,138,638	21.42%	3,265,363	61.43%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,338,183	319,091	23.85%	144,185	10.77%	400,240	29.91%	474,667	35.47%	0	0.00%	0	0.00%
Mobility limitations	1,753,157	9,956	0.57%	244,116	13.92%	510,812	29.14%	988,274	56.37%	0	0.00%	0	0.00%
No mobility limitations	3,424,288	181,683	5.31%	588,636	17.19%	864,968	25.26%	1,789,001	52.24%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	4,171,546	268,312	6.43%	322,223	7.72%	847,357	20.31%	2,733,654	65.53%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,600,880	22,614	1.41%	244,662	15.28%	420,300	26.25%	913,304	57.05%	0	0.00%	0	0.00%
No	9,086,294	756,428	8.32%	1,054,498	11.61%	2,203,077	24.25%	5,072,291	55.82%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	2,370,770	350,416	14.78%	232,914	9.82%	599,292	25.28%	1,188,148	50.12%	0	0.00%	0	0.00%
Once a day	790,356	69,420	8.78%	10,782	1.36%	220,561	27.91%	489,592	61.95%	0	0.00%	0	0.00%
2 or more time a week	1,063,506	70,999	6.68%	89,796	8.44%	333,521	31.36%	569,190	53.52%	0	0.00%	0	0.00%
Once a week	657,234	8,970	1.36%	140,381	21.36%	97,060	14.77%	410,822	62.51%	0	0.00%	0	0.00%
1-3 times a month	1,777,533	98,836	5.56%	60,778	3.42%	655,773	36.89%	962,146	54.13%	0	0.00%	0	0.00%
Never	3,975,387	128,012	3.22%	764,509	19.23%	717,168	18.04%	2,365,697	59.51%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	483,850	94,828	19.60%	12,321	2.55%	212,783	43.98%	163,918	33.88%	0	0.00%	0	0.00%
Never	10,029,659	538,028	5.36%	1,286,839	12.83%	2,400,284	23.93%	5,804,508	57.87%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	11,272,360	872,118	7.74%	1,286,804	11.42%	2,980,955	26.44%	6,095,451	54.07%	37,032	0.33%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	479,511	0	0.00%	24,497	5.11%	74,630	15.56%	380,385	79.33%	0	0.00%	0	0.00%
30-39	945,757	11,602	1.23%	141,150	14.92%	63,821	6.75%	729,183	77.10%	0	0.00%	0	0.00%
40-49	1,992,599	161,992	8.13%	12,631	0.63%	549,521	27.58%	1,268,455	63.66%	0	0.00%	0	0.00%
50-59	2,400,891	242,946	10.12%	305,483	12.72%	684,973	28.53%	1,167,488	48.63%	0	0.00%	0	0.00%
60-64	1,268,582	20,832	1.64%	87,489	6.90%	319,739	25.20%	833,653	65.72%	6,869	0.54%	0	0.00%
65-69	1,336,379	114,998	8.61%	188,460	14.10%	395,728	29.61%	637,192	47.68%	0	0.00%	0	0.00%
70-74	916,570	16,073	1.75%	239,114	26.09%	317,286	34.62%	344,098	37.54%	0	0.00%	0	0.00%
75-79	832,791	85,045	10.21%	144,872	17.40%	303,512	36.45%	299,362	35.95%	0	0.00%	0	0.00%
80+	1,083,571	218,630	20.18%	143,107	13.21%	271,745	25.08%	419,926	38.75%	30,163	2.78%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,076,308	39,642	3.68%	84,254	7.83%	258,559	24.02%	693,853	64.47%	0	0.00%	0	0.00%
Other Hispanic	565,954	22,383	3.95%	90,841	16.05%	187,889	33.20%	264,841	46.80%	0	0.00%	0	0.00%
Non-Hispanic White	7,368,578	674,651	9.16%	1,004,064	13.63%	1,664,230	22.59%	3,995,470	54.22%	30,163	0.41%	0	0.00%
Non-Hispanic Black	1,759,216	70,206	3.99%	107,645	6.12%	515,210	29.29%	1,059,287	60.21%	6,869	0.39%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	1,544,257	121,319	7.86%	279,573	18.10%	452,407	29.30%	684,088	44.30%	6,869	0.44%	0	0.00%
9-11th Grade	1,924,983	153,026	7.95%	255,383	13.27%	671,571	34.89%	845,003	43.90%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	7,549,271	459,984	6.09%	751,848	9.96%	1,830,546	24.25%	4,476,730	59.30%	30,163	0.40%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	7,134,742	501,379	7.03%	799,108	11.20%	2,180,007	30.55%	3,654,248	51.22%	0	0.00%	0	0.00%
Widowed	739,099	113,087	15.30%	89,521	12.11%	187,113	25.32%	342,509	46.34%	6,869	0.93%	0	0.00%
Divorced	1,624,983	171,530	10.56%	120,720	7.43%	282,444	17.38%	1,020,125	62.78%	30,163	1.86%	0	0.00%
Separated	353,160	18,360	5.20%	56,591	16.02%	72,595	20.56%	205,613	58.22%	0	0.00%	0	0.00%
Never Married	692,371	59,605	8.61%	7,641	1.10%	166,509	24.05%	458,616	66.24%	0	0.00%	0	0.00%
Living with Partner	678,726	8,156	1.20%	207,481	30.57%	92,286	13.60%	370,803	54.63%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,157,599	152,476	13.17%	112,764	9.74%	226,368	19.55%	665,990	57.53%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,277,055	70,986	5.56%	44,785	3.51%	574,476	44.98%	579,939	45.41%	6,869	0.54%	0	0.00%
Poverty income ratio 1 or greater	8,837,707	648,656	7.34%	1,129,256	12.78%	2,180,110	24.67%	4,849,522	54.87%	30,163	0.34%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	197,574	9,484	4.80%	17,276	8.74%	53,860	27.26%	86,790	43.93%	30,163	15.27%	0	0.00%
Normal: less than 25	2,464,262	177,590	7.21%	119,320	4.84%	668,157	27.11%	1,499,196	60.84%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	3,755,813	212,433	5.66%	617,200	16.43%	1,091,811	29.07%	1,834,370	48.84%	0	0.00%	0	0.00%
Obese: 30 or higher	4,854,712	472,611	9.74%	533,009	10.98%	1,167,127	24.04%	2,675,095	55.10%	6,869	0.14%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	164,562	42,771	25.99%	37,643	22.87%	34,222	20.80%	43,056	26.16%	6,869	4.17%	0	0.00%
Glycohemoglobin < 5.7%	5,700,739	345,306	6.06%	589,820	10.35%	1,372,538	24.08%	3,393,075	59.52%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	3,774,940	292,408	7.75%	497,597	13.18%	1,035,805	27.44%	1,918,967	50.83%	30,163	0.80%	0	0.00%
Glycohemoglobin 6.5% or higher	1,632,120	191,633	11.74%	161,744	9.91%	538,389	32.99%	740,354	45.36%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,511,112	385,471	15.35%	239,692	9.55%	631,833	25.16%	1,254,116	49.94%	0	0.00%	0	0.00%
No	8,761,248	486,647	5.55%	1,047,113	11.95%	2,349,121	26.81%	4,841,335	55.26%	37,032	0.42%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	7,327,311	448,835	6.13%	958,082	13.08%	1,712,530	23.37%	4,177,701	57.02%	30,163	0.41%	0	0.00%
No	3,907,220	423,283	10.83%	328,722	8.41%	1,230,595	31.50%	1,917,750	49.08%	6,869	0.18%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,194,367	538,707	10.37%	688,357	13.25%	1,115,835	21.48%	2,814,436	54.18%	37,032	0.71%	0	0.00%
No	6,077,994	333,411	5.49%	598,448	9.85%	1,865,119	30.69%	3,281,015	53.98%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,293,992	329,810	14.38%	196,835	8.58%	647,603	28.23%	1,089,581	47.50%	30,163	1.31%	0	0.00%
Mobility limitations	1,878,864	188,014	10.01%	272,990	14.53%	622,205	33.12%	788,786	41.98%	6,869	0.37%	0	0.00%
No mobility limitations	2,913,602	202,561	6.95%	439,971	15.10%	670,788	23.02%	1,600,282	54.92%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	4,148,073	151,732	3.66%	377,009	9.09%	1,002,529	24.17%	2,616,802	63.08%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,229,491	133,824	10.88%	131,559	10.70%	317,264	25.80%	616,681	50.16%	30,163	2.45%	0	0.00%
No	10,013,034	738,294	7.37%	1,125,410	11.24%	2,663,691	26.60%	5,478,770	54.72%	6,869	0.07%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	10,687,174	779,043	7.29%	1,299,159	12.16%	2,623,377	24.55%	5,985,596	56.01%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	1,378,698	144,381	10.47%	203,704	14.78%	444,940	32.27%	585,672	42.48%	0	0.00%	0	0.00%
Once a day	1,429,554	12,631	0.88%	37,003	2.59%	403,613	28.23%	976,307	68.29%	0	0.00%	0	0.00%
2 or more time a week	1,103,780	144,988	13.14%	143,190	12.97%	302,297	27.39%	513,304	46.50%	0	0.00%	0	0.00%
Once a week	763,501	7,219	0.95%	64,995	8.51%	258,860	33.90%	432,428	56.64%	0	0.00%	0	0.00%
1-3 times a month	1,361,707	251,725	18.49%	269,389	19.78%	197,090	14.47%	643,503	47.26%	0	0.00%	0	0.00%
Never	5,100,445	294,327	5.77%	531,357	10.42%	1,340,822	26.29%	2,927,069	57.39%	6,869	0.13%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	463,418	95,422	20.59%	29,073	6.27%	152,196	32.84%	186,727	40.29%	0	0.00%	0	0.00%
Never	10,535,088	752,631	7.14%	1,207,238	11.46%	2,795,425	26.53%	5,772,924	54.80%	6,869	0.07%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	11,866,036	828,068	6.98%	1,389,825	11.71%	3,206,225	27.02%	6,441,919	54.29%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	533,407	0	0.00%	0	0.00%	117,095	21.95%	416,312	78.05%	0	0.00%	0	0.00%
30-39	1,171,423	0	0.00%	40,810	3.48%	264,067	22.54%	866,546	73.97%	0	0.00%	0	0.00%
40-49	1,952,867	58,494	3.00%	180,010	9.22%	478,497	24.50%	1,235,866	63.28%	0	0.00%	0	0.00%
50-59	2,571,971	268,547	10.44%	368,574	14.33%	628,324	24.43%	1,306,526	50.80%	0	0.00%	0	0.00%
60-64	1,611,652	79,563	4.94%	226,455	14.05%	403,856	25.06%	901,778	55.95%	0	0.00%	0	0.00%
65-69	1,077,976	102,725	9.53%	151,950	14.10%	363,816	33.75%	459,486	42.62%	0	0.00%	0	0.00%
70-74	881,903	158,626	17.99%	120,743	13.69%	300,361	34.06%	302,174	34.26%	0	0.00%	0	0.00%
75-79	898,836	65,756	7.32%	90,073	10.02%	291,667	32.45%	451,340	50.21%	0	0.00%	0	0.00%
80+	1,124,309	94,358	8.39%	211,209	18.79%	316,849	28.18%	501,893	44.64%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	1,095,519	39,473	3.60%	54,764	5.00%	285,804	26.09%	715,478	65.31%	0	0.00%	0	0.00%
Other Hispanic	505,517	19,255	3.81%	34,145	6.75%	96,176	19.03%	355,942	70.41%	0	0.00%	0	0.00%
Non-Hispanic White	7,963,037	696,827	8.75%	938,030	11.78%	2,240,597	28.14%	4,087,584	51.33%	0	0.00%	0	0.00%
Non-Hispanic Black	2,016,334	56,942	2.82%	297,830	14.77%	520,469	25.81%	1,141,093	56.59%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	285,629	15,572	5.45%	65,056	22.78%	63,179	22.12%	141,822	49.65%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	300,834	0	0.00%	142,891	47.50%	74,702	24.83%	83,240	27.67%	0	0.00%	0	0.00%
Less Than 9th Grade	1,128,418	119,877	10.62%	90,813	8.05%	286,878	25.42%	630,851	55.91%	0	0.00%	0	0.00%
9-11th Grade	1,543,160	187,436	12.15%	247,032	16.01%	504,056	32.66%	604,635	39.18%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	8,893,625	520,755	5.86%	909,089	10.22%	2,340,589	26.32%	5,123,193	57.61%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	11,866,036	828,068	6.98%	1,389,825	11.71%	3,206,225	27.02%	6,441,919	54.29%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	7,967,730	586,723	7.36%	978,721	12.28%	2,128,421	26.71%	4,273,866	53.64%	0	0.00%	0	0.00%
Widowed	394,066	102,098	25.91%	64,780	16.44%	114,885	29.15%	112,303	28.50%	0	0.00%	0	0.00%
Divorced	1,191,296	52,393	4.40%	104,191	8.75%	351,181	29.48%	683,530	57.38%	0	0.00%	0	0.00%
Separated	210,011	0	0.00%	30,853	14.69%	59,155	28.17%	120,003	57.14%	0	0.00%	0	0.00%
Never Married	1,253,248	31,054	2.48%	134,256	10.71%	384,166	30.65%	703,772	56.16%	0	0.00%	0	0.00%
Living with Partner	747,208	12,061	1.61%	77,024	10.31%	109,678	14.68%	548,446	73.40%	0	0.00%	0	0.00%
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,150,775	12,491	1.09%	102,609	8.92%	375,076	32.59%	660,599	57.40%	0	0.00%	0	0.00%
Poverty income ratio less than 1	1,301,649	52,483	4.03%	84,796	6.51%	208,142	15.99%	956,229	73.46%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	9,413,612	763,095	8.11%	1,202,419	12.77%	2,623,006	27.86%	4,825,091	51.26%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	2,669,139	106,166	3.98%	253,948	9.51%	545,961	20.45%	1,763,064	66.05%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	3,558,307	171,168	4.81%	226,719	6.37%	1,084,597	30.48%	2,075,824	58.34%	0	0.00%	0	0.00%
Obese: 30 or higher	5,368,642	511,461	9.53%	805,233	15.00%	1,478,798	27.55%	2,573,149	47.93%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	6,431,032	271,671	4.22%	453,859	7.06%	1,648,202	25.63%	4,057,300	63.09%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	3,324,840	351,613	10.58%	608,585	18.30%	885,190	26.62%	1,479,453	44.50%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	2,018,694	204,784	10.14%	297,083	14.72%	651,557	32.28%	865,270	42.86%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,236,721	172,422	7.71%	479,595	21.44%	642,912	28.74%	941,792	42.11%	0	0.00%	0	0.00%
No	9,629,315	655,646	6.81%	910,230	9.45%	2,563,313	26.62%	5,500,127	57.12%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	11,866,036	828,068	6.98%	1,389,825	11.71%	3,206,225	27.02%	6,441,919	54.29%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	7,157,497	564,159	7.88%	979,270	13.68%	1,862,208	26.02%	3,751,859	52.42%	0	0.00%	0	0.00%
No	4,708,540	263,909	5.60%	410,554	8.72%	1,344,017	28.54%	2,690,060	57.13%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	5,361,569	408,310	7.62%	881,296	16.44%	1,552,445	28.96%	2,519,518	46.99%	0	0.00%	0	0.00%
No	6,493,583	419,758	6.46%	508,528	7.83%	1,653,779	25.47%	3,911,518	60.24%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	2,015,691	240,245	11.92%	292,845	14.53%	577,417	28.65%	905,184	44.91%	0	0.00%	0	0.00%
Mobility limitations	1,776,064	229,868	12.94%	218,359	12.29%	639,415	36.00%	688,422	38.76%	0	0.00%	0	0.00%
No mobility limitations	3,469,531	133,146	3.84%	609,847	17.58%	1,034,977	29.83%	1,691,560	48.75%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	4,604,751	224,808	4.88%	268,774	5.84%	954,416	20.73%	3,156,754	68.55%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,762,717	161,044	9.14%	454,885	25.81%	512,250	29.06%	634,538	36.00%	0	0.00%	0	0.00%
No	10,096,494	667,024	6.61%	934,940	9.26%	2,687,149	26.61%	5,807,381	57.52%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	1,426,665	137,637	9.65%	139,841	9.80%	407,040	28.53%	742,146	52.02%	0	0.00%	0	0.00%
Once a day	830,369	62,248	7.50%	50,449	6.08%	290,054	34.93%	427,618	51.50%	0	0.00%	0	0.00%
2 or more time a week	992,469	0	0.00%	166,212	16.75%	221,766	22.34%	604,491	60.91%	0	0.00%	0	0.00%
Once a week	829,676	121,163	14.60%	107,577	12.97%	182,839	22.04%	418,097	50.39%	0	0.00%	0	0.00%
1-3 times a month	2,165,283	95,544	4.41%	147,635	6.82%	585,295	27.03%	1,336,809	61.74%	0	0.00%	0	0.00%
Never	5,366,897	363,089	6.77%	755,301	14.07%	1,366,387	25.46%	2,882,121	53.70%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M03a - Frequency of bladder control problems among those who responded 'yes' to urge urinary incontinence among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	11,866,036	828,068	6.98%	1,389,825	11.71%	3,206,225	27.02%	6,441,919	54.29%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	392,062	13,265	3.38%	36,394	9.28%	150,365	38.35%	192,037	48.98%	0	0.00%	0	0.00%
Never	11,040,333	766,416	6.94%	1,268,118	11.49%	2,799,748	25.36%	6,206,051	56.21%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on KIQ.045: How frequently does this (referring to KIQ.044) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?