

Table M04 - Prevalence of mixed urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	82,861,861	2,305,061	2.78%	2.20%	3.36%	80,556,800	97.22%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	311,061	0	0.00%	0.00%	0.00%	311,061	100.00%
20-29	11,424,981	0	0.00%	0.00%	0.00%	11,424,981	100.00%
30-39	17,162,219	156,434	0.91%	( 0.24%)	2.06%	17,005,786	99.09%
40-49	21,499,211	169,904	0.79%	( 0.38%)	1.96%	21,329,307	99.21%
50-59	17,107,411	404,942	2.37%	0.21%	4.53%	16,702,469	97.63%
60-64	4,047,823	244,787	6.05%	1.88%	10.21%	3,803,036	93.95%
65-69	3,123,223	260,203	8.33%	2.64%	14.02%	2,863,020	91.67%
70-74	3,029,815	364,192	12.02%	2.99%	21.05%	2,665,623	87.98%
75-79	2,514,807	300,780	11.96%	4.75%	19.17%	2,214,027	88.04%
80-84	1,722,977	217,898	12.65%	6.98%	18.32%	1,505,079	87.35%
85+	918,333	185,922	20.25%	6.90%	33.60%	732,411	79.75%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	6,712,363	98,936	1.47%	0.36%	2.59%	6,613,427	98.53%
Other Hispanic	4,509,971	97,150	2.15%	( 0.26%)	4.56%	4,412,820	97.85%
Non-Hispanic White	60,704,443	1,857,584	3.06%	2.09%	4.03%	58,846,858	96.94%
Non-Hispanic Black	8,015,474	251,390	3.14%	0.59%	5.68%	7,764,084	96.86%
Other Race - Including Multi-Racial	2,919,610	0	0.00%	0.00%	0.00%	2,919,610	100.00%
<b>Education</b>	-	-	-	-	-	-	-
Missing	875,696	24,453	2.79%	( 3.59%)	9.17%	851,243	97.21%
Less Than 9th Grade	5,790,082	543,028	9.38%	4.46%	14.30%	5,247,055	90.62%
9-11th Grade	9,157,055	282,887	3.09%	0.96%	5.21%	8,874,168	96.91%
High School Grad/GED or Equivalent	66,935,057	1,454,693	2.17%	1.45%	2.90%	65,480,364	97.83%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	777,526	21,480	2.76%	( 3.29%)	8.81%	756,047	97.24%
Married	54,968,480	1,674,895	3.05%	2.40%	3.70%	53,293,585	96.95%
Widowed	3,484,033	292,798	8.40%	1.97%	14.84%	3,191,235	91.60%
Divorced	6,845,124	242,256	3.54%	( 0.20%)	7.28%	6,602,868	96.46%
Separated	2,054,234	2,737	0.13%	( 0.15%)	0.42%	2,051,498	99.87%
Never Married	9,291,638	18,704	0.20%	( 0.12%)	0.52%	9,272,934	99.80%
Living with Partner	5,424,742	52,191	0.96%	( 0.99%)	2.91%	5,372,551	99.04%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	4,559,321	141,495	3.10%	( 0.71%)	6.92%	4,417,826	96.90%
Poverty income ratio less than 1	8,596,310	314,223	3.66%	1.78%	5.53%	8,282,087	96.34%
Poverty income ratio 1 or greater	69,706,229	1,849,343	2.65%	2.00%	3.31%	67,856,887	97.35%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	2,785,488	456,937	16.40%	5.89%	26.92%	2,328,551	83.60%
Normal: less than 25	25,061,639	360,607	1.44%	0.75%	2.12%	24,701,031	98.56%
Overweight: 25.0 - 29.9	33,519,832	646,217	1.93%	0.86%	3.00%	32,873,615	98.07%
Obese: 30 or higher	21,494,902	841,300	3.91%	2.44%	5.39%	20,653,602	96.09%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	2,773,096	48,080	1.73%	( 0.16%)	3.63%	2,725,016	98.27%
Glycohemoglobin < 5.7%	65,352,681	1,326,263	2.03%	1.42%	2.63%	64,026,419	97.97%
Glycohemoglobin 5.7% - 6.4%	9,906,441	458,279	4.63%	3.00%	6.25%	9,448,162	95.37%
Glycohemoglobin 6.5% or higher	4,829,643	472,439	9.78%	3.05%	16.51%	4,357,203	90.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	82,861,861	2,305,061	2.78%	2.20%	3.36%	80,556,800	97.22%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	5,365,838	476,287	8.88%	3.12%	14.63%	4,889,551	91.12%
No	77,496,023	1,828,774	2.36%	1.85%	2.87%	75,667,249	97.64%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	45,717,574	1,513,325	3.31%	2.49%	4.13%	44,204,249	96.69%
No	37,057,126	791,736	2.14%	1.36%	2.92%	36,265,390	97.86%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	1,036,586	0	0.00%	0.00%	0.00%	1,036,586	100.00%
Yes	18,889,863	1,170,717	6.20%	4.17%	8.22%	17,719,146	93.80%
No	62,838,671	1,134,344	1.81%	1.18%	2.43%	61,704,328	98.19%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	4,552,204	576,826	12.67%	7.11%	18.24%	3,975,379	87.33%
Mobility limitations	5,391,917	722,776	13.40%	6.71%	20.10%	4,669,141	86.60%
No mobility limitations	15,962,311	790,113	4.95%	2.49%	7.41%	15,172,199	95.05%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	56,940,813	200,732	0.35%	( 0.10%)	0.80%	56,740,081	99.65%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	82,861,861	2,305,061	2.78%	2.20%	3.36%	80,556,800	97.22%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	82,861,861	2,305,061	2.78%	2.20%	3.36%	80,556,800	97.22%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	82,861,861	2,305,061	2.78%	2.20%	3.36%	80,556,800	97.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	81,587,046	3,025,438	3.71%	2.50%	4.92%	78,561,608	96.29%
<b>Age at screening</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	538,199	0	0.00%	0.00%	0.00%	538,199	100.00%
20-29	11,585,094	65,286	0.56%	( 0.26%)	1.39%	11,519,808	99.44%
30-39	16,202,663	239,956	1.48%	0.12%	2.84%	15,962,707	98.52%
40-49	19,495,889	335,238	1.72%	0.16%	3.28%	19,160,650	98.28%
50-59	15,656,581	303,600	1.94%	0.61%	3.27%	15,352,981	98.06%
60-64	4,879,364	213,733	4.38%	1.30%	7.46%	4,665,631	95.62%
65-69	4,860,365	361,371	7.44%	2.57%	12.30%	4,498,994	92.56%
70-74	3,086,687	452,358	14.66%	8.79%	20.52%	2,634,329	85.34%
75-79	2,552,001	547,062	21.44%	10.43%	32.44%	2,004,939	78.56%
80-84	1,722,779	320,478	18.60%	8.77%	28.44%	1,402,301	81.40%
85+	1,007,424	186,356	18.50%	2.71%	34.28%	821,068	81.50%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	6,565,119	137,224	2.09%	0.63%	3.55%	6,427,895	97.91%
Other Hispanic	2,542,163	67,083	2.64%	0.17%	5.11%	2,475,080	97.36%
Non-Hispanic White	60,511,937	2,340,202	3.87%	2.52%	5.22%	58,171,735	96.13%
Non-Hispanic Black	7,586,602	399,025	5.26%	2.67%	7.85%	7,187,577	94.74%
Other Race - Including Multi-Racial	4,381,225	81,904	1.87%	( 1.10%)	4.84%	4,299,321	98.13%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,671,854	45,896	2.75%	( 0.89%)	6.38%	1,625,959	97.25%
Less Than 9th Grade	5,497,165	338,834	6.16%	2.55%	9.78%	5,158,330	93.84%
9-11th Grade	8,409,336	442,044	5.26%	1.08%	9.43%	7,967,292	94.74%
High School Grad/GED or Equivalent	65,698,647	2,198,664	3.35%	2.16%	4.54%	63,499,983	96.65%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	810,294	6,188	0.76%	( 0.84%)	2.37%	804,106	99.24%
Married	53,783,193	2,097,210	3.90%	2.47%	5.33%	51,685,983	96.10%
Widowed	3,014,662	271,335	9.00%	1.89%	16.11%	2,743,327	91.00%
Divorced	6,687,276	153,135	2.29%	( 0.03%)	4.61%	6,534,140	97.71%
Separated	1,276,504	105,472	8.26%	( 0.51%)	17.04%	1,171,032	91.74%
Never Married	10,203,659	364,192	3.57%	0.32%	6.82%	9,839,467	96.43%
Living with Partner	5,419,747	27,906	0.51%	( 0.54%)	1.57%	5,391,842	99.49%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	4,209,934	301,143	7.15%	3.49%	10.81%	3,908,792	92.85%
Poverty income ratio less than 1	8,449,158	265,671	3.14%	0.82%	5.47%	8,183,487	96.86%
Poverty income ratio 1 or greater	68,927,954	2,458,625	3.57%	2.09%	5.04%	66,469,329	96.43%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	795,850	102,948	12.94%	( 3.44%)	29.31%	692,902	87.06%
Normal: less than 25	25,353,550	912,049	3.60%	2.17%	5.02%	24,441,501	96.40%
Overweight: 25.0 - 29.9	31,787,701	972,448	3.06%	1.45%	4.67%	30,815,253	96.94%
Obese: 30 or higher	23,649,946	1,037,993	4.39%	2.48%	6.29%	22,611,953	95.61%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	81,587,046	3,025,438	3.71%	2.50%	4.92%	78,561,608	96.29%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	2,159,779	78,565	3.64%	( 0.64%)	7.92%	2,081,214	96.36%
Glycohemoglobin < 5.7%	63,568,107	1,867,637	2.94%	1.84%	4.04%	61,700,470	97.06%
Glycohemoglobin 5.7% - 6.4%	11,119,844	841,914	7.57%	3.94%	11.21%	10,277,931	92.43%
Glycohemoglobin 6.5% or higher	4,739,316	237,322	5.01%	1.19%	8.83%	4,501,994	94.99%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	5,241,223	499,939	9.54%	5.03%	14.05%	4,741,284	90.46%
No	76,345,823	2,525,499	3.31%	2.18%	4.43%	73,820,324	96.69%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	46,783,271	2,212,678	4.73%	3.29%	6.17%	44,570,593	95.27%
No	34,765,562	800,929	2.30%	1.21%	3.40%	33,964,633	97.70%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	755,859	0	0.00%	0.00%	0.00%	755,859	100.00%
Yes	23,781,141	1,706,423	7.18%	4.34%	10.01%	22,074,718	92.82%
No	56,993,923	1,319,015	2.31%	1.60%	3.03%	55,674,908	97.69%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	7,345,644	882,175	12.01%	6.91%	17.11%	6,463,469	87.99%
Mobility limitations	6,684,849	666,481	9.97%	6.00%	13.94%	6,018,368	90.03%
No mobility limitations	17,399,529	1,072,482	6.16%	3.60%	8.73%	16,327,047	93.84%
N/A - Respondent not asked mobility limitation questions	50,157,025	404,301	0.81%	0.13%	1.48%	49,752,724	99.19%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
N/A for these years of data	81,587,046	3,025,438	3.71%	2.50%	4.92%	78,561,608	96.29%
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
N/A for these years of data	81,587,046	3,025,438	3.71%	2.50%	4.92%	78,561,608	96.29%
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
N/A for these years of data	81,587,046	3,025,438	3.71%	2.50%	4.92%	78,561,608	96.29%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	82,565,210	1,403,468	1.70%	0.88%	2.52%	81,161,742	98.30%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	362,271	0	0.00%	0.00%	0.00%	362,271	100.00%
20-29	13,838,661	63,691	0.46%	( 0.12%)	1.04%	13,774,969	99.54%
30-39	16,735,271	90,019	0.54%	( 0.33%)	1.40%	16,645,252	99.46%
40-49	19,618,066	371,561	1.89%	( 0.04%)	3.83%	19,246,505	98.11%
50-59	14,937,078	217,883	1.46%	( 0.09%)	3.00%	14,719,195	98.54%
60-64	5,491,073	25,338	0.46%	( 0.16%)	1.08%	5,465,735	99.54%
65-69	4,039,777	123,864	3.07%	0.41%	5.72%	3,915,913	96.93%
70-74	3,273,717	191,655	5.85%	2.04%	9.67%	3,082,062	94.15%
75-79	1,890,710	146,362	7.74%	0.64%	14.84%	1,744,348	92.26%
80-84	1,620,348	162,313	10.02%	( 2.77%)	22.80%	1,458,035	89.98%
85+	758,239	10,782	1.42%	( 1.46%)	4.31%	747,457	98.58%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,032,548	77,475	1.10%	0.15%	2.06%	6,955,073	98.90%
Other Hispanic	2,647,840	70,865	2.68%	( 2.02%)	7.38%	2,576,975	97.32%
Non-Hispanic White	60,876,643	1,114,528	1.83%	0.78%	2.88%	59,762,115	98.17%
Non-Hispanic Black	8,282,136	140,600	1.70%	1.19%	2.20%	8,141,535	98.30%
Other Race - Including Multi-Racial	3,726,043	0	0.00%	0.00%	0.00%	3,726,043	100.00%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,104,602	0	0.00%	0.00%	0.00%	1,104,602	100.00%
Less Than 9th Grade	5,431,179	22,335	0.41%	( 0.16%)	0.99%	5,408,844	99.59%
9-11th Grade	9,087,527	212,515	2.34%	0.81%	3.87%	8,875,012	97.66%
High School Grad/GED or Equivalent	66,810,640	1,155,741	1.73%	0.72%	2.74%	65,654,899	98.27%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	951,242	12,877	1.35%	( 0.92%)	3.63%	938,365	98.65%
Married	52,372,865	726,455	1.39%	0.59%	2.18%	51,646,410	98.61%
Widowed	2,569,871	52,510	2.04%	( 1.12%)	5.21%	2,517,361	97.96%
Divorced	7,687,426	178,084	2.32%	( 0.49%)	5.12%	7,509,342	97.68%
Separated	2,348,521	108,280	4.61%	( 4.98%)	14.21%	2,240,241	95.39%
Never Married	8,747,874	224,342	2.56%	( 0.08%)	5.21%	8,523,532	97.44%
Living with Partner	7,887,410	100,919	1.28%	( 1.15%)	3.71%	7,786,491	98.72%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	2,353,664	76,574	3.25%	( 0.51%)	7.01%	2,277,090	96.75%
Poverty income ratio less than 1	7,905,730	221,599	2.80%	( 0.43%)	6.03%	7,684,131	97.20%
Poverty income ratio 1 or greater	72,305,816	1,105,295	1.53%	0.67%	2.38%	71,200,521	98.47%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	915,669	0	0.00%	0.00%	0.00%	915,669	100.00%
Normal: less than 25	23,297,411	192,122	0.82%	( 0.20%)	1.85%	23,105,289	99.18%
Overweight: 25.0 - 29.9	31,749,626	405,456	1.28%	0.42%	2.14%	31,344,170	98.72%
Obese: 30 or higher	26,602,504	805,889	3.03%	0.92%	5.14%	25,796,615	96.97%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,307,812	0	0.00%	0.00%	0.00%	3,307,812	100.00%
Glycohemoglobin < 5.7%	63,312,374	751,605	1.19%	0.57%	1.81%	62,560,768	98.81%
Glycohemoglobin 5.7% - 6.4%	11,579,354	330,193	2.85%	0.75%	4.96%	11,249,161	97.15%
Glycohemoglobin 6.5% or higher	4,365,671	321,670	7.37%	1.12%	13.62%	4,044,000	92.63%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	82,565,210	1,403,468	1.70%	0.88%	2.52%	81,161,742	98.30%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	5,551,706	262,444	4.73%	0.29%	9.16%	5,289,261	95.27%
No	77,013,504	1,141,024	1.48%	0.66%	2.30%	75,872,480	98.52%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	45,296,005	760,868	1.68%	0.83%	2.53%	44,535,136	98.32%
No	37,269,205	642,600	1.72%	0.37%	3.08%	36,626,605	98.28%
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	21,715,279	936,576	4.31%	1.76%	6.86%	20,778,703	95.69%
No	60,728,047	466,892	0.77%	0.24%	1.30%	60,261,155	99.23%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	5,724,886	307,753	5.38%	0.53%	10.22%	5,417,134	94.62%
Mobility limitations	4,235,028	278,162	6.57%	2.05%	11.09%	3,956,866	93.43%
No mobility limitations	16,807,617	403,056	2.40%	1.20%	3.59%	16,404,561	97.60%
N/A - Respondent not asked mobility limitation questions	55,797,678	414,497	0.74%	0.03%	1.46%	55,383,182	99.26%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,154,746	171,495	2.79%	( 0.21%)	5.79%	5,983,251	97.21%
No	76,294,361	1,231,973	1.61%	0.74%	2.49%	75,062,388	98.39%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	317,258	9,956	3.14%	( 3.77%)	10.05%	307,302	96.86%
2 or more times a day	11,506,127	400,473	3.48%	0.68%	6.28%	11,105,654	96.52%
Once a day	7,582,496	80,218	1.06%	( 0.41%)	2.52%	7,502,278	98.94%
2 or more time a week	4,951,461	253,833	5.13%	0.15%	10.11%	4,697,628	94.87%
Once a week	4,730,157	166,516	3.52%	( 0.46%)	7.50%	4,563,641	96.48%
1-3 times a month	7,612,041	123,691	1.62%	0.01%	3.24%	7,488,350	98.38%
Never	45,572,854	368,781	0.81%	0.09%	1.53%	45,204,073	99.19%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	317,258	9,956	3.14%	( 3.77%)	10.05%	307,302	96.86%
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*
Never	81,685,219	1,285,081	1.57%	0.79%	2.35%	80,400,138	98.43%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	84,047,163	2,238,601	2.66%	1.80%	3.52%	81,808,562	97.34%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	566,180	0	0.00%	0.00%	0.00%	566,180	100.00%
20-29	13,763,466	24,497	0.18%	( 0.19%)	0.55%	13,738,969	99.82%
30-39	16,110,637	121,728	0.76%	( 0.07%)	1.58%	15,988,909	99.24%
40-49	19,529,956	446,777	2.29%	( 0.14%)	4.71%	19,083,180	97.71%
50-59	16,469,888	531,527	3.23%	1.82%	4.64%	15,938,361	96.77%
60-64	6,127,236	194,131	3.17%	1.05%	5.28%	5,933,104	96.83%
65-69	4,223,104	341,147	8.08%	1.03%	15.13%	3,881,958	91.92%
70-74	2,764,113	158,854	5.75%	1.09%	10.41%	2,605,259	94.25%
75-79	2,097,243	157,676	7.52%	2.83%	12.20%	1,939,566	92.48%
80+	2,395,340	262,264	10.95%	4.50%	17.40%	2,133,076	89.05%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,996,334	121,892	1.52%	0.01%	3.04%	7,874,442	98.48%
Other Hispanic	4,020,331	81,768	2.03%	( 0.35%)	4.42%	3,938,564	97.97%
Non-Hispanic White	58,744,416	1,514,494	2.58%	1.59%	3.56%	57,229,921	97.42%
Non-Hispanic Black	8,063,224	226,682	2.81%	0.82%	4.81%	7,836,542	97.19%
Other Race - Including Multi-Racial	5,222,858	293,764	5.62%	( 3.30%)	14.54%	4,929,094	94.38%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,147,831	44,574	3.88%	( 0.77%)	8.54%	1,103,257	96.12%
Less Than 9th Grade	5,287,150	295,432	5.59%	2.46%	8.72%	4,991,718	94.41%
9-11th Grade	10,195,422	387,113	3.80%	2.04%	5.56%	9,808,309	96.20%
High School Grad/GED or Equivalent	67,361,899	1,511,482	2.24%	1.29%	3.20%	65,850,417	97.76%
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	1,232,787	0	0.00%	0.00%	0.00%	1,232,787	100.00%
Married	53,481,118	1,547,568	2.89%	1.66%	4.12%	51,933,550	97.11%
Widowed	2,594,443	143,849	5.54%	1.17%	9.92%	2,450,594	94.46%
Divorced	6,851,144	110,591	1.61%	0.29%	2.94%	6,740,553	98.39%
Separated	1,950,636	190,980	9.79%	( 0.30%)	19.88%	1,759,656	90.21%
Never Married	11,726,576	112,153	0.96%	( 0.01%)	1.92%	11,614,423	99.04%
Living with Partner	6,025,088	133,460	2.22%	0.05%	4.38%	5,891,628	97.78%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	5,974,161	413,643	6.92%	2.86%	10.98%	5,560,518	93.08%
Poverty income ratio less than 1	10,369,571	409,307	3.95%	0.47%	7.43%	9,960,264	96.05%
Poverty income ratio 1 or greater	67,703,431	1,415,651	2.09%	1.33%	2.85%	66,287,780	97.91%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	633,121	35,914	5.67%	( 3.44%)	14.79%	597,206	94.33%
Normal: less than 25	24,327,266	469,681	1.93%	0.85%	3.02%	23,857,585	98.07%
Overweight: 25.0 - 29.9	33,014,084	677,650	2.05%	0.45%	3.66%	32,336,434	97.95%
Obese: 30 or higher	26,072,693	1,055,356	4.05%	2.31%	5.78%	25,017,337	95.95%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	3,442,085	74,759	2.17%	0.58%	3.77%	3,367,326	97.83%
Glycohemoglobin < 5.7%	57,199,430	979,321	1.71%	0.99%	2.44%	56,220,109	98.29%
Glycohemoglobin 5.7% - 6.4%	17,547,492	761,604	4.34%	2.70%	5.98%	16,785,888	95.66%
Glycohemoglobin 6.5% or higher	5,858,157	422,918	7.22%	0.62%	13.82%	5,435,239	92.78%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	84,047,163	2,238,601	2.66%	1.80%	3.52%	81,808,562	97.34%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,194,520	411,164	6.64%	2.08%	11.19%	5,783,356	93.36%
No	77,852,644	1,827,437	2.35%	1.57%	3.13%	76,025,207	97.65%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	44,952,245	1,517,708	3.38%	2.11%	4.64%	43,434,537	96.62%
No	39,064,893	720,893	1.85%	0.95%	2.74%	38,344,000	98.15%
Don't know	*	*	*	*	*	*	*
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	23,077,882	1,023,247	4.43%	2.78%	6.09%	22,054,635	95.57%
No	60,866,669	1,215,354	2.00%	1.04%	2.95%	59,651,314	98.00%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	7,093,315	705,520	9.95%	5.09%	14.80%	6,387,795	90.05%
Mobility limitations	5,298,935	607,942	11.47%	5.89%	17.06%	4,690,994	88.53%
No mobility limitations	15,862,314	330,970	2.09%	0.61%	3.57%	15,531,344	97.91%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	55,792,600	594,169	1.06%	0.15%	1.98%	55,198,430	98.94%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,806,639	464,811	6.83%	2.28%	11.38%	6,341,828	93.17%
No	77,028,361	1,743,954	2.26%	1.49%	3.04%	75,284,407	97.74%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	531,084	37,943	7.14%	( 2.98%)	17.27%	493,141	92.86%
2 or more times a day	9,146,384	416,901	4.56%	1.65%	7.47%	8,729,483	95.44%
Once a day	6,842,679	104,676	1.53%	( 0.20%)	3.26%	6,738,003	98.47%
2 or more time a week	4,872,324	291,566	5.98%	2.60%	9.37%	4,580,758	94.02%
Once a week	3,809,094	270,295	7.10%	( 0.14%)	14.33%	3,538,799	92.90%
1-3 times a month	6,688,465	174,110	2.60%	0.61%	4.59%	6,514,355	97.40%
Never	51,793,242	943,111	1.82%	0.87%	2.77%	50,850,131	98.18%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	531,084	37,943	7.14%	( 2.98%)	17.27%	493,141	92.86%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	641,292	169,834	26.48%	( 8.33%)	61.30%	471,458	73.52%
Never	82,256,056	1,991,848	2.42%	1.64%	3.20%	80,264,208	97.58%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).



Table M04 - Prevalence of mixed urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	83,496,634	1,744,161	2.09%	1.52%	2.66%	81,752,473	97.91%
<b>Age at screening</b>	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*
20-29	12,417,242	70,629	0.57%	( 0.19%)	1.33%	12,346,613	99.43%
30-39	15,393,062	172,227	1.12%	( 0.11%)	2.35%	15,220,835	98.88%
40-49	19,044,118	83,542	0.44%	( 0.13%)	1.00%	18,960,576	99.56%
50-59	18,767,403	434,453	2.31%	0.74%	3.89%	18,332,950	97.69%
60-64	5,334,674	45,493	0.85%	( 0.33%)	2.04%	5,289,181	99.15%
65-69	5,043,134	216,414	4.29%	1.09%	7.49%	4,826,720	95.71%
70-74	2,927,120	192,008	6.56%	1.87%	11.25%	2,735,112	93.44%
75-79	2,279,097	249,917	10.97%	2.41%	19.52%	2,029,180	89.03%
80+	2,034,840	279,479	13.73%	3.24%	24.23%	1,755,361	86.27%
<b>Race/ethnicity</b>	-	-	-	-	-	-	-
Mexican American	7,942,513	57,760	0.73%	( 0.22%)	1.67%	7,884,753	99.27%
Other Hispanic	4,393,341	85,606	1.95%	( 0.66%)	4.55%	4,307,734	98.05%
Non-Hispanic White	57,877,241	1,250,156	2.16%	1.48%	2.84%	56,627,086	97.84%
Non-Hispanic Black	7,930,382	325,291	4.10%	2.71%	5.49%	7,605,090	95.90%
Other Race - Including Multi-Racial	5,353,157	25,347	0.47%	0.17%	0.77%	5,327,810	99.53%
<b>Education</b>	-	-	-	-	-	-	-
Missing	1,129,081	70,472	6.24%	( 4.77%)	17.25%	1,058,609	93.76%
Less Than 9th Grade	5,472,566	206,041	3.76%	1.97%	5.56%	5,266,525	96.24%
9-11th Grade	8,756,435	194,426	2.22%	0.34%	4.10%	8,562,008	97.78%
High School Grad/GED or Equivalent	67,992,090	1,273,222	1.87%	1.34%	2.40%	66,718,869	98.13%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Marital status</b>	-	-	-	-	-	-	-
Missing	271,492	15,549	5.73%	( 6.72%)	18.17%	255,943	94.27%
Married	52,693,097	1,071,140	2.03%	1.26%	2.81%	51,621,957	97.97%
Widowed	2,865,482	92,987	3.25%	( 1.56%)	8.05%	2,772,495	96.75%
Divorced	7,630,334	205,361	2.69%	0.18%	5.20%	7,424,974	97.31%
Separated	2,037,540	62,760	3.08%	( 0.98%)	7.14%	1,974,780	96.92%
Never Married	11,195,125	183,795	1.64%	0.78%	2.50%	11,011,329	98.36%
Living with Partner	6,763,717	84,378	1.25%	( 0.16%)	2.66%	6,679,339	98.75%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
<b>Poverty income ratio</b>	-	-	-	-	-	-	-
Missing	5,315,778	51,368	0.97%	( 0.59%)	2.52%	5,264,409	99.03%
Poverty income ratio less than 1	9,911,195	295,166	2.98%	1.45%	4.50%	9,616,030	97.02%
Poverty income ratio 1 or greater	68,269,661	1,397,626	2.05%	1.41%	2.69%	66,872,034	97.95%
<b>Body mass index</b>	-	-	-	-	-	-	-
Missing	658,587	36,087	5.48%	( 3.34%)	14.30%	622,500	94.52%
Normal: less than 25	21,885,457	578,929	2.65%	1.30%	3.99%	21,306,529	97.35%
Overweight: 25.0 - 29.9	32,535,423	470,545	1.45%	0.57%	2.32%	32,064,879	98.55%
Obese: 30 or higher	28,417,166	658,600	2.32%	1.06%	3.58%	27,758,566	97.68%
<b>Glycohemoglobin</b>	-	-	-	-	-	-	-
Missing	2,524,945	33,513	1.33%	( 0.63%)	3.28%	2,491,432	98.67%
Glycohemoglobin < 5.7%	55,007,584	860,915	1.57%	0.75%	2.38%	54,146,669	98.43%
Glycohemoglobin 5.7% - 6.4%	19,445,791	483,350	2.49%	1.33%	3.64%	18,962,441	97.51%
Glycohemoglobin 6.5% or higher	6,518,314	366,383	5.62%	1.78%	9.46%	6,151,932	94.38%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M04 - Prevalence of mixed urinary incontinence among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
<b>Total</b>	83,496,634	1,744,161	2.09%	1.52%	2.66%	81,752,473	97.91%
<b>Diabetes</b>	-	-	-	-	-	-	-
Yes	6,818,210	414,248	6.08%	1.94%	10.21%	6,403,962	93.92%
No	76,678,424	1,329,912	1.73%	1.19%	2.28%	75,348,511	98.27%
<b>Smoked at least 100 cigarettes in life</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	40,357,688	1,128,012	2.80%	1.67%	3.92%	39,229,675	97.20%
No	43,138,946	616,148	1.43%	0.88%	1.98%	42,522,798	98.57%
<b>High blood pressure</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	23,886,929	957,627	4.01%	2.65%	5.37%	22,929,302	95.99%
No	59,509,272	786,534	1.32%	0.80%	1.84%	58,722,738	98.68%
Don't know	*	*	*	*	*	*	*
<b>Mobility limitation</b>	-	-	-	-	-	-	-
Missing	5,909,147	322,715	5.46%	2.98%	7.95%	5,586,432	94.54%
Mobility limitations	5,451,945	447,760	8.21%	4.97%	11.45%	5,004,185	91.79%
No mobility limitations	16,599,968	599,579	3.61%	1.72%	5.50%	16,000,390	96.39%
N/A - Respondent not asked mobility limitation questions	55,535,573	374,107	0.67%	0.13%	1.22%	55,161,466	99.33%
<b>Sleep disorder</b>	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	6,217,978	360,298	5.79%	0.32%	11.27%	5,857,680	94.21%
No	77,082,132	1,383,863	1.80%	1.09%	2.50%	75,698,269	98.20%
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - gas</b>	-	-	-	-	-	-	-
Missing	253,049	48,387	19.12%	( 5.96%)	44.20%	204,661	80.88%
2 or more times a day	7,986,704	367,031	4.60%	1.11%	8.09%	7,619,673	95.40%
Once a day	7,784,349	127,709	1.64%	0.25%	3.03%	7,656,640	98.36%
2 or more time a week	5,865,983	178,869	3.05%	0.38%	5.72%	5,687,115	96.95%
Once a week	4,097,733	180,356	4.40%	( 0.95%)	9.76%	3,917,378	95.60%
1-3 times a month	8,167,068	339,647	4.16%	1.61%	6.71%	7,827,422	95.84%
Never	49,278,072	502,163	1.02%	0.60%	1.44%	48,775,909	98.98%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
<b>Bowel leakage - stool</b>	-	-	-	-	-	-	-
Missing	253,049	48,387	19.12%	( 5.96%)	44.20%	204,661	80.88%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	575,394	155,439	27.01%	10.41%	43.62%	419,954	72.99%
Never	82,281,081	1,510,165	1.84%	1.22%	2.45%	80,770,916	98.16%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: \* Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).