

Table M05 - Prevalence of insensate urine loss among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	90,411,698	2,952,597	3.27%	2.58%	3.95%	87,459,101	96.73%
Age at screening	-	-	-	-	-	-	-
10-19	318,425	7,364	2.31%	(2.96%)	7.59%	311,061	97.69%
20-29	11,767,942	30,739	0.26%	(0.30%)	0.83%	11,737,203	99.74%
30-39	18,212,868	52,207	0.29%	(0.18%)	0.75%	18,160,661	99.71%
40-49	22,595,280	335,502	1.48%	0.04%	2.93%	22,259,779	98.52%
50-59	18,540,781	683,981	3.69%	1.82%	5.56%	17,856,800	96.31%
60-64	4,648,416	249,754	5.37%	1.50%	9.25%	4,398,663	94.63%
65-69	3,936,729	305,063	7.75%	3.33%	12.17%	3,631,667	92.25%
70-74	4,053,453	437,470	10.79%	4.44%	17.14%	3,615,983	89.21%
75-79	2,934,099	348,796	11.89%	6.08%	17.69%	2,585,303	88.11%
80-84	2,102,740	247,548	11.77%	7.21%	16.33%	1,855,192	88.23%
85+	1,300,965	254,173	19.54%	8.42%	30.66%	1,046,792	80.46%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,055,890	157,894	2.24%	0.04%	4.43%	6,897,996	97.76%
Other Hispanic	4,938,236	129,497	2.62%	(0.40%)	5.64%	4,808,739	97.38%
Non-Hispanic White	66,271,493	2,346,031	3.54%	2.49%	4.59%	63,925,462	96.46%
Non-Hispanic Black	9,016,771	245,085	2.72%	0.89%	4.55%	8,771,686	97.28%
Other Race - Including Multi-Racial	3,129,308	74,091	2.37%	(2.05%)	6.79%	3,055,218	97.63%
Education	-	-	-	-	-	-	-
Missing	933,223	35,669	3.82%	(2.77%)	10.41%	897,554	96.18%
Less Than 9th Grade	6,516,375	407,756	6.26%	3.96%	8.56%	6,108,619	93.74%
9-11th Grade	10,142,865	376,336	3.71%	0.61%	6.82%	9,766,529	96.29%
High School Grad/GED or Equivalent	72,672,406	2,100,490	2.89%	1.96%	3.82%	70,571,915	97.11%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	909,023	38,014	4.18%	(4.66%)	13.03%	871,009	95.82%
Married	59,642,236	2,095,650	3.51%	2.68%	4.35%	57,546,586	96.49%
Widowed	4,052,855	327,563	8.08%	3.24%	12.92%	3,725,291	91.92%
Divorced	7,804,451	317,231	4.06%	0.30%	7.83%	7,487,220	95.94%
Separated	2,163,795	2,737	0.13%	(0.14%)	0.40%	2,161,058	99.87%
Never Married	10,212,194	169,351	1.66%	0.24%	3.07%	10,042,843	98.34%
Living with Partner	5,531,980	2,051	0.04%	(0.04%)	0.12%	5,529,929	99.96%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,980,213	192,238	3.86%	(0.76%)	8.48%	4,787,975	96.14%
Poverty income ratio less than 1	9,683,534	526,018	5.43%	2.46%	8.41%	9,157,517	94.57%
Poverty income ratio 1 or greater	75,747,951	2,234,342	2.95%	2.24%	3.66%	73,513,609	97.05%
Body mass index	-	-	-	-	-	-	-
Missing	3,214,827	475,220	14.78%	7.70%	21.86%	2,739,607	85.22%
Normal: less than 25	27,368,328	581,300	2.12%	0.72%	3.53%	26,787,028	97.88%
Overweight: 25.0 - 29.9	36,211,291	1,050,439	2.90%	1.65%	4.15%	35,160,852	97.10%
Obese: 30 or higher	23,617,252	845,638	3.58%	2.58%	4.58%	22,771,613	96.42%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,165,821	86,141	2.72%	0.93%	4.51%	3,079,681	97.28%
Glycohemoglobin < 5.7%	69,910,044	1,920,404	2.75%	1.95%	3.55%	67,989,640	97.25%
Glycohemoglobin 5.7% - 6.4%	11,200,635	580,748	5.18%	2.24%	8.13%	10,619,887	94.82%
Glycohemoglobin 6.5% or higher	6,135,198	365,304	5.95%	2.14%	9.77%	5,769,894	94.05%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2001 - 2002

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	83,496,634	1,744,161	2.09%	1.52%	2.66%	81,752,473	97.91%
Diabetes	-	-	-	-	-	-	-
Yes	6,600,753	426,056	6.45%	3.24%	9.67%	6,174,698	93.55%
No	83,810,945	2,526,542	3.01%	2.37%	3.66%	81,284,403	96.99%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	51,264,387	2,001,836	3.90%	3.10%	4.71%	49,262,551	96.10%
No	39,060,150	950,762	2.43%	1.46%	3.41%	38,109,389	97.57%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	1,046,623	0	0.00%	0.00%	0.00%	1,046,623	100.00%
Yes	21,541,683	1,336,170	6.20%	4.61%	7.79%	20,205,513	93.80%
No	67,726,653	1,616,427	2.39%	1.41%	3.36%	66,110,225	97.61%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	5,360,610	627,786	11.71%	5.08%	18.35%	4,732,823	88.29%
Mobility limitations	6,911,054	587,782	8.50%	4.48%	12.53%	6,323,272	91.50%
No mobility limitations	18,453,406	955,378	5.18%	3.44%	6.92%	17,498,028	94.82%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	59,672,014	767,037	1.29%	0.57%	2.00%	58,904,977	98.71%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	90,411,698	2,952,597	3.27%	2.58%	3.95%	87,459,101	96.73%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	90,411,698	2,952,597	3.27%	2.58%	3.95%	87,459,101	96.73%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	90,411,698	2,952,597	3.27%	2.58%	3.95%	87,459,101	96.73%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	91,083,006	4,001,349	4.39%	3.07%	5.71%	87,081,657	95.61%
Age at screening	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
10-19	538,199	0	0.00%	0.00%	0.00%	538,199	100.00%
20-29	12,110,825	359,906	2.97%	0.42%	5.52%	11,750,919	97.03%
30-39	17,822,488	445,312	2.50%	0.61%	4.39%	17,377,176	97.50%
40-49	20,941,149	618,108	2.95%	0.44%	5.46%	20,323,041	97.05%
50-59	17,868,567	761,563	4.26%	1.23%	7.29%	17,107,004	95.74%
60-64	5,666,205	183,436	3.24%	0.50%	5.97%	5,482,769	96.76%
65-69	5,762,216	191,846	3.33%	(0.25%)	6.91%	5,570,369	96.67%
70-74	3,988,818	616,203	15.45%	8.29%	22.61%	3,372,615	84.55%
75-79	3,030,438	344,309	11.36%	2.76%	19.97%	2,686,129	88.64%
80-84	2,000,307	324,332	16.21%	8.38%	24.04%	1,675,976	83.79%
85+	1,353,795	156,335	11.55%	1.07%	22.03%	1,197,460	88.45%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,316,409	222,797	3.05%	0.78%	5.31%	7,093,612	96.95%
Other Hispanic	2,977,019	174,353	5.86%	(0.59%)	12.30%	2,802,665	94.14%
Non-Hispanic White	66,896,218	3,073,126	4.59%	2.89%	6.30%	63,823,092	95.41%
Non-Hispanic Black	9,127,400	441,952	4.84%	2.91%	6.78%	8,685,447	95.16%
Other Race - Including Multi-Racial	4,765,962	89,120	1.87%	(2.28%)	6.02%	4,676,841	98.13%
Education	-	-	-	-	-	-	-
Missing	1,796,573	42,500	2.37%	(2.88%)	7.62%	1,754,073	97.63%
Less Than 9th Grade	6,534,412	421,518	6.45%	2.61%	10.29%	6,112,894	93.55%
9-11th Grade	9,052,486	302,972	3.35%	1.22%	5.48%	8,749,514	96.65%
High School Grad/GED or Equivalent	73,351,938	3,234,359	4.41%	2.97%	5.85%	70,117,578	95.59%
Refused	*	*	*	*	*	*	*
Don't Know	335,074	0	0.00%	0.00%	0.00%	335,074	100.00%
Marital status	-	-	-	-	-	-	-
Missing	1,019,622	6,188	0.61%	(0.68%)	1.89%	1,013,434	99.39%
Married	59,766,248	2,509,326	4.20%	2.78%	5.62%	57,256,922	95.80%
Widowed	3,574,691	318,198	8.90%	0.39%	17.42%	3,256,493	91.10%
Divorced	7,657,708	352,289	4.60%	0.25%	8.95%	7,305,419	95.40%
Separated	1,505,477	81,800	5.43%	(0.09%)	10.96%	1,423,677	94.57%
Never Married	11,206,160	575,185	5.13%	2.02%	8.24%	10,630,975	94.87%
Living with Partner	5,961,389	158,363	2.66%	(0.32%)	5.63%	5,803,025	97.34%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	4,481,624	176,165	3.93%	1.50%	6.36%	4,305,459	96.07%
Poverty income ratio less than 1	9,324,225	263,321	2.82%	1.34%	4.31%	9,060,905	97.18%
Poverty income ratio 1 or greater	77,277,157	3,561,863	4.61%	3.07%	6.15%	73,715,294	95.39%
Body mass index	-	-	-	-	-	-	-
Missing	960,921	206,289	21.47%	(0.07%)	43.01%	754,633	78.53%
Normal: less than 25	27,566,205	1,194,639	4.33%	2.19%	6.47%	26,371,566	95.67%
Overweight: 25.0 - 29.9	34,644,037	1,317,812	3.80%	1.83%	5.78%	33,326,225	96.20%
Obese: 30 or higher	27,911,843	1,282,609	4.60%	3.36%	5.83%	26,629,234	95.40%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2003 - 2004

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	91,083,006	4,001,349	4.39%	3.07%	5.71%	87,081,657	95.61%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,412,745	53,255	2.21%	(1.24%)	5.65%	2,359,490	97.79%
Glycohemoglobin < 5.7%	69,767,673	2,645,183	3.79%	2.59%	4.99%	67,122,490	96.21%
Glycohemoglobin 5.7% - 6.4%	12,802,604	852,392	6.66%	3.09%	10.23%	11,950,212	93.34%
Glycohemoglobin 6.5% or higher	6,099,984	450,519	7.39%	1.98%	12.79%	5,649,465	92.61%
Diabetes	-	-	-	-	-	-	-
Yes	7,009,428	487,264	6.95%	2.55%	11.35%	6,522,164	93.05%
No	84,073,578	3,514,085	4.18%	2.96%	5.40%	80,559,493	95.82%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,688,389	2,223,533	4.22%	2.52%	5.92%	50,464,856	95.78%
No	38,356,405	1,765,985	4.60%	2.29%	6.92%	36,590,420	95.40%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	789,967	0	0.00%	0.00%	0.00%	789,967	100.00%
Yes	27,630,586	1,692,258	6.12%	3.74%	8.51%	25,938,329	93.88%
No	62,556,780	2,309,092	3.69%	2.56%	4.82%	60,247,689	96.31%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	9,336,477	1,035,708	11.09%	6.34%	15.84%	8,300,769	88.91%
Mobility limitations	7,921,738	423,581	5.35%	1.36%	9.33%	7,498,157	94.65%
No mobility limitations	19,554,028	962,764	4.92%	3.16%	6.68%	18,591,264	95.08%
N/A - Respondent not asked mobility limitation questions	54,270,763	1,579,297	2.91%	1.10%	4.72%	52,691,467	97.09%
Sleep disorder	-	-	-	-	-	-	-
N/A for these years of data	91,083,006	4,001,349	4.39%	3.07%	5.71%	87,081,657	95.61%
Bowel leakage - gas	-	-	-	-	-	-	-
N/A for these years of data	91,083,006	4,001,349	4.39%	3.07%	5.71%	87,081,657	95.61%
Bowel leakage - stool	-	-	-	-	-	-	-
N/A for these years of data	91,083,006	4,001,349	4.39%	3.07%	5.71%	87,081,657	95.61%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,607,576	3,675,867	3.93%	2.62%	5.23%	89,931,709	96.07%
Age at screening	-	-	-	-	-	-	-
10-19	384,918	0	0.00%	0.00%	0.00%	384,918	100.00%
20-29	14,540,015	279,671	1.92%	(0.37%)	4.21%	14,260,345	98.08%
30-39	17,503,545	252,156	1.44%	0.12%	2.77%	17,251,389	98.56%
40-49	21,809,592	850,679	3.90%	1.72%	6.08%	20,958,912	96.10%
50-59	18,142,193	443,000	2.44%	0.40%	4.48%	17,699,193	97.56%
60-64	6,373,461	341,489	5.36%	2.60%	8.12%	6,031,973	94.64%
65-69	5,297,573	383,111	7.23%	4.31%	10.15%	4,914,461	92.77%
70-74	3,970,856	362,711	9.13%	4.16%	14.11%	3,608,146	90.87%
75-79	2,536,289	403,615	15.91%	6.85%	24.98%	2,132,674	84.09%
80-84	2,002,541	238,686	11.92%	4.53%	19.31%	1,763,855	88.08%
85+	1,046,592	120,749	11.54%	1.94%	21.13%	925,843	88.46%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	7,979,822	157,047	1.97%	(0.33%)	4.27%	7,822,774	98.03%
Other Hispanic	2,862,199	58,017	2.03%	(2.21%)	6.27%	2,804,182	97.97%
Non-Hispanic White	69,050,169	3,207,870	4.65%	3.10%	6.20%	65,842,299	95.35%
Non-Hispanic Black	9,553,122	252,933	2.65%	0.90%	4.40%	9,300,189	97.35%
Other Race - Including Multi-Racial	4,162,265	0	0.00%	0.00%	0.00%	4,162,265	100.00%
Education	-	-	-	-	-	-	-
Missing	1,125,917	0	0.00%	0.00%	0.00%	1,125,917	100.00%
Less Than 9th Grade	6,333,269	192,180	3.03%	0.07%	5.99%	6,141,089	96.97%
9-11th Grade	10,260,776	361,303	3.52%	1.42%	5.62%	9,899,474	96.48%
High School Grad/GED or Equivalent	75,722,078	3,113,414	4.11%	2.64%	5.58%	72,608,664	95.89%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,042,842	8,970	0.86%	(1.00%)	2.72%	1,033,871	99.14%
Married	60,157,438	2,669,633	4.44%	2.78%	6.10%	57,487,805	95.56%
Widowed	2,879,447	223,802	7.77%	0.23%	15.32%	2,655,645	92.23%
Divorced	9,188,198	309,873	3.37%	0.55%	6.19%	8,878,325	96.63%
Separated	2,712,975	90,524	3.34%	(0.59%)	7.26%	2,622,451	96.66%
Never Married	9,472,270	203,635	2.15%	(0.79%)	5.09%	9,268,635	97.85%
Living with Partner	8,115,581	130,604	1.61%	(0.92%)	4.14%	7,984,977	98.39%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	2,977,906	204,300	6.86%	0.19%	13.54%	2,773,605	93.14%
Poverty income ratio less than 1	9,079,370	334,660	3.69%	1.20%	6.17%	8,744,711	96.31%
Poverty income ratio 1 or greater	81,550,300	3,136,908	3.85%	2.56%	5.13%	78,413,393	96.15%
Body mass index	-	-	-	-	-	-	-
Missing	1,009,735	53,177	5.27%	(2.20%)	12.74%	956,557	94.73%
Normal: less than 25	25,820,546	692,850	2.68%	0.83%	4.54%	25,127,695	97.32%
Overweight: 25.0 - 29.9	35,878,979	1,669,382	4.65%	2.57%	6.74%	34,209,597	95.35%
Obese: 30 or higher	30,898,317	1,260,458	4.08%	1.64%	6.51%	29,637,860	95.92%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,564,155	44,480	1.25%	(1.44%)	3.94%	3,519,675	98.75%
Glycohemoglobin < 5.7%	70,219,466	2,677,627	3.81%	2.27%	5.35%	67,541,839	96.19%
Glycohemoglobin 5.7% - 6.4%	14,222,346	711,748	5.00%	3.29%	6.72%	13,510,598	95.00%
Glycohemoglobin 6.5% or higher	5,601,610	242,013	4.32%	1.29%	7.35%	5,359,597	95.68%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2005 - 2006

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	93,607,576	3,675,867	3.93%	2.62%	5.23%	89,931,709	96.07%
Diabetes	-	-	-	-	-	-	-
Yes	6,979,396	350,895	5.03%	0.95%	9.11%	6,628,501	94.97%
No	86,628,181	3,324,972	3.84%	2.45%	5.23%	83,303,208	96.16%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,967,879	2,663,334	5.03%	3.02%	7.04%	50,304,545	94.97%
No	40,639,698	1,012,533	2.49%	1.48%	3.50%	39,627,164	97.51%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	26,812,046	1,675,760	6.25%	4.03%	8.47%	25,136,285	93.75%
No	66,617,068	1,943,528	2.92%	1.16%	4.67%	64,673,539	97.08%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	6,866,793	596,541	8.69%	3.58%	13.80%	6,270,252	91.31%
Mobility limitations	5,930,657	425,613	7.18%	4.17%	10.18%	5,505,043	92.82%
No mobility limitations	20,256,288	1,337,177	6.60%	4.83%	8.37%	18,919,111	93.40%
N/A - Respondent not asked mobility limitation questions	60,553,839	1,316,537	2.17%	0.77%	3.58%	59,237,302	97.83%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,936,022	458,050	5.77%	1.10%	10.44%	7,477,972	94.23%
No	85,555,452	3,217,818	3.76%	2.27%	5.25%	82,337,634	96.24%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	372,256	33,688	9.05%	(2.23%)	20.33%	338,567	90.95%
2 or more times a day	13,735,930	799,653	5.82%	1.57%	10.07%	12,936,277	94.18%
Once a day	8,509,610	218,128	2.56%	(0.23%)	5.36%	8,291,482	97.44%
2 or more time a week	5,956,076	452,832	7.60%	1.23%	13.98%	5,503,244	92.40%
Once a week	5,314,628	353,190	6.65%	0.70%	12.60%	4,961,438	93.35%
1-3 times a month	9,361,331	700,822	7.49%	3.62%	11.35%	8,660,510	92.51%
Never	50,064,929	1,117,555	2.23%	1.11%	3.36%	48,947,374	97.77%
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	372,256	33,688	9.05%	(2.23%)	20.33%	338,567	90.95%
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	749,457	227,289	30.33%	4.35%	56.30%	522,168	69.67%
Never	92,089,049	3,365,300	3.65%	2.27%	5.03%	88,723,748	96.35%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,900,379	3,989,857	4.20%	3.10%	5.31%	90,910,522	95.80%
Age at screening	-	-	-	-	-	-	-
10-19	581,889	0	0.00%	0.00%	0.00%	581,889	100.00%
20-29	14,324,402	70,913	0.50%	(0.16%)	1.15%	14,253,489	99.50%
30-39	17,120,661	89,941	0.53%	(0.18%)	1.23%	17,030,720	99.47%
40-49	21,351,091	813,283	3.81%	0.97%	6.65%	20,537,808	96.19%
50-59	18,864,707	981,371	5.20%	2.03%	8.38%	17,883,336	94.80%
60-64	7,462,217	327,050	4.38%	(0.29%)	9.05%	7,135,167	95.62%
65-69	5,293,210	428,262	8.09%	4.41%	11.77%	4,864,949	91.91%
70-74	3,760,917	392,888	10.45%	4.98%	15.91%	3,368,029	89.55%
75-79	2,921,173	439,686	15.05%	3.51%	26.59%	2,481,487	84.95%
80+	3,220,111	446,463	13.86%	8.54%	19.19%	2,773,648	86.14%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,009,819	221,844	2.46%	0.38%	4.55%	8,787,974	97.54%
Other Hispanic	4,534,772	101,012	2.23%	0.89%	3.57%	4,433,761	97.77%
Non-Hispanic White	65,865,926	2,895,648	4.40%	3.10%	5.70%	62,970,278	95.60%
Non-Hispanic Black	9,812,309	443,564	4.52%	2.43%	6.61%	9,368,746	95.48%
Other Race - Including Multi-Racial	5,677,553	327,789	5.77%	(0.76%)	12.30%	5,349,763	94.23%
Education	-	-	-	-	-	-	-
Missing	1,275,540	9,066	0.71%	(0.81%)	2.23%	1,266,475	99.29%
Less Than 9th Grade	6,679,302	453,466	6.79%	2.76%	10.82%	6,225,836	93.21%
9-11th Grade	11,881,477	369,185	3.11%	1.51%	4.71%	11,512,292	96.89%
High School Grad/GED or Equivalent	74,927,632	3,158,139	4.21%	3.00%	5.43%	71,769,493	95.79%
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	1,282,066	0	0.00%	0.00%	0.00%	1,282,066	100.00%
Married	60,660,936	2,495,814	4.11%	2.52%	5.71%	58,165,123	95.89%
Widowed	3,201,326	249,784	7.80%	0.33%	15.28%	2,951,542	92.20%
Divorced	8,397,982	585,044	6.97%	2.44%	11.50%	7,812,937	93.03%
Separated	2,134,533	96,709	4.53%	(0.76%)	9.82%	2,037,824	95.47%
Never Married	12,388,270	174,055	1.40%	0.54%	2.27%	12,214,215	98.60%
Living with Partner	6,649,894	388,451	5.84%	0.92%	10.77%	6,261,443	94.16%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,832,632	390,439	5.71%	3.04%	8.39%	6,442,192	94.29%
Poverty income ratio less than 1	11,426,636	252,425	2.21%	1.20%	3.22%	11,174,211	97.79%
Poverty income ratio 1 or greater	76,641,111	3,346,992	4.37%	3.12%	5.61%	73,294,118	95.63%
Body mass index	-	-	-	-	-	-	-
Missing	778,736	36,914	4.74%	(3.18%)	12.66%	741,822	95.26%
Normal: less than 25	26,748,202	1,128,857	4.22%	2.39%	6.05%	25,619,345	95.78%
Overweight: 25.0 - 29.9	36,904,215	1,171,399	3.17%	2.26%	4.09%	35,732,816	96.83%
Obese: 30 or higher	30,469,225	1,652,687	5.42%	3.03%	7.82%	28,816,539	94.58%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	3,558,454	39,301	1.10%	(0.27%)	2.48%	3,519,153	98.90%
Glycohemoglobin < 5.7%	62,899,045	1,753,796	2.79%	1.63%	3.95%	61,145,249	97.21%
Glycohemoglobin 5.7% - 6.4%	21,160,974	1,557,264	7.36%	4.60%	10.12%	19,603,710	92.64%
Glycohemoglobin 6.5% or higher	7,281,906	639,496	8.78%	4.02%	13.54%	6,642,410	91.22%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2007 - 2008

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	94,900,379	3,989,857	4.20%	3.10%	5.31%	90,910,522	95.80%
Diabetes	-	-	-	-	-	-	-
Yes	8,462,529	585,592	6.92%	3.57%	10.27%	7,876,937	93.08%
No	86,437,849	3,404,264	3.94%	2.81%	5.07%	83,033,585	96.06%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	52,022,370	2,506,294	4.82%	3.06%	6.58%	49,516,076	95.18%
No	42,810,155	1,483,563	3.47%	2.11%	4.82%	41,326,592	96.53%
Don't know	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,186,914	2,454,018	8.71%	5.44%	11.97%	25,732,896	91.29%
No	66,610,852	1,535,839	2.31%	1.67%	2.94%	65,075,014	97.69%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	8,937,010	755,983	8.46%	5.61%	11.31%	8,181,027	91.54%
Mobility limitations	6,878,098	706,352	10.27%	6.67%	13.87%	6,171,746	89.73%
No mobility limitations	18,940,300	958,623	5.06%	3.02%	7.11%	17,981,677	94.94%
Refused/Don't know	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	60,107,142	1,568,898	2.61%	1.36%	3.86%	58,538,243	97.39%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,808,478	653,360	8.37%	2.74%	13.99%	7,155,119	91.63%
No	86,879,737	3,336,497	3.84%	2.73%	4.95%	83,543,240	96.16%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	574,280	49,251	8.58%	(2.22%)	19.37%	525,029	91.42%
2 or more times a day	10,308,368	653,142	6.34%	2.38%	10.30%	9,655,226	93.66%
Once a day	8,264,954	334,691	4.05%	1.69%	6.41%	7,930,263	95.95%
2 or more time a week	5,908,062	358,953	6.08%	2.15%	10.00%	5,549,109	93.92%
Once a week	4,371,150	77,192	1.77%	(1.03%)	4.56%	4,293,958	98.23%
1-3 times a month	8,302,501	853,343	10.28%	4.19%	16.37%	7,449,157	89.72%
Never	56,789,894	1,663,284	2.93%	1.93%	3.93%	55,126,610	97.07%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	574,280	49,251	8.58%	(2.22%)	19.37%	525,029	91.42%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	1,037,565	150,158	14.47%	0.92%	28.02%	887,407	85.53%
Never	92,552,323	3,790,447	4.10%	3.01%	5.18%	88,761,876	95.90%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,526,374	4,165,544	4.36%	3.28%	5.45%	91,360,830	95.64%
Age at screening	-	-	-	-	-	-	-
10-19	297,636	0	0.00%	0.00%	0.00%	297,636	100.00%
20-29	12,944,492	255,494	1.97%	(0.23%)	4.18%	12,688,998	98.03%
30-39	16,576,959	216,356	1.31%	0.35%	2.26%	16,360,603	98.69%
40-49	21,324,506	923,292	4.33%	1.60%	7.06%	20,401,214	95.67%
50-59	21,456,557	756,720	3.53%	1.47%	5.58%	20,699,837	96.47%
60-64	7,051,206	532,739	7.56%	2.01%	13.10%	6,518,467	92.44%
65-69	6,200,612	482,971	7.79%	2.43%	13.15%	5,717,641	92.21%
70-74	3,723,264	253,350	6.80%	1.51%	12.10%	3,469,915	93.20%
75-79	2,936,105	265,394	9.04%	2.58%	15.49%	2,670,711	90.96%
80+	3,015,038	479,229	15.89%	9.79%	22.00%	2,535,809	84.11%
Race/ethnicity	-	-	-	-	-	-	-
Mexican American	9,079,208	133,891	1.47%	0.30%	2.65%	8,945,317	98.53%
Other Hispanic	4,899,222	92,668	1.89%	0.00%	3.78%	4,806,554	98.11%
Non-Hispanic White	66,059,416	3,293,559	4.99%	3.48%	6.49%	62,765,857	95.01%
Non-Hispanic Black	9,655,413	513,411	5.32%	3.44%	7.19%	9,142,002	94.68%
Other Race - Including Multi-Racial	5,833,115	132,015	2.26%	(1.48%)	6.00%	5,701,100	97.74%
Education	-	-	-	-	-	-	-
Missing	1,359,443	109,205	8.03%	(2.82%)	18.89%	1,250,238	91.97%
Less Than 9th Grade	6,477,166	167,456	2.59%	(0.48%)	5.65%	6,309,710	97.41%
9-11th Grade	10,241,048	514,877	5.03%	1.62%	8.44%	9,726,171	94.97%
High School Grad/GED or Equivalent	77,302,256	3,374,006	4.36%	3.24%	5.49%	73,928,250	95.64%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Marital status	-	-	-	-	-	-	-
Missing	313,184	15,549	4.96%	(5.58%)	15.51%	297,636	95.04%
Married	61,034,010	3,022,250	4.95%	3.54%	6.36%	58,011,760	95.05%
Widowed	3,241,681	265,282	8.18%	3.70%	12.67%	2,976,400	91.82%
Divorced	8,786,406	198,398	2.26%	0.22%	4.30%	8,588,008	97.74%
Separated	2,184,790	52,985	2.43%	(1.18%)	6.03%	2,131,806	97.57%
Never Married	12,431,390	347,896	2.80%	0.82%	4.78%	12,083,494	97.20%
Living with Partner	7,478,020	247,614	3.31%	0.06%	6.56%	7,230,406	96.69%
Refused	*	*	*	*	*	*	*
Don't Know	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-
Missing	6,719,530	248,215	3.69%	0.85%	6.54%	6,471,315	96.31%
Poverty income ratio less than 1	10,991,125	345,047	3.14%	0.80%	5.48%	10,646,078	96.86%
Poverty income ratio 1 or greater	77,815,720	3,572,283	4.59%	3.36%	5.83%	74,243,437	95.41%
Body mass index	-	-	-	-	-	-	-
Missing	914,576	78,878	8.62%	(5.45%)	22.70%	835,698	91.38%
Normal: less than 25	24,656,831	1,292,395	5.24%	2.98%	7.50%	23,364,436	94.76%
Overweight: 25.0 - 29.9	36,168,956	1,161,381	3.21%	1.90%	4.53%	35,007,575	96.79%
Obese: 30 or higher	33,786,012	1,632,890	4.83%	2.88%	6.79%	32,153,122	95.17%
Glycohemoglobin	-	-	-	-	-	-	-
Missing	2,643,943	48,089	1.82%	(1.06%)	4.69%	2,595,854	98.18%
Glycohemoglobin < 5.7%	61,620,636	2,228,781	3.62%	2.39%	4.85%	59,391,855	96.38%
Glycohemoglobin 5.7% - 6.4%	23,018,912	1,372,628	5.96%	3.92%	8.01%	21,646,283	94.04%
Glycohemoglobin 6.5% or higher	8,242,884	516,045	6.26%	3.27%	9.25%	7,726,839	93.74%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M05 - Prevalence of insensate urine loss among men, 2009 - 2010

	Total (N)	Yes (N)	Yes (%)	Lower 95% CI	Upper 95% CI	No (N)	No (%)
Total	95,526,374	4,165,544	4.36%	3.28%	5.45%	91,360,830	95.64%
Diabetes	-	-	-	-	-	-	-
Yes	8,780,629	640,241	7.29%	4.48%	10.10%	8,140,388	92.71%
No	86,745,745	3,525,303	4.06%	2.94%	5.18%	83,220,442	95.94%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	47,419,960	2,237,513	4.72%	3.32%	6.12%	45,182,448	95.28%
No	48,106,414	1,928,031	4.01%	2.36%	5.66%	46,178,382	95.99%
High blood pressure	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	28,837,492	1,814,026	6.29%	4.32%	8.26%	27,023,466	93.71%
No	66,577,565	2,351,518	3.53%	2.46%	4.60%	64,226,047	96.47%
Don't know	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-
Missing	7,905,122	697,283	8.82%	5.28%	12.36%	7,207,838	91.18%
Mobility limitations	6,892,982	719,633	10.44%	6.34%	14.54%	6,173,348	89.56%
No mobility limitations	19,994,371	1,235,244	6.18%	3.71%	8.65%	18,759,127	93.82%
N/A - Respondent not asked mobility limitation questions	60,733,899	1,513,383	2.49%	1.26%	3.72%	59,220,516	97.51%
Sleep disorder	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*
Yes	7,676,778	785,278	10.23%	3.83%	16.63%	6,891,500	89.77%
No	87,646,247	3,380,266	3.86%	2.82%	4.90%	84,265,981	96.14%
Don't know	*	*	*	*	*	*	*
Bowel leakage - gas	-	-	-	-	-	-	-
Missing	459,339	91,102	19.83%	(5.25%)	44.92%	368,238	80.17%
2 or more times a day	9,148,208	485,649	5.31%	2.11%	8.51%	8,662,559	94.69%
Once a day	8,770,661	547,595	6.24%	2.65%	9.84%	8,223,066	93.76%
2 or more time a week	6,773,718	456,027	6.73%	0.38%	13.09%	6,317,691	93.27%
Once a week	4,927,442	277,244	5.63%	0.30%	10.95%	4,650,198	94.37%
1-3 times a month	10,278,875	795,351	7.74%	2.45%	13.03%	9,483,524	92.26%
Never	55,104,454	1,512,576	2.74%	1.67%	3.82%	53,591,878	97.26%
Refused	*	*	*	*	*	*	*
Don't know	*	*	*	*	*	*	*
Bowel leakage - stool	-	-	-	-	-	-	-
Missing	459,339	91,102	19.83%	(5.25%)	44.92%	368,238	80.17%
2 or more times a day	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*
1-3 times a month	844,643	174,180	20.62%	1.62%	39.62%	670,463	79.38%
Never	93,686,486	3,887,069	4.15%	3.09%	5.21%	89,799,417	95.85%
Don't know	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).