

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
10-19	*	*	*	*	*	*	*	*	*	*	*	*	*
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	*	*	*	*	*	*	*	*	*	*	*	*	*
50-59	683,981	17,605	2.57%	431,622	63.10%	34,787	5.09%	199,968	29.24%	0	0.00%	0	0.00%
60-64	249,754	43,082	17.25%	26,370	10.56%	48,359	19.36%	131,943	52.83%	0	0.00%	0	0.00%
65-69	305,063	72,094	23.63%	143,694	47.10%	45,976	15.07%	43,298	14.19%	0	0.00%	0	0.00%
70-74	437,470	140,939	32.22%	64,757	14.80%	131,420	30.04%	100,354	22.94%	0	0.00%	0	0.00%
75-79	348,796	147,129	42.18%	68,873	19.75%	94,494	27.09%	38,301	10.98%	0	0.00%	0	0.00%
80-84	247,548	39,667	16.02%	91,956	37.15%	63,778	25.76%	52,148	21.07%	0	0.00%	0	0.00%
85+	254,173	118,409	46.59%	73,630	28.97%	38,681	15.22%	23,453	9.23%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	157,894	35,995	22.80%	37,320	23.64%	14,394	9.12%	70,184	44.45%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	2,346,031	520,285	22.18%	683,095	29.12%	473,841	20.20%	668,809	28.51%	0	0.00%	0	0.00%
Non-Hispanic Black	245,085	62,666	25.57%	101,756	41.52%	18,653	7.61%	62,010	25.30%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	407,756	213,807	52.44%	118,241	29.00%	38,858	9.53%	36,850	9.04%	0	0.00%	0	0.00%
9-11th Grade	376,336	97,449	25.89%	85,280	22.66%	79,698	21.18%	113,909	30.27%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	2,100,490	300,553	14.31%	686,273	32.67%	388,333	18.49%	725,331	34.53%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,095,650	461,199	22.01%	751,956	35.88%	311,876	14.88%	570,619	27.23%	0	0.00%	0	0.00%
Widowed	327,563	87,376	26.67%	54,286	16.57%	125,225	38.23%	60,676	18.52%	0	0.00%	0	0.00%
Divorced	317,231	6,473	2.04%	47,420	14.95%	62,423	19.68%	200,915	63.33%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	169,351	43,161	25.49%	74,947	44.26%	7,364	4.35%	43,879	25.91%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	526,018	83,145	15.81%	121,041	23.01%	98,397	18.71%	223,434	42.48%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	2,234,342	531,724	23.80%	797,521	35.69%	408,491	18.28%	496,605	22.23%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	475,220	208,760	43.93%	141,665	29.81%	63,606	13.38%	61,190	12.88%	0	0.00%	0	0.00%
Normal: less than 25	581,300	87,013	14.97%	179,964	30.96%	79,576	13.69%	234,747	40.38%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,050,439	127,993	12.18%	300,635	28.62%	304,059	28.95%	317,752	30.25%	0	0.00%	0	0.00%
Obese: 30 or higher	845,638	217,245	25.69%	306,346	36.23%	59,648	7.05%	262,400	31.03%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	1,920,404	350,649	18.26%	566,193	29.48%	238,579	12.42%	764,983	39.83%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	580,748	142,555	24.55%	220,731	38.01%	146,432	25.21%	71,031	12.23%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	365,304	86,615	23.71%	128,025	35.05%	110,589	30.27%	40,075	10.97%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	426,056	102,378	24.03%	186,861	43.86%	98,839	23.20%	37,978	8.91%	0	0.00%	0	0.00%
No	2,526,542	538,634	21.32%	741,748	29.36%	408,049	16.15%	838,111	33.17%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2001 - 2002

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,001,836	393,655	19.66%	693,742	34.66%	341,318	17.05%	573,119	28.63%	0	0.00%	0	0.00%
No	950,762	247,356	26.02%	234,866	24.70%	165,570	17.41%	302,970	31.87%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,336,170	391,479	29.30%	349,552	26.16%	335,678	25.12%	259,461	19.42%	0	0.00%	0	0.00%
No	1,616,427	249,532	15.44%	579,056	35.82%	171,211	10.59%	616,628	38.15%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	627,786	209,825	33.42%	103,419	16.47%	109,979	17.52%	204,562	32.58%	0	0.00%	0	0.00%
Mobility limitations	587,782	193,717	32.96%	176,300	29.99%	119,749	20.37%	98,016	16.68%	0	0.00%	0	0.00%
No mobility limitations	955,378	203,772	21.33%	298,164	31.21%	227,765	23.84%	225,676	23.62%	0	0.00%	0	0.00%
Refused/Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
N/A - Respondent not asked mobility limitation questions	767,037	19,082	2.49%	350,725	45.72%	49,394	6.44%	347,835	45.35%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	2,952,597	641,011	21.71%	928,608	31.45%	506,888	17.17%	876,089	29.67%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	445,312	97,823	21.97%	88,777	19.94%	24,738	5.56%	233,974	52.54%	0	0.00%	0	0.00%
40-49	618,108	34,621	5.60%	151,143	24.45%	47,203	7.64%	385,142	62.31%	0	0.00%	0	0.00%
50-59	761,563	119,135	15.64%	122,733	16.12%	190,526	25.02%	329,169	43.22%	0	0.00%	0	0.00%
60-64	183,436	112,851	61.52%	28,707	15.65%	41,878	22.83%	0	0.00%	0	0.00%	0	0.00%
65-69	*	*	*	*	*	*	*	*	*	*	*	*	*
70-74	616,203	134,521	21.83%	250,638	40.67%	28,725	4.66%	202,319	32.83%	0	0.00%	0	0.00%
75-79	344,309	128,947	37.45%	160,647	46.66%	2,145	0.62%	52,570	15.27%	0	0.00%	0	0.00%
80-84	324,332	197,159	60.79%	16,867	5.20%	64,675	19.94%	45,631	14.07%	0	0.00%	0	0.00%
85+	*	*	*	*	*	*	*	*	*	*	*	*	*
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	222,797	49,277	22.12%	48,130	21.60%	40,441	18.15%	84,948	38.13%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	3,073,126	815,481	26.54%	774,260	25.19%	473,989	15.42%	1,009,396	32.85%	0	0.00%	0	0.00%
Non-Hispanic Black	441,952	99,911	22.61%	140,156	31.71%	108,564	24.56%	93,322	21.12%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	421,518	188,934	44.82%	125,083	29.67%	45,521	10.80%	61,979	14.70%	0	0.00%	0	0.00%
9-11th Grade	302,972	178,305	58.85%	43,290	14.29%	42,966	14.18%	38,411	12.68%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,234,359	624,439	19.31%	775,964	23.99%	534,507	16.53%	1,299,450	40.18%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,509,326	592,257	23.60%	644,710	25.69%	300,752	11.99%	971,608	38.72%	0	0.00%	0	0.00%
Widowed	318,198	211,897	66.59%	55,217	17.35%	14,592	4.59%	36,492	11.47%	0	0.00%	0	0.00%
Divorced	352,289	119,034	33.79%	58,393	16.58%	87,385	24.80%	87,476	24.83%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	575,185	59,371	10.32%	171,856	29.88%	119,832	20.83%	224,125	38.97%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	263,321	91,578	34.78%	33,527	12.73%	106,837	40.57%	31,377	11.92%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	3,561,863	836,745	23.49%	888,094	24.93%	492,959	13.84%	1,344,065	37.73%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,194,639	333,608	27.93%	176,597	14.78%	213,374	17.86%	471,060	39.43%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,317,812	247,884	18.81%	333,023	25.27%	275,922	20.94%	460,984	34.98%	0	0.00%	0	0.00%
Obese: 30 or higher	1,282,609	327,733	25.55%	366,096	28.54%	133,698	10.42%	455,082	35.48%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	2,645,183	692,093	26.16%	687,181	25.98%	374,550	14.16%	891,359	33.70%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	852,392	232,395	27.26%	108,575	12.74%	198,363	23.27%	313,059	36.73%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	450,519	91,482	20.31%	166,789	37.02%	26,883	5.97%	165,365	36.71%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	487,264	176,786	36.28%	80,112	16.44%	66,946	13.74%	163,420	33.54%	0	0.00%	0	0.00%
No	3,514,085	839,184	23.88%	882,434	25.11%	556,047	15.82%	1,236,420	35.18%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a - Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2003 - 2004

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,223,533	819,950	36.88%	560,130	25.19%	359,073	16.15%	484,380	21.78%	0	0.00%	0	0.00%
No	1,765,985	184,189	10.43%	402,416	22.79%	263,921	14.94%	915,460	51.84%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,692,258	523,607	30.94%	524,170	30.97%	302,863	17.90%	341,617	20.19%	0	0.00%	0	0.00%
No	2,309,092	492,362	21.32%	438,376	18.98%	320,131	13.86%	1,058,223	45.83%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	1,035,708	486,554	46.98%	158,122	15.27%	136,008	13.13%	255,023	24.62%	0	0.00%	0	0.00%
Mobility limitations	423,581	250,408	59.12%	85,751	20.24%	59,892	14.14%	27,530	6.50%	0	0.00%	0	0.00%
No mobility limitations	962,764	229,711	23.86%	364,518	37.86%	164,472	17.08%	204,062	21.20%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,579,297	49,296	3.12%	354,154	22.42%	262,622	16.63%	913,224	57.82%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
N/A for these years of data	4,001,349	1,015,969	25.39%	962,546	24.06%	622,994	15.57%	1,399,840	34.98%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	850,679	130,902	15.39%	382,513	44.97%	0	0.00%	337,264	39.65%	0	0.00%	0	0.00%
50-59	*	*	*	*	*	*	*	*	*	*	*	*	*
60-64	341,489	4,008	1.17%	27,837	8.15%	101,524	29.73%	208,120	60.94%	0	0.00%	0	0.00%
65-69	383,111	70,999	18.53%	0	0.00%	157,844	41.20%	154,269	40.27%	0	0.00%	0	0.00%
70-74	362,711	21,351	5.89%	44,954	12.39%	77,418	21.34%	218,988	60.38%	0	0.00%	0	0.00%
75-79	*	*	*	*	*	*	*	*	*	*	*	*	*
80-84	238,686	40,804	17.10%	44,884	18.80%	49,147	20.59%	103,850	43.51%	0	0.00%	0	0.00%
85+	*	*	*	*	*	*	*	*	*	*	*	*	*
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	157,047	4,008	2.55%	18,866	12.01%	108,100	68.83%	26,073	16.60%	0	0.00%	0	0.00%
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	3,207,870	462,567	14.42%	691,471	21.56%	577,222	17.99%	1,476,610	46.03%	0	0.00%	0	0.00%
Non-Hispanic Black	252,933	0	0.00%	8,970	3.55%	108,185	42.77%	135,777	53.68%	0	0.00%	0	0.00%
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Less Than 9th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*
9-11th Grade	361,303	4,008	1.11%	44,146	12.22%	62,934	17.42%	250,214	69.25%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,113,414	450,002	14.45%	606,532	19.48%	706,630	22.70%	1,350,250	43.37%	0	0.00%	0	0.00%
Don't Know	*	*	*	*	*	*	*	*	*	*	*	*	*

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	2,669,633	262,667	9.84%	522,178	19.56%	441,654	16.54%	1,443,134	54.06%	0	0.00%	0	0.00%
Widowed	*	*	*	*	*	*	*	*	*	*	*	*	*
Divorced	*	*	*	*	*	*	*	*	*	*	*	*	*
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	*	*	*	*	*	*	*	*	*	*	*	*	*
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio less than 1	334,660	0	0.00%	74,602	22.29%	87,808	26.24%	172,249	51.47%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	3,136,908	466,575	14.87%	535,150	17.06%	631,677	20.14%	1,503,505	47.93%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	692,850	89,721	12.95%	94,756	13.68%	181,461	26.19%	326,913	47.18%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,669,382	173,241	10.38%	481,749	28.86%	334,775	20.05%	679,617	40.71%	0	0.00%	0	0.00%
Obese: 30 or higher	1,260,458	203,614	16.15%	123,199	9.77%	256,160	20.32%	677,486	53.75%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	2,677,627	313,686	11.72%	524,649	19.59%	434,110	16.21%	1,405,182	52.48%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	711,748	140,325	19.72%	131,313	18.45%	256,983	36.11%	183,127	25.73%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	242,013	12,565	5.19%	18,866	7.80%	102,414	42.32%	108,167	44.69%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	350,895	148,013	42.18%	0	0.00%	102,787	29.29%	100,095	28.53%	0	0.00%	0	0.00%
No	3,324,972	318,562	9.58%	719,308	21.63%	690,721	20.77%	1,596,382	48.01%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,663,334	454,010	17.05%	503,128	18.89%	489,717	18.39%	1,216,479	45.68%	0	0.00%	0	0.00%
No	1,012,533	12,565	1.24%	216,180	21.35%	303,791	30.00%	479,998	47.41%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,675,760	313,609	18.71%	320,298	19.11%	249,780	14.91%	792,073	47.27%	0	0.00%	0	0.00%
No	1,943,528	152,966	7.87%	399,009	20.53%	487,149	25.07%	904,404	46.53%	0	0.00%	0	0.00%
Don't know	*	*	*	*	*	*	*	*	*	*	*	*	*
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	596,541	231,927	38.88%	153,971	25.81%	97,939	16.42%	112,704	18.89%	0	0.00%	0	0.00%
Mobility limitations	425,613	0	0.00%	100,215	23.55%	85,351	20.05%	240,047	56.40%	0	0.00%	0	0.00%
No mobility limitations	1,337,177	151,103	11.30%	82,609	6.18%	363,049	27.15%	740,417	55.37%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,316,537	83,545	6.35%	382,513	29.05%	247,169	18.77%	603,309	45.83%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	458,050	34,914	7.62%	45,471	9.93%	90,114	19.67%	287,551	62.78%	0	0.00%	0	0.00%
No	3,217,818	431,662	13.41%	673,837	20.94%	703,394	21.86%	1,408,926	43.79%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	799,653	212,356	26.56%	120,089	15.02%	140,776	17.60%	326,432	40.82%	0	0.00%	0	0.00%
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	700,822	4,008	0.57%	194,231	27.71%	139,841	19.95%	362,741	51.76%	0	0.00%	0	0.00%
Never	1,117,555	62,313	5.58%	101,585	9.09%	333,035	29.80%	620,621	55.53%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2005 - 2006

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a day	*	*	*	*	*	*	*	*	*	*	*	*	*
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	3,365,300	404,420	12.02%	708,141	21.04%	745,492	22.15%	1,507,247	44.79%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,989,857	488,975	12.26%	528,399	13.24%	1,183,017	29.65%	1,745,660	43.75%	43,806	1.10%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	813,283	27,326	3.36%	156,332	19.22%	229,004	28.16%	400,621	49.26%	0	0.00%	0	0.00%
50-59	981,371	115,577	11.78%	153,138	15.60%	186,694	19.02%	506,073	51.57%	19,889	2.03%	0	0.00%
60-64	327,050	0	0.00%	7,722	2.36%	174,496	53.35%	144,833	44.28%	0	0.00%	0	0.00%
65-69	428,262	65,101	15.20%	30,159	7.04%	162,332	37.90%	170,669	39.85%	0	0.00%	0	0.00%
70-74	392,888	50,561	12.87%	62,724	15.96%	166,835	42.46%	112,768	28.70%	0	0.00%	0	0.00%
75-79	439,686	119,699	27.22%	48,550	11.04%	58,986	13.42%	212,452	48.32%	0	0.00%	0	0.00%
80+	446,463	110,711	24.80%	69,774	15.63%	146,280	32.76%	95,780	21.45%	23,917	5.36%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	221,844	38,490	17.35%	38,152	17.20%	24,395	11.00%	120,808	54.46%	0	0.00%	0	0.00%
Other Hispanic	101,012	24,128	23.89%	9,034	8.94%	28,613	28.33%	39,236	38.84%	0	0.00%	0	0.00%
Non-Hispanic White	2,895,648	343,829	11.87%	440,188	15.20%	685,245	23.66%	1,402,469	48.43%	23,917	0.83%	0	0.00%
Non-Hispanic Black	443,564	34,700	7.82%	41,025	9.25%	199,342	44.94%	148,607	33.50%	19,889	4.48%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	453,466	109,182	24.08%	98,317	21.68%	98,436	21.71%	147,532	32.53%	0	0.00%	0	0.00%
9-11th Grade	369,185	94,008	25.46%	20,348	5.51%	138,219	37.44%	96,721	26.20%	19,889	5.39%	0	0.00%
High School Grad/GED or Equivalent	3,158,139	285,786	9.05%	409,734	12.97%	937,297	29.68%	1,501,407	47.54%	23,917	0.76%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Married	2,495,814	245,082	9.82%	416,179	16.68%	878,207	35.19%	912,540	36.56%	43,806	1.76%	0	0.00%
Widowed	249,784	61,732	24.71%	0	0.00%	54,865	21.96%	133,187	53.32%	0	0.00%	0	0.00%
Divorced	585,044	143,693	24.56%	40,765	6.97%	58,538	10.01%	342,048	58.47%	0	0.00%	0	0.00%
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	*	*	*	*	*	*	*	*	*	*	*	*	*
Living with Partner	388,451	0	0.00%	50,778	13.07%	87,629	22.56%	250,044	64.37%	0	0.00%	0	0.00%
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	390,439	120,133	30.77%	28,674	7.34%	126,915	32.51%	114,717	29.38%	0	0.00%	0	0.00%
Poverty income ratio less than 1	252,425	13,693	5.42%	44,686	17.70%	97,074	38.46%	96,972	38.42%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	3,346,992	355,148	10.61%	455,039	13.60%	959,028	28.65%	1,533,971	45.83%	43,806	1.31%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,128,857	210,214	18.62%	106,068	9.40%	386,319	34.22%	406,367	36.00%	19,889	1.76%	0	0.00%
Overweight: 25.0 - 29.9	1,171,399	141,671	12.09%	179,310	15.31%	367,436	31.37%	482,982	41.23%	0	0.00%	0	0.00%
Obese: 30 or higher	1,652,687	127,606	7.72%	243,021	14.70%	401,833	24.31%	856,310	51.81%	23,917	1.45%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	1,753,796	264,125	15.06%	157,949	9.01%	598,373	34.12%	709,433	40.45%	23,917	1.36%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,557,264	150,757	9.68%	290,753	18.67%	367,735	23.61%	748,018	48.03%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	639,496	74,093	11.59%	79,697	12.46%	210,020	32.84%	275,686	43.11%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	585,592	87,726	14.98%	49,267	8.41%	202,194	34.53%	246,405	42.08%	0	0.00%	0	0.00%
No	3,404,264	401,249	11.79%	479,132	14.07%	980,823	28.81%	1,499,254	44.04%	43,806	1.29%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,506,294	349,909	13.96%	363,802	14.52%	691,498	27.59%	1,101,085	43.93%	0	0.00%	0	0.00%
No	1,483,563	139,066	9.37%	164,597	11.09%	491,520	33.13%	644,575	43.45%	43,806	2.95%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,454,018	271,053	11.05%	342,629	13.96%	905,442	36.90%	934,894	38.10%	0	0.00%	0	0.00%
No	1,535,839	217,922	14.19%	185,770	12.10%	277,575	18.07%	810,765	52.79%	43,806	2.85%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	755,983	231,325	30.60%	110,753	14.65%	192,826	25.51%	197,161	26.08%	23,917	3.16%	0	0.00%
Mobility limitations	706,352	97,653	13.82%	98,896	14.00%	294,219	41.65%	195,696	27.71%	19,889	2.82%	0	0.00%
No mobility limitations	958,623	134,669	14.05%	113,069	11.79%	277,992	29.00%	432,893	45.16%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,568,898	25,328	1.61%	205,680	13.11%	417,980	26.64%	919,910	58.63%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	653,360	64,209	9.83%	47,007	7.19%	52,664	8.06%	469,591	71.87%	19,889	3.04%	0	0.00%
No	3,336,497	424,766	12.73%	481,392	14.43%	1,130,354	33.88%	1,276,069	38.25%	23,917	0.72%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	653,142	94,044	14.40%	31,598	4.84%	297,604	45.56%	229,895	35.20%	0	0.00%	0	0.00%
Once a day	334,691	28,835	8.62%	28,070	8.39%	28,751	8.59%	249,036	74.41%	0	0.00%	0	0.00%
2 or more time a week	358,953	0	0.00%	108,452	30.21%	186,144	51.86%	64,358	17.93%	0	0.00%	0	0.00%
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	853,343	77,278	9.06%	70,256	8.23%	341,761	40.05%	340,131	39.86%	23,917	2.80%	0	0.00%
Never	1,663,284	271,972	16.35%	277,670	16.69%	316,242	19.01%	797,400	47.94%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2007 - 2008

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	3,675,867	466,575	12.69%	719,308	19.57%	793,508	21.59%	1,696,477	46.15%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	3,790,447	394,851	10.42%	528,399	13.94%	1,119,805	29.54%	1,723,476	45.47%	23,917	0.63%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,165,544	757,405	18.18%	470,432	11.29%	937,200	22.50%	2,000,507	48.03%	0	0.00%	0	0.00%
Age at screening	-	-	-	-	-	-	-	-	-	-	-	-	-
20-29	*	*	*	*	*	*	*	*	*	*	*	*	*
30-39	*	*	*	*	*	*	*	*	*	*	*	*	*
40-49	923,292	37,984	4.11%	34,901	3.78%	119,452	12.94%	730,955	79.17%	0	0.00%	0	0.00%
50-59	756,720	283,655	37.48%	130,832	17.29%	115,671	15.29%	226,562	29.94%	0	0.00%	0	0.00%
60-64	532,739	82,781	15.54%	129,858	24.38%	98,646	18.52%	221,454	41.57%	0	0.00%	0	0.00%
65-69	482,971	121,642	25.19%	21,723	4.50%	261,391	54.12%	78,215	16.19%	0	0.00%	0	0.00%
70-74	253,350	23,962	9.46%	47,499	18.75%	105,531	41.65%	76,358	30.14%	0	0.00%	0	0.00%
75-79	265,394	45,923	17.30%	0	0.00%	110,982	41.82%	108,489	40.88%	0	0.00%	0	0.00%
80+	479,229	50,907	10.62%	87,239	18.20%	118,606	24.75%	222,477	46.42%	0	0.00%	0	0.00%
Race/ethnicity	-	-	-	-	-	-	-	-	-	-	-	-	-
Mexican American	*	*	*	*	*	*	*	*	*	*	*	*	*
Other Hispanic	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White	3,293,559	627,371	19.05%	367,925	11.17%	716,613	21.76%	1,581,651	48.02%	0	0.00%	0	0.00%
Non-Hispanic Black	513,411	55,283	10.77%	87,889	17.12%	117,022	22.79%	253,217	49.32%	0	0.00%	0	0.00%
Other Race - Including Multi-Racial	*	*	*	*	*	*	*	*	*	*	*	*	*
Education	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Less Than 9th Grade	*	*	*	*	*	*	*	*	*	*	*	*	*
9-11th Grade	514,877	171,887	33.38%	49,867	9.69%	73,385	14.25%	219,737	42.68%	0	0.00%	0	0.00%
High School Grad/GED or Equivalent	3,374,006	544,125	16.13%	311,231	9.22%	779,817	23.11%	1,738,834	51.54%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,165,544	757,405	18.18%	470,432	11.29%	937,200	22.50%	2,000,507	48.03%	0	0.00%	0	0.00%
Marital status	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Married	3,022,250	557,123	18.43%	381,576	12.63%	649,528	21.49%	1,434,023	47.45%	0	0.00%	0	0.00%
Widowed	265,282	64,391	24.27%	49,398	18.62%	74,611	28.13%	76,881	28.98%	0	0.00%	0	0.00%
Divorced	*	*	*	*	*	*	*	*	*	*	*	*	*
Separated	*	*	*	*	*	*	*	*	*	*	*	*	*
Never Married	347,896	40,829	11.74%	0	0.00%	41,403	11.90%	265,663	76.36%	0	0.00%	0	0.00%
Living with Partner	*	*	*	*	*	*	*	*	*	*	*	*	*
Refused	*	*	*	*	*	*	*	*	*	*	*	*	*
Poverty income ratio	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	248,215	75,488	30.41%	14,618	5.89%	83,370	33.59%	74,738	30.11%	0	0.00%	0	0.00%
Poverty income ratio less than 1	345,047	9,775	2.83%	42,801	12.40%	93,004	26.95%	199,466	57.81%	0	0.00%	0	0.00%
Poverty income ratio 1 or greater	3,572,283	672,142	18.82%	413,012	11.56%	760,825	21.30%	1,726,303	48.32%	0	0.00%	0	0.00%
Body mass index	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Normal: less than 25	1,292,395	73,895	5.72%	72,621	5.62%	428,941	33.19%	716,938	55.47%	0	0.00%	0	0.00%
Overweight: 25.0 - 29.9	1,161,381	203,429	17.52%	170,414	14.67%	134,452	11.58%	653,086	56.23%	0	0.00%	0	0.00%
Obese: 30 or higher	1,632,890	480,081	29.40%	170,646	10.45%	373,807	22.89%	608,356	37.26%	0	0.00%	0	0.00%
Glycohemoglobin	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
Glycohemoglobin < 5.7%	2,228,781	261,131	11.72%	190,104	8.53%	460,494	20.66%	1,317,052	59.09%	0	0.00%	0	0.00%
Glycohemoglobin 5.7% - 6.4%	1,372,628	422,014	30.74%	168,920	12.31%	219,613	16.00%	562,082	40.95%	0	0.00%	0	0.00%
Glycohemoglobin 6.5% or higher	516,045	74,260	14.39%	79,976	15.50%	240,435	46.59%	121,374	23.52%	0	0.00%	0	0.00%
Diabetes	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	640,241	180,840	28.25%	79,976	12.49%	312,904	48.87%	66,521	10.39%	0	0.00%	0	0.00%
No	3,525,303	576,566	16.36%	390,455	11.08%	624,296	17.71%	1,933,986	54.86%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,165,544	757,405	18.18%	470,432	11.29%	937,200	22.50%	2,000,507	48.03%	0	0.00%	0	0.00%
Smoked at least 100 cigarettes in life	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	2,237,513	410,800	18.36%	370,943	16.58%	433,699	19.38%	1,022,070	45.68%	0	0.00%	0	0.00%
No	1,928,031	346,605	17.98%	99,488	5.16%	503,501	26.11%	978,438	50.75%	0	0.00%	0	0.00%
High blood pressure	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1,814,026	289,360	15.95%	265,400	14.63%	461,494	25.44%	797,772	43.98%	0	0.00%	0	0.00%
No	2,351,518	468,046	19.90%	205,031	8.72%	475,706	20.23%	1,202,735	51.15%	0	0.00%	0	0.00%
Mobility limitation	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	697,283	102,124	14.65%	222,052	31.85%	226,789	32.52%	146,320	20.98%	0	0.00%	0	0.00%
Mobility limitations	719,633	134,661	18.71%	80,496	11.19%	214,837	29.85%	289,639	40.25%	0	0.00%	0	0.00%
No mobility limitations	1,235,244	268,086	21.70%	131,122	10.62%	377,099	30.53%	458,937	37.15%	0	0.00%	0	0.00%
N/A - Respondent not asked mobility limitation questions	1,513,383	252,534	16.69%	36,762	2.43%	118,475	7.83%	1,105,612	73.06%	0	0.00%	0	0.00%
Sleep disorder	-	-	-	-	-	-	-	-	-	-	-	-	-
Yes	785,278	290,320	36.97%	151,781	19.33%	174,714	22.25%	168,464	21.45%	0	0.00%	0	0.00%
No	3,380,266	467,086	13.82%	318,651	9.43%	762,486	22.56%	1,832,044	54.20%	0	0.00%	0	0.00%
Bowel leakage - gas	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more times a day	485,649	115,914	23.87%	88,855	18.30%	137,025	28.21%	143,855	29.62%	0	0.00%	0	0.00%
Once a day	547,595	120,327	21.97%	12,081	2.21%	88,988	16.25%	326,199	59.57%	0	0.00%	0	0.00%
2 or more time a week	456,027	0	0.00%	31,431	6.89%	119,836	26.28%	304,760	66.83%	0	0.00%	0	0.00%
Once a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	795,351	61,769	7.77%	56,751	7.14%	187,785	23.61%	489,046	61.49%	0	0.00%	0	0.00%
Never	1,512,576	287,355	19.00%	232,896	15.40%	300,483	19.87%	691,842	45.74%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.047: How frequently does this (referring to KIQ.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?

Table M05a -Frequency of bladder control problems among those who responded 'yes' to insensate urine loss among men, 2009 - 2010

	Total	Every day/ night (N)	Every day/ night (%)	Few per week (N)	Few per week (%)	Few per month (N)	Few per month (%)	Less than once a month (N)	Less than once a month (%)	Refused to Answer or Don't Know (N)	Refused to Answer or Don't Know (%)	Missing (N)	Missing (%)
Total	4,165,544	757,405	18.18%	470,432	11.29%	937,200	22.50%	2,000,507	48.03%	0	0.00%	0	0.00%
Bowel leakage - stool	-	-	-	-	-	-	-	-	-	-	-	-	-
Missing	*	*	*	*	*	*	*	*	*	*	*	*	*
2 or more time a week	*	*	*	*	*	*	*	*	*	*	*	*	*
1-3 times a month	*	*	*	*	*	*	*	*	*	*	*	*	*
Never	3,887,069	698,789	17.98%	429,242	11.04%	860,276	22.13%	1,898,762	48.85%	0	0.00%	0	0.00%

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.047: How frequently does this (referring to KI.Q.046) occur? Would you say this occurs...every day, a few times a week, a few times a month, or a few times a year?