

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 30-39 | 1,245,096 | 454,685 | 36.52% | 523,029 | 42.01% | 198,855 | 15.97% | 38,014 | 3.05% | 30,513 | 2.45% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 1,282,331 | 597,613 | 46.60% | 333,750 | 26.03% | 255,712 | 19.94% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 95,257 | 7.43% |
| 50-59 | 2,024,486 | 641,594 | 31.69% | 652,328 | 32.22% | 422,812 | 20.88% | 159,788 | 7.89% | 147,963 | 7.31% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 929,881 | 188,500 | 20.27% | 459,038 | 49.37% | 199,730 | 21.48% | 54,701 | 5.88% | 27,912 | 3.00% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,080,369 | 308,004 | 28.51% | 421,963 | 39.06% | 252,861 | 23.41% | 33,873 | 3.14% | 63,668 | 5.89% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,373,568 | 338,895 | 24.67% | 551,219 | 40.13% | 348,917 | 25.40% | 43,943 | 3.20% | 76,466 | 5.57% | 14,128 | 1.03% | 0 | 0.00% |
| 75-79 | 758,372 | 95,832 | 12.64% | 309,228 | 40.78% | 244,006 | 32.17% | 84,071 | 11.09% | 25,235 | 3.33% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 624,458 | 229,505 | 36.75% | 171,479 | 27.46% | 150,954 | 24.17% | 54,878 | 8.79% | 17,643 | 2.83% | 0 | 0.00% | 0 | 0.00% |
| 85+ | 566,783 | 193,213 | 34.09% | 230,500 | 40.67% | 55,118 | 9.72% | 29,512 | 5.21% | 58,440 | 10.31% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 475,436 | 145,030 | 30.50% | 189,115 | 39.78% | 119,430 | 25.12% | 19,764 | 4.16% | 2,097 | 0.44% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 553,379 | 255,510 | 46.17% | 146,882 | 26.54% | 68,615 | 12.40% | 7,286 | 1.32% | 75,086 | 13.57% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 7,730,812 | 2,266,831 | 29.32% | 2,946,017 | 38.11% | 1,645,658 | 21.29% | 444,802 | 5.75% | 332,246 | 4.30% | 0 | 0.00% | 95,257 | 1.23% |
| Non-Hispanic Black | 1,311,346 | 491,237 | 37.46% | 429,144 | 32.73% | 311,498 | 23.75% | 26,927 | 2.05% | 38,411 | 2.93% | 14,128 | 1.08% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,274,348 | 361,469 | 28.37% | 492,441 | 38.64% | 198,997 | 15.62% | 146,444 | 11.49% | 74,998 | 5.89% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,393,333 | 421,946 | 30.28% | 513,840 | 36.88% | 252,331 | 18.11% | 105,350 | 7.56% | 99,866 | 7.17% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 7,469,140 | 2,408,357 | 32.24% | 2,796,398 | 37.44% | 1,635,038 | 21.89% | 246,986 | 3.31% | 272,976 | 3.65% | 14,128 | 0.19% | 95,257 | 1.28% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 6,617,196 | 1,865,739 | 28.20% | 2,576,351 | 38.93% | 1,442,539 | 21.80% | 373,050 | 5.64% | 345,390 | 5.22% | 14,128 | 0.21% | 0 | 0.00% |
| Widowed | 861,620 | 252,099 | 29.26% | 344,511 | 39.98% | 162,978 | 18.92% | 24,817 | 2.88% | 77,215 | 8.96% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,262,656 | 456,658 | 36.17% | 441,022 | 34.93% | 226,741 | 17.96% | 42,978 | 3.40% | 0 | 0.00% | 0 | 0.00% | 95,257 | 7.54% |
| Separated | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Never Married | 983,139 | 376,419 | 38.29% | 286,285 | 29.12% | 277,329 | 28.21% | 17,870 | 1.82% | 25,235 | 2.57% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 593,049 | 236,654 | 39.90% | 164,284 | 27.70% | 91,241 | 15.39% | 5,612 | 0.95% | 0 | 0.00% | 0 | 0.00% | 95,257 | 16.06% |
| Poverty income ratio less than 1 | 1,516,688 | 591,004 | 38.97% | 571,305 | 37.67% | 200,378 | 13.21% | 12,061 | 0.80% | 141,940 | 9.36% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 8,156,673 | 2,423,794 | 29.72% | 3,067,089 | 37.60% | 1,864,655 | 22.86% | 481,107 | 5.90% | 305,900 | 3.75% | 14,128 | 0.17% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 874,992 | 215,893 | 24.67% | 279,367 | 31.93% | 233,134 | 26.64% | 41,055 | 4.69% | 105,542 | 12.06% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 2,739,084 | 801,293 | 29.25% | 1,194,736 | 43.62% | 525,621 | 19.19% | 73,143 | 2.67% | 49,035 | 1.79% | 0 | 0.00% | 95,257 | 3.48% |
| Overweight: 25.0 - 29.9 | 3,551,109 | 1,233,523 | 34.74% | 1,220,223 | 34.36% | 820,992 | 23.12% | 231,730 | 6.53% | 30,513 | 0.86% | 14,128 | 0.40% | 0 | 0.00% |
| Obese: 30 or higher | 3,101,225 | 1,000,743 | 32.27% | 1,108,352 | 35.74% | 576,528 | 18.59% | 152,852 | 4.93% | 262,749 | 8.47% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 416,345 | 294,500 | 70.73% | 73,784 | 17.72% | 2,737 | 0.66% | 14,812 | 3.56% | 30,513 | 7.33% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 6,339,137 | 1,996,593 | 31.50% | 2,163,676 | 34.13% | 1,543,634 | 24.35% | 385,056 | 6.07% | 154,921 | 2.44% | 0 | 0.00% | 95,257 | 1.50% |
| Glycohemoglobin 5.7% - 6.4% | 1,757,161 | 448,999 | 25.55% | 741,597 | 42.20% | 377,445 | 21.48% | 63,324 | 3.60% | 125,796 | 7.16% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,753,766 | 511,360 | 29.16% | 823,621 | 46.96% | 232,459 | 13.25% | 35,588 | 2.03% | 136,610 | 7.79% | 14,128 | 0.81% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,695,100 | 449,780 | 26.53% | 883,009 | 52.09% | 203,709 | 12.02% | 78,991 | 4.66% | 79,611 | 4.70% | 0 | 0.00% | 0 | 0.00% |
| No | 8,571,309 | 2,801,672 | 32.69% | 2,919,669 | 34.06% | 1,952,565 | 22.78% | 419,789 | 4.90% | 368,229 | 4.30% | 14,128 | 0.16% | 95,257 | 1.11% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 7,293,348 | 2,365,729 | 32.44% | 2,679,045 | 36.73% | 1,479,740 | 20.29% | 385,900 | 5.29% | 368,806 | 5.06% | 14,128 | 0.19% | 0 | 0.00% |
| No | 2,973,061 | 885,723 | 29.79% | 1,123,633 | 37.79% | 676,534 | 22.76% | 112,880 | 3.80% | 79,034 | 2.66% | 0 | 0.00% | 95,257 | 3.20% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Yes | 3,876,071 | 1,072,463 | 27.67% | 1,519,011 | 39.19% | 775,122 | 20.00% | 188,025 | 4.85% | 321,451 | 8.29% | 0 | 0.00% | 0 | 0.00% |
| No | 6,380,301 | 2,168,953 | 33.99% | 2,283,667 | 35.79% | 1,381,152 | 21.65% | 310,755 | 4.87% | 126,389 | 1.98% | 14,128 | 0.22% | 95,257 | 1.49% |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,480,096 | 187,485 | 12.67% | 643,384 | 43.47% | 265,530 | 17.94% | 113,434 | 7.66% | 256,135 | 17.31% | 14,128 | 0.95% | 0 | 0.00% |
| Mobility limitations | 2,270,983 | 656,701 | 28.92% | 771,751 | 33.98% | 556,411 | 24.50% | 198,094 | 8.72% | 88,025 | 3.88% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 3,383,951 | 1,023,668 | 30.25% | 1,371,821 | 40.54% | 809,179 | 23.91% | 106,116 | 3.14% | 73,167 | 2.16% | 0 | 0.00% | 0 | 0.00% |
| Refused/Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| N/A - Respondent not asked mobility limitation questions | 3,116,764 | 1,368,983 | 43.92% | 1,015,722 | 32.59% | 525,154 | 16.85% | 81,135 | 2.60% | 30,513 | 0.98% | 0 | 0.00% | 95,257 | 3.06% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 13,702,607 | 5,146,367 | 37.56% | 4,630,460 | 33.79% | 2,208,640 | 16.12% | 1,023,404 | 7.47% | 693,736 | 5.06% | 0 | 0.00% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 20-29 | 920,444 | 353,337 | 38.39% | 284,975 | 30.96% | 118,735 | 12.90% | 91,429 | 9.93% | 71,968 | 7.82% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,961,991 | 829,089 | 42.26% | 369,534 | 18.83% | 479,814 | 24.46% | 144,669 | 7.37% | 138,885 | 7.08% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 1,948,077 | 830,659 | 42.64% | 755,413 | 38.78% | 188,613 | 9.68% | 126,132 | 6.47% | 47,260 | 2.43% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 2,843,199 | 1,089,396 | 38.32% | 1,046,470 | 36.81% | 465,182 | 16.36% | 111,534 | 3.92% | 130,617 | 4.59% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,000,574 | 342,605 | 34.24% | 316,645 | 31.65% | 129,912 | 12.98% | 130,479 | 13.04% | 80,933 | 8.09% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,335,420 | 562,393 | 42.11% | 499,360 | 37.39% | 137,506 | 10.30% | 61,495 | 4.60% | 74,665 | 5.59% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,398,131 | 454,416 | 32.50% | 529,412 | 37.87% | 203,435 | 14.55% | 164,460 | 11.76% | 46,407 | 3.32% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 1,080,370 | 303,967 | 28.14% | 453,349 | 41.96% | 266,421 | 24.66% | 54,374 | 5.03% | 2,260 | 0.21% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 662,074 | 200,038 | 30.21% | 249,377 | 37.67% | 116,317 | 17.57% | 75,838 | 11.45% | 20,504 | 3.10% | 0 | 0.00% | 0 | 0.00% |
| 85+ | 552,328 | 180,467 | 32.67% | 125,925 | 22.80% | 102,705 | 18.59% | 62,994 | 11.41% | 80,236 | 14.53% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 987,200 | 437,506 | 44.32% | 337,136 | 34.15% | 129,775 | 13.15% | 45,551 | 4.61% | 37,231 | 3.77% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 501,938 | 87,439 | 17.42% | 263,672 | 52.53% | 0 | 0.00% | 83,745 | 16.68% | 67,083 | 13.36% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 9,809,316 | 3,507,441 | 35.76% | 3,309,192 | 33.74% | 1,744,047 | 17.78% | 745,183 | 7.60% | 503,453 | 5.13% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic Black | 2,024,497 | 994,922 | 49.14% | 489,638 | 24.19% | 305,044 | 15.07% | 148,925 | 7.36% | 85,969 | 4.25% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,425,778 | 558,146 | 39.15% | 435,491 | 30.54% | 203,895 | 14.30% | 145,575 | 10.21% | 82,673 | 5.80% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,149,203 | 424,891 | 36.97% | 359,113 | 31.25% | 130,769 | 11.38% | 122,050 | 10.62% | 112,380 | 9.78% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 10,919,458 | 4,150,483 | 38.01% | 3,765,586 | 34.49% | 1,837,518 | 16.83% | 752,424 | 6.89% | 413,447 | 3.79% | 0 | 0.00% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 8,855,567 | 3,362,037 | 37.97% | 3,059,260 | 34.55% | 1,497,854 | 16.91% | 559,171 | 6.31% | 377,246 | 4.26% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 869,715 | 261,032 | 30.01% | 219,573 | 25.25% | 175,747 | 20.21% | 94,057 | 10.81% | 119,306 | 13.72% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,217,057 | 494,786 | 40.65% | 380,979 | 31.30% | 135,409 | 11.13% | 205,883 | 16.92% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Separated | 334,445 | 170,503 | 50.98% | 71,464 | 21.37% | 18,209 | 5.44% | 64,901 | 19.41% | 9,367 | 2.80% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 1,587,594 | 458,536 | 28.88% | 686,318 | 43.23% | 306,850 | 19.33% | 62,727 | 3.95% | 73,162 | 4.61% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 622,715 | 225,583 | 36.23% | 177,430 | 28.49% | 74,570 | 11.97% | 30,477 | 4.89% | 114,655 | 18.41% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 572,832 | 233,552 | 40.77% | 120,176 | 20.98% | 125,895 | 21.98% | 14,160 | 2.47% | 79,049 | 13.80% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,153,158 | 539,977 | 46.83% | 158,675 | 13.76% | 210,919 | 18.29% | 69,810 | 6.05% | 173,779 | 15.07% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,976,617 | 4,372,839 | 36.51% | 4,351,608 | 36.33% | 1,871,826 | 15.63% | 939,434 | 7.84% | 440,909 | 3.68% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 268,019 | 69,485 | 25.93% | 21,496 | 8.02% | 21,276 | 7.94% | 155,763 | 58.12% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 3,529,345 | 1,194,516 | 33.85% | 1,379,014 | 39.07% | 518,379 | 14.69% | 129,142 | 3.66% | 308,295 | 8.74% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 4,301,128 | 2,013,203 | 46.81% | 1,491,959 | 34.69% | 339,424 | 7.89% | 350,877 | 8.16% | 105,665 | 2.46% | 0 | 0.00% | 0 | 0.00% |
| Obese: 30 or higher | 5,604,114 | 1,869,163 | 33.35% | 1,737,991 | 31.01% | 1,329,562 | 23.72% | 387,623 | 6.92% | 279,776 | 4.99% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 361,588 | 133,498 | 36.92% | 70,215 | 19.42% | 123,143 | 34.06% | 0 | 0.00% | 34,732 | 9.61% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 8,890,742 | 3,485,326 | 39.20% | 3,147,608 | 35.40% | 1,278,785 | 14.38% | 633,019 | 7.12% | 346,004 | 3.89% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 2,728,162 | 834,481 | 30.59% | 993,531 | 36.42% | 601,732 | 22.06% | 218,532 | 8.01% | 79,886 | 2.93% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,722,115 | 693,061 | 40.24% | 419,106 | 24.34% | 204,980 | 11.90% | 171,853 | 9.98% | 233,115 | 13.54% | 0 | 0.00% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,199,006 | 785,784 | 35.73% | 629,900 | 28.64% | 295,462 | 13.44% | 211,023 | 9.60% | 276,837 | 12.59% | 0 | 0.00% | 0 | 0.00% |
| No | 11,503,602 | 4,360,583 | 37.91% | 4,000,560 | 34.78% | 1,913,179 | 16.63% | 812,380 | 7.06% | 416,900 | 3.62% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 3,251,452 | 31.67% | 3,802,678 | 37.04% | 2,156,274 | 21.00% | 498,780 | 4.86% | 447,840 | 4.36% | 14,128 | 0.14% | 95,257 | 0.93% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 8,509,424 | 3,257,577 | 38.28% | 2,559,123 | 30.07% | 1,352,409 | 15.89% | 791,892 | 9.31% | 548,422 | 6.44% | 0 | 0.00% | 0 | 0.00% |
| No | 5,181,352 | 1,888,790 | 36.45% | 2,071,336 | 39.98% | 856,231 | 16.53% | 219,681 | 4.24% | 145,314 | 2.80% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Yes | 6,058,883 | 2,176,557 | 35.92% | 1,832,374 | 30.24% | 1,108,544 | 18.30% | 534,197 | 8.82% | 407,211 | 6.72% | 0 | 0.00% | 0 | 0.00% |
| No | 7,560,067 | 2,935,702 | 38.83% | 2,769,041 | 36.63% | 1,100,096 | 14.55% | 468,703 | 6.20% | 286,525 | 3.79% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,964,807 | 664,860 | 22.43% | 918,072 | 30.97% | 569,736 | 19.22% | 527,784 | 17.80% | 284,354 | 9.59% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 1,853,409 | 582,240 | 31.41% | 502,327 | 27.10% | 499,581 | 26.95% | 98,651 | 5.32% | 170,609 | 9.21% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 3,470,630 | 1,385,969 | 39.93% | 1,345,341 | 38.76% | 460,834 | 13.28% | 133,342 | 3.84% | 145,144 | 4.18% | 0 | 0.00% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 5,413,761 | 2,513,298 | 46.42% | 1,864,719 | 34.44% | 678,489 | 12.53% | 263,627 | 4.87% | 93,629 | 1.73% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 5,146,367 | 37.56% | 4,630,460 | 33.79% | 2,208,640 | 16.12% | 1,023,404 | 7.47% | 693,736 | 5.06% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 5,146,367 | 37.56% | 4,630,460 | 33.79% | 2,208,640 | 16.12% | 1,023,404 | 7.47% | 693,736 | 5.06% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 5,146,367 | 37.56% | 4,630,460 | 33.79% | 2,208,640 | 16.12% | 1,023,404 | 7.47% | 693,736 | 5.06% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 6,274,123 | 44.43% | 4,896,101 | 34.67% | 2,151,733 | 15.24% | 527,345 | 3.73% | 246,932 | 1.75% | 26,515 | 0.19% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 899,897 | 554,584 | 61.63% | 255,539 | 28.40% | 65,003 | 7.22% | 0 | 0.00% | 24,771 | 2.75% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,031,971 | 415,966 | 40.31% | 302,630 | 29.33% | 285,821 | 27.70% | 27,554 | 2.67% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,978,059 | 1,505,710 | 50.56% | 715,874 | 24.04% | 418,180 | 14.04% | 320,115 | 10.75% | 18,179 | 0.61% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,614,824 | 1,777,040 | 49.16% | 974,867 | 26.97% | 718,996 | 19.89% | 35,642 | 0.99% | 108,280 | 3.00% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,085,089 | 250,338 | 23.07% | 526,819 | 48.55% | 223,342 | 20.58% | 28,011 | 2.58% | 56,579 | 5.21% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,466,777 | 653,409 | 44.55% | 706,007 | 48.13% | 31,297 | 2.13% | 66,242 | 4.52% | 9,823 | 0.67% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,031,394 | 517,357 | 50.16% | 414,016 | 40.14% | 39,125 | 3.79% | 39,545 | 3.83% | 21,351 | 2.07% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 959,797 | 244,668 | 25.49% | 491,011 | 51.16% | 224,118 | 23.35% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 733,159 | 248,727 | 33.93% | 375,064 | 51.16% | 64,667 | 8.82% | 10,236 | 1.40% | 7,950 | 1.08% | 26,515 | 3.62% | 0 | 0.00% |
| 85+ | 299,135 | 106,324 | 35.54% | 111,626 | 37.32% | 81,185 | 27.14% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,116,500 | 648,146 | 58.05% | 323,331 | 28.96% | 64,254 | 5.75% | 80,769 | 7.23% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic White | 10,764,351 | 4,382,199 | 40.71% | 4,183,270 | 38.86% | 1,539,581 | 14.30% | 446,576 | 4.15% | 186,210 | 1.73% | 26,515 | 0.25% | 0 | 0.00% |
| Non-Hispanic Black | 1,520,452 | 925,270 | 60.85% | 222,158 | 14.61% | 312,302 | 20.54% | 0 | 0.00% | 60,722 | 3.99% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,110,708 | 561,931 | 50.59% | 475,946 | 42.85% | 63,335 | 5.70% | 9,496 | 0.85% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,514,077 | 698,590 | 46.14% | 523,975 | 34.61% | 259,465 | 17.14% | 24,098 | 1.59% | 7,950 | 0.53% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 11,429,497 | 4,983,316 | 43.60% | 3,861,906 | 33.79% | 1,825,026 | 15.97% | 493,751 | 4.32% | 238,983 | 2.09% | 26,515 | 0.23% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 6,274,123 | 44.43% | 4,896,101 | 34.67% | 2,151,733 | 15.24% | 527,345 | 3.73% | 246,932 | 1.75% | 26,515 | 0.19% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 9,694,781 | 4,091,741 | 42.21% | 3,440,084 | 35.48% | 1,739,490 | 17.94% | 330,551 | 3.41% | 66,401 | 0.68% | 26,515 | 0.27% | 0 | 0.00% |
| Widowed | 577,904 | 321,249 | 55.59% | 180,329 | 31.20% | 68,376 | 11.83% | 0 | 0.00% | 7,950 | 1.38% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,797,233 | 937,255 | 52.15% | 613,007 | 34.11% | 116,069 | 6.46% | 130,902 | 7.28% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Separated | 472,734 | 244,634 | 51.75% | 160,393 | 33.93% | 46,356 | 9.81% | 0 | 0.00% | 21,351 | 4.52% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 999,423 | 476,158 | 47.64% | 361,739 | 36.19% | 57,915 | 5.79% | 60,660 | 6.07% | 42,950 | 4.30% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 437,370 | 194,115 | 44.38% | 10,123 | 2.31% | 119,620 | 27.35% | 5,231 | 1.20% | 108,280 | 24.76% | 0 | 0.00% | 0 | 0.00% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 811,536 | 489,346 | 60.30% | 208,731 | 25.72% | 113,458 | 13.98% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,447,959 | 767,033 | 52.97% | 500,095 | 34.54% | 91,975 | 6.35% | 88,856 | 6.14% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,863,253 | 5,017,743 | 42.30% | 4,187,274 | 35.30% | 1,946,300 | 16.41% | 438,489 | 3.70% | 246,932 | 2.08% | 26,515 | 0.22% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Normal: less than 25 | 3,052,898 | 1,263,379 | 41.38% | 1,091,654 | 35.76% | 442,124 | 14.48% | 139,512 | 4.57% | 116,229 | 3.81% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 5,408,869 | 2,136,288 | 39.50% | 2,153,033 | 39.81% | 861,631 | 15.93% | 174,823 | 3.23% | 56,579 | 1.05% | 26,515 | 0.49% | 0 | 0.00% |
| Obese: 30 or higher | 5,534,849 | 2,810,709 | 50.78% | 1,635,283 | 29.55% | 801,723 | 14.49% | 213,010 | 3.85% | 74,124 | 1.34% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 300,823 | 167,634 | 55.73% | 83,126 | 27.63% | 44,480 | 14.79% | 5,582 | 1.86% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 9,038,263 | 3,880,094 | 42.93% | 3,233,392 | 35.77% | 1,279,630 | 14.16% | 467,808 | 5.18% | 150,823 | 1.67% | 26,515 | 0.29% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 3,182,425 | 1,438,200 | 45.19% | 1,147,875 | 36.07% | 546,585 | 17.18% | 10,236 | 0.32% | 39,530 | 1.24% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,601,238 | 788,195 | 49.22% | 431,708 | 26.96% | 281,038 | 17.55% | 43,718 | 2.73% | 56,579 | 3.53% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 6,274,123 | 44.43% | 4,896,101 | 34.67% | 2,151,733 | 15.24% | 527,345 | 3.73% | 246,932 | 1.75% | 26,515 | 0.19% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,702,435 | 966,827 | 56.79% | 461,032 | 27.08% | 166,212 | 9.76% | 30,434 | 1.79% | 77,930 | 4.58% | 0 | 0.00% | 0 | 0.00% |
| No | 12,420,313 | 5,307,296 | 42.73% | 4,435,069 | 35.71% | 1,985,520 | 15.99% | 496,910 | 4.00% | 169,002 | 1.36% | 26,515 | 0.21% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 9,608,565 | 4,038,394 | 42.03% | 3,361,706 | 34.99% | 1,512,787 | 15.74% | 481,467 | 5.01% | 214,211 | 2.23% | 0 | 0.00% | 0 | 0.00% |
| No | 4,514,183 | 2,235,729 | 49.53% | 1,534,394 | 33.99% | 638,946 | 14.15% | 45,877 | 1.02% | 32,721 | 0.72% | 26,515 | 0.59% | 0 | 0.00% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 6,581,507 | 2,520,777 | 38.30% | 2,362,301 | 35.89% | 1,281,379 | 19.47% | 359,747 | 5.47% | 57,303 | 0.87% | 0 | 0.00% | 0 | 0.00% |
| No | 7,484,663 | 3,753,345 | 50.15% | 2,533,800 | 33.85% | 870,354 | 11.63% | 167,598 | 2.24% | 133,051 | 1.78% | 26,515 | 0.35% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,617,259 | 390,454 | 24.14% | 516,952 | 31.96% | 494,219 | 30.56% | 137,704 | 8.51% | 77,930 | 4.82% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 2,018,270 | 922,555 | 45.71% | 811,108 | 40.19% | 284,607 | 14.10% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 4,434,930 | 1,732,858 | 39.07% | 2,016,915 | 45.48% | 448,374 | 10.11% | 192,495 | 4.34% | 17,772 | 0.40% | 26,515 | 0.60% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 6,052,290 | 3,228,255 | 53.34% | 1,551,126 | 25.63% | 924,533 | 15.28% | 197,146 | 3.26% | 151,230 | 2.50% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,079,229 | 677,106 | 32.57% | 846,686 | 40.72% | 513,724 | 24.71% | 23,534 | 1.13% | 18,179 | 0.87% | 0 | 0.00% | 0 | 0.00% |
| No | 12,043,519 | 5,597,017 | 46.47% | 4,049,415 | 33.62% | 1,638,009 | 13.60% | 503,810 | 4.18% | 228,753 | 1.90% | 26,515 | 0.22% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 6,274,123 | 44.43% | 4,896,101 | 34.67% | 2,151,733 | 15.24% | 527,345 | 3.73% | 246,932 | 1.75% | 26,515 | 0.19% | 0 | 0.00% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 2,904,617 | 1,171,933 | 40.35% | 1,040,148 | 35.81% | 445,291 | 15.33% | 239,295 | 8.24% | 7,950 | 0.27% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,026,938 | 644,894 | 62.80% | 240,417 | 23.41% | 95,214 | 9.27% | 46,413 | 4.52% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,451,505 | 558,779 | 38.50% | 362,477 | 24.97% | 494,608 | 34.08% | 35,642 | 2.46% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 1,065,210 | 233,091 | 21.88% | 511,013 | 47.97% | 186,312 | 17.49% | 0 | 0.00% | 108,280 | 10.17% | 26,515 | 2.49% | 0 | 0.00% |
| 1-3 times a month | 2,228,101 | 933,089 | 41.88% | 876,750 | 39.35% | 215,539 | 9.67% | 127,964 | 5.74% | 74,757 | 3.36% | 0 | 0.00% | 0 | 0.00% |
| Never | 5,381,424 | 2,709,816 | 50.35% | 1,841,562 | 34.22% | 696,069 | 12.93% | 78,032 | 1.45% | 55,945 | 1.04% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 598,247 | 70,436 | 11.77% | 371,604 | 62.12% | 33,825 | 5.65% | 122,382 | 20.46% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 13,338,270 | 6,181,165 | 46.34% | 4,473,285 | 33.54% | 2,026,761 | 15.20% | 404,963 | 3.04% | 225,581 | 1.69% | 26,515 | 0.20% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 6,738,730 | 45.30% | 4,693,232 | 31.55% | 2,122,591 | 14.27% | 852,436 | 5.73% | 386,354 | 2.60% | 62,237 | 0.42% | 19,889 | 0.13% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 604,177 | 412,907 | 68.34% | 166,773 | 27.60% | 0 | 0.00% | 0 | 0.00% | 24,497 | 4.05% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,125,718 | 791,878 | 70.34% | 152,468 | 13.54% | 62,001 | 5.51% | 97,220 | 8.64% | 22,151 | 1.97% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,794,224 | 1,351,597 | 48.37% | 583,940 | 20.90% | 709,444 | 25.39% | 90,787 | 3.25% | 58,456 | 2.09% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,385,303 | 1,494,907 | 44.16% | 1,131,078 | 33.41% | 483,523 | 14.28% | 202,598 | 5.98% | 53,308 | 1.57% | 0 | 0.00% | 19,889 | 0.59% |
| 60-64 | 1,707,386 | 743,310 | 43.53% | 677,068 | 39.66% | 186,788 | 10.94% | 85,629 | 5.02% | 7,722 | 0.45% | 6,869 | 0.40% | 0 | 0.00% |
| 65-69 | 1,518,446 | 503,764 | 33.18% | 695,808 | 45.82% | 175,987 | 11.59% | 69,968 | 4.61% | 72,919 | 4.80% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,283,484 | 560,925 | 43.70% | 350,384 | 27.30% | 174,736 | 13.61% | 128,066 | 9.98% | 63,276 | 4.93% | 6,097 | 0.48% | 0 | 0.00% |
| 75-79 | 1,207,845 | 391,386 | 32.40% | 548,271 | 45.39% | 174,699 | 14.46% | 56,330 | 4.66% | 30,951 | 2.56% | 6,208 | 0.51% | 0 | 0.00% |
| 80+ | 1,233,178 | 488,056 | 39.58% | 371,735 | 30.14% | 155,412 | 12.60% | 121,839 | 9.88% | 53,074 | 4.30% | 43,062 | 3.49% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,208,868 | 607,559 | 50.26% | 395,408 | 32.71% | 66,631 | 5.51% | 73,989 | 6.12% | 59,073 | 4.89% | 6,208 | 0.51% | 0 | 0.00% |
| Other Hispanic | 624,985 | 256,810 | 41.09% | 253,573 | 40.57% | 27,060 | 4.33% | 59,293 | 9.49% | 22,151 | 3.54% | 6,097 | 0.98% | 0 | 0.00% |
| Non-Hispanic White | 10,135,298 | 4,619,666 | 45.58% | 3,223,988 | 31.81% | 1,410,450 | 13.92% | 577,504 | 5.70% | 273,528 | 2.70% | 30,163 | 0.30% | 0 | 0.00% |
| Non-Hispanic Black | 2,092,450 | 1,125,410 | 53.78% | 461,047 | 22.03% | 324,685 | 15.52% | 141,649 | 6.77% | 0 | 0.00% | 19,768 | 0.94% | 19,889 | 0.95% |
| Other Race - Including Multi-Racial | 813,867 | 129,286 | 15.89% | 359,215 | 44.14% | 293,764 | 36.09% | 0 | 0.00% | 31,602 | 3.88% | 0 | 0.00% | 0 | 0.00% |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,853,886 | 927,415 | 50.03% | 469,131 | 25.31% | 235,725 | 12.72% | 106,477 | 5.74% | 102,060 | 5.51% | 13,077 | 0.71% | 0 | 0.00% |
| 9-11th Grade | 2,135,076 | 1,085,274 | 50.83% | 511,219 | 23.94% | 344,993 | 16.16% | 144,402 | 6.76% | 29,298 | 1.37% | 0 | 0.00% | 19,889 | 0.93% |
| High School Grad/GED or Equivalent | 10,623,592 | 4,498,401 | 42.34% | 3,677,607 | 34.62% | 1,541,873 | 14.51% | 601,557 | 5.66% | 254,995 | 2.40% | 49,159 | 0.46% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KI.Q.042 and KI.Q.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 6,738,730 | 45.30% | 4,693,232 | 31.55% | 2,122,591 | 14.27% | 852,436 | 5.73% | 386,354 | 2.60% | 62,237 | 0.42% | 19,889 | 0.13% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 9,743,242 | 4,417,138 | 45.34% | 3,048,201 | 31.29% | 1,599,237 | 16.41% | 507,552 | 5.21% | 126,021 | 1.29% | 25,204 | 0.26% | 19,889 | 0.20% |
| Widowed | 884,467 | 359,816 | 40.68% | 325,242 | 36.77% | 78,842 | 8.91% | 64,917 | 7.34% | 48,780 | 5.52% | 6,869 | 0.78% | 0 | 0.00% |
| Divorced | 1,942,388 | 895,697 | 46.11% | 648,184 | 33.37% | 88,831 | 4.57% | 130,453 | 6.72% | 149,060 | 7.67% | 30,163 | 1.55% | 0 | 0.00% |
| Separated | 423,395 | 170,484 | 40.27% | 156,798 | 37.03% | 53,689 | 12.68% | 31,283 | 7.39% | 11,141 | 2.63% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 818,909 | 414,939 | 50.67% | 269,732 | 32.94% | 86,235 | 10.53% | 48,003 | 5.86% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 1,013,788 | 447,085 | 44.10% | 229,366 | 22.62% | 215,757 | 21.28% | 70,228 | 6.93% | 51,351 | 5.07% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,314,414 | 461,437 | 35.11% | 235,444 | 17.91% | 350,463 | 26.66% | 137,904 | 10.49% | 122,958 | 9.35% | 6,208 | 0.47% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,579,500 | 766,200 | 48.51% | 415,397 | 26.30% | 232,550 | 14.72% | 94,608 | 5.99% | 57,779 | 3.66% | 12,967 | 0.82% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,981,554 | 5,511,093 | 46.00% | 4,042,391 | 33.74% | 1,539,578 | 12.85% | 619,923 | 5.17% | 205,618 | 1.72% | 43,062 | 0.36% | 19,889 | 0.17% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 211,693 | 110,074 | 52.00% | 27,430 | 12.96% | 44,026 | 20.80% | 0 | 0.00% | 0 | 0.00% | 30,163 | 14.25% | 0 | 0.00% |
| Normal: less than 25 | 3,280,900 | 1,622,323 | 49.45% | 1,034,755 | 31.54% | 349,461 | 10.65% | 161,663 | 4.93% | 86,600 | 2.64% | 6,208 | 0.19% | 19,889 | 0.61% |
| Overweight: 25.0 - 29.9 | 5,264,464 | 2,043,567 | 38.82% | 1,762,013 | 33.47% | 1,190,681 | 22.62% | 175,003 | 3.32% | 87,103 | 1.65% | 6,097 | 0.12% | 0 | 0.00% |
| Obese: 30 or higher | 6,118,411 | 2,962,766 | 48.42% | 1,869,034 | 30.55% | 538,423 | 8.80% | 515,770 | 8.43% | 212,651 | 3.48% | 19,768 | 0.32% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 182,953 | 54,153 | 29.60% | 65,315 | 35.70% | 29,837 | 16.31% | 6,889 | 3.77% | 0 | 0.00% | 6,869 | 3.75% | 19,889 | 10.87% |
| Glycohemoglobin < 5.7% | 7,603,718 | 3,270,082 | 43.01% | 2,795,356 | 36.76% | 997,865 | 13.12% | 385,112 | 5.06% | 155,303 | 2.04% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 5,052,896 | 2,566,624 | 50.80% | 1,198,208 | 23.71% | 789,775 | 15.63% | 322,975 | 6.39% | 126,155 | 2.50% | 49,159 | 0.97% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 2,035,901 | 847,870 | 41.65% | 634,353 | 31.16% | 305,114 | 14.99% | 137,459 | 6.75% | 104,897 | 5.15% | 6,208 | 0.30% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,791,313 | 1,149,673 | 41.19% | 1,075,983 | 38.55% | 323,162 | 11.58% | 112,394 | 4.03% | 104,897 | 3.76% | 25,204 | 0.90% | 0 | 0.00% |
| No | 12,084,156 | 5,589,057 | 46.25% | 3,617,249 | 29.93% | 1,799,429 | 14.89% | 740,042 | 6.12% | 281,458 | 2.33% | 37,032 | 0.31% | 19,889 | 0.16% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 6,738,730 | 45.30% | 4,693,232 | 31.55% | 2,122,591 | 14.27% | 852,436 | 5.73% | 386,354 | 2.60% | 62,237 | 0.42% | 19,889 | 0.13% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 9,678,148 | 4,552,934 | 47.04% | 3,030,971 | 31.32% | 1,108,555 | 11.45% | 603,276 | 6.23% | 333,252 | 3.44% | 49,159 | 0.51% | 0 | 0.00% |
| No | 5,159,491 | 2,147,966 | 41.63% | 1,662,261 | 32.22% | 1,014,036 | 19.65% | 249,159 | 4.83% | 53,102 | 1.03% | 13,077 | 0.25% | 19,889 | 0.39% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 7,126,875 | 3,021,137 | 42.39% | 2,593,918 | 36.40% | 938,335 | 13.17% | 414,096 | 5.81% | 110,052 | 1.54% | 49,338 | 0.69% | 0 | 0.00% |
| No | 7,748,593 | 3,717,592 | 47.98% | 2,099,314 | 27.09% | 1,184,256 | 15.28% | 438,340 | 5.66% | 276,302 | 3.57% | 12,899 | 0.17% | 19,889 | 0.26% |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,826,339 | 1,146,603 | 40.57% | 845,257 | 29.91% | 424,592 | 15.02% | 240,264 | 8.50% | 120,464 | 4.26% | 49,159 | 1.74% | 0 | 0.00% |
| Mobility limitations | 2,377,127 | 927,405 | 39.01% | 816,318 | 34.34% | 270,725 | 11.39% | 296,900 | 12.49% | 39,019 | 1.64% | 6,869 | 0.29% | 19,889 | 0.84% |
| No mobility limitations | 3,828,510 | 1,697,152 | 44.33% | 1,370,922 | 35.81% | 511,840 | 13.37% | 98,471 | 2.57% | 143,917 | 3.76% | 6,208 | 0.16% | 0 | 0.00% |
| Refused/Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| N/A - Respondent not asked mob. lim. questions | 5,805,663 | 2,929,740 | 50.46% | 1,660,735 | 28.61% | 915,434 | 15.77% | 216,800 | 3.73% | 82,953 | 1.43% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,782,327 | 736,242 | 41.31% | 502,749 | 28.21% | 254,550 | 14.28% | 160,361 | 9.00% | 78,373 | 4.40% | 30,163 | 1.69% | 19,889 | 1.12% |
| No | 13,063,306 | 6,002,488 | 45.95% | 4,190,483 | 32.08% | 1,868,041 | 14.30% | 662,239 | 5.07% | 307,981 | 2.36% | 32,074 | 0.25% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 6,738,730 | 45.30% | 4,693,232 | 31.55% | 2,122,591 | 14.27% | 852,436 | 5.73% | 386,354 | 2.60% | 62,237 | 0.42% | 19,889 | 0.13% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 1,880,197 | 960,637 | 51.09% | 468,030 | 24.89% | 212,901 | 11.32% | 114,394 | 6.08% | 124,235 | 6.61% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,620,551 | 699,681 | 43.18% | 778,495 | 48.04% | 98,924 | 6.10% | 43,452 | 2.68% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,385,683 | 500,033 | 36.09% | 456,934 | 32.98% | 216,896 | 15.65% | 160,026 | 11.55% | 51,795 | 3.74% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 844,703 | 436,643 | 51.69% | 294,714 | 34.89% | 113,346 | 13.42% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 1-3 times a month | 2,186,284 | 1,012,785 | 46.32% | 674,551 | 30.85% | 270,449 | 12.37% | 207,879 | 9.51% | 7,722 | 0.35% | 12,899 | 0.59% | 0 | 0.00% |
| Never | 6,823,374 | 3,099,826 | 45.43% | 2,008,261 | 29.43% | 1,172,921 | 17.19% | 326,686 | 4.79% | 202,603 | 2.97% | 13,077 | 0.19% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 566,107 | 381,051 | 67.31% | 18,144 | 3.21% | 130,443 | 23.04% | 36,468 | 6.44% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 14,035,507 | 6,248,899 | 44.52% | 4,603,316 | 32.80% | 1,954,994 | 13.93% | 815,968 | 5.81% | 386,354 | 2.75% | 25,976 | 0.19% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 827,931 | 474,953 | 57.37% | 193,021 | 23.31% | 124,290 | 15.01% | 35,666 | 4.31% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,431,660 | 739,246 | 51.64% | 558,958 | 39.04% | 62,575 | 4.37% | 70,881 | 4.95% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,979,005 | 1,869,663 | 62.76% | 706,509 | 23.72% | 263,824 | 8.86% | 70,967 | 2.38% | 68,041 | 2.28% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,262,484 | 1,489,318 | 45.65% | 902,784 | 27.67% | 477,848 | 14.65% | 235,656 | 7.22% | 156,879 | 4.81% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,986,235 | 1,110,434 | 55.91% | 678,293 | 34.15% | 120,652 | 6.07% | 17,821 | 0.90% | 59,036 | 2.97% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,488,541 | 664,856 | 44.66% | 551,947 | 37.08% | 185,717 | 12.48% | 41,151 | 2.76% | 44,871 | 3.01% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,114,946 | 486,715 | 43.65% | 385,096 | 34.54% | 76,246 | 6.84% | 132,827 | 11.91% | 34,061 | 3.05% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 976,453 | 477,947 | 48.95% | 375,937 | 38.50% | 44,578 | 4.57% | 61,041 | 6.25% | 16,951 | 1.74% | 0 | 0.00% | 0 | 0.00% |
| 80+ | 1,366,355 | 434,192 | 31.78% | 636,083 | 46.55% | 228,095 | 16.69% | 44,280 | 3.24% | 23,705 | 1.73% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,243,153 | 759,418 | 61.09% | 339,840 | 27.34% | 61,138 | 4.92% | 48,224 | 3.88% | 34,534 | 2.78% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 611,745 | 465,210 | 76.05% | 85,028 | 13.90% | 29,243 | 4.78% | 19,774 | 3.23% | 12,491 | 2.04% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 10,871,907 | 4,960,497 | 45.63% | 3,795,944 | 34.92% | 1,238,014 | 11.39% | 576,484 | 5.30% | 300,968 | 2.77% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic Black | 2,146,928 | 1,106,834 | 51.55% | 682,581 | 31.79% | 234,292 | 10.91% | 67,671 | 3.15% | 55,550 | 2.59% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | 601,568 | 476,474 | 79.21% | 85,237 | 14.17% | 21,138 | 3.51% | 18,720 | 3.11% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 300,834 | 45,747 | 15.21% | 132,210 | 43.95% | 87,610 | 29.12% | 0 | 0.00% | 35,267 | 11.72% | 0 | 0.00% | 0 | 0.00% |
| Less Than 9th Grade | 1,222,885 | 713,181 | 58.32% | 321,438 | 26.29% | 109,573 | 8.96% | 40,822 | 3.34% | 37,870 | 3.10% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,874,404 | 990,082 | 52.82% | 538,756 | 28.74% | 141,375 | 7.54% | 153,071 | 8.17% | 51,120 | 2.73% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 12,077,179 | 6,019,423 | 49.84% | 3,996,224 | 33.09% | 1,245,267 | 10.31% | 536,980 | 4.45% | 279,285 | 2.31% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 10,682,851 | 5,487,638 | 51.37% | 3,467,681 | 32.46% | 1,102,267 | 10.32% | 343,609 | 3.22% | 281,656 | 2.64% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 569,758 | 221,799 | 38.93% | 260,218 | 45.67% | 87,741 | 15.40% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,496,763 | 702,659 | 46.95% | 364,111 | 24.33% | 222,838 | 14.89% | 116,019 | 7.75% | 91,136 | 6.09% | 0 | 0.00% | 0 | 0.00% |
| Separated | 210,011 | 136,255 | 64.88% | 47,522 | 22.63% | 0 | 0.00% | 26,234 | 12.49% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 1,602,517 | 798,258 | 49.81% | 592,212 | 36.96% | 70,848 | 4.42% | 110,448 | 6.89% | 30,751 | 1.92% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 810,924 | 372,524 | 45.94% | 239,839 | 29.58% | 100,131 | 12.35% | 98,431 | 12.14% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,455,121 | 814,490 | 55.97% | 443,703 | 30.49% | 159,224 | 10.94% | 25,213 | 1.73% | 12,491 | 0.86% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,596,007 | 765,211 | 47.95% | 537,396 | 33.67% | 137,663 | 8.63% | 122,530 | 7.68% | 33,207 | 2.08% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 12,424,174 | 6,188,732 | 49.81% | 4,007,529 | 32.26% | 1,286,937 | 10.36% | 583,131 | 4.69% | 357,845 | 2.88% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 292,076 | 156,700 | 53.65% | 111,671 | 38.23% | 0 | 0.00% | 0 | 0.00% | 23,705 | 8.12% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 4,066,463 | 1,936,994 | 47.63% | 1,499,553 | 36.88% | 281,438 | 6.92% | 193,242 | 4.75% | 155,236 | 3.82% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 4,788,091 | 2,636,180 | 55.06% | 1,360,417 | 28.41% | 513,607 | 10.73% | 222,019 | 4.64% | 55,868 | 1.17% | 0 | 0.00% | 0 | 0.00% |
| Obese: 30 or higher | 6,328,671 | 3,038,558 | 48.01% | 2,016,988 | 31.87% | 788,781 | 12.46% | 315,612 | 4.99% | 168,733 | 2.67% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Glycohemoglobin < 5.7% | 8,629,123 | 4,484,564 | 51.97% | 2,726,278 | 31.59% | 803,674 | 9.31% | 416,956 | 4.83% | 197,651 | 2.29% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 4,513,678 | 2,059,672 | 45.63% | 1,504,385 | 33.33% | 533,679 | 11.82% | 229,348 | 5.08% | 186,594 | 4.13% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 2,163,331 | 1,147,153 | 53.03% | 696,236 | 32.18% | 216,075 | 9.99% | 84,569 | 3.91% | 19,298 | 0.89% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,480,826 | 1,096,987 | 44.22% | 978,189 | 39.43% | 280,213 | 11.30% | 84,569 | 3.41% | 40,869 | 1.65% | 0 | 0.00% | 0 | 0.00% |
| No | 12,994,476 | 6,671,446 | 51.34% | 4,010,440 | 30.86% | 1,303,612 | 10.03% | 646,304 | 4.97% | 362,674 | 2.79% | 0 | 0.00% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 8,875,386 | 4,486,987 | 50.56% | 2,725,635 | 30.71% | 939,691 | 10.59% | 529,462 | 5.97% | 193,611 | 2.18% | 0 | 0.00% | 0 | 0.00% |
| No | 6,599,916 | 3,281,446 | 49.72% | 2,262,993 | 34.29% | 644,134 | 9.76% | 201,411 | 3.05% | 209,932 | 3.18% | 0 | 0.00% | 0 | 0.00% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 6,397,998 | 3,004,142 | 46.95% | 1,963,403 | 30.69% | 760,448 | 11.89% | 313,486 | 4.90% | 356,518 | 5.57% | 0 | 0.00% | 0 | 0.00% |
| No | 9,066,420 | 4,753,407 | 52.43% | 3,025,226 | 33.37% | 823,377 | 9.08% | 417,387 | 4.60% | 47,024 | 0.52% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,512,390 | 1,288,888 | 51.30% | 714,797 | 28.45% | 326,811 | 13.01% | 97,891 | 3.90% | 84,003 | 3.34% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 2,045,907 | 651,644 | 31.85% | 757,583 | 37.03% | 370,990 | 18.13% | 217,932 | 10.65% | 47,758 | 2.33% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 4,531,815 | 2,041,143 | 45.04% | 1,724,000 | 38.04% | 468,409 | 10.34% | 162,752 | 3.59% | 135,511 | 2.99% | 0 | 0.00% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 6,385,189 | 3,786,757 | 59.31% | 1,792,248 | 28.07% | 417,614 | 6.54% | 252,298 | 3.95% | 136,271 | 2.13% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,963,555 | 670,672 | 34.16% | 502,300 | 25.58% | 514,962 | 26.23% | 123,731 | 6.30% | 151,891 | 7.74% | 0 | 0.00% | 0 | 0.00% |
| No | 13,504,920 | 7,097,761 | 52.56% | 4,479,503 | 33.17% | 1,068,862 | 7.91% | 607,143 | 4.50% | 251,652 | 1.86% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).

Table M06 - Level of bother among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 1,691,877 | 873,422 | 51.62% | 569,880 | 33.68% | 119,439 | 7.06% | 84,265 | 4.98% | 44,871 | 2.65% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,484,183 | 655,196 | 44.15% | 476,291 | 32.09% | 327,661 | 22.08% | 25,035 | 1.69% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,294,731 | 596,129 | 46.04% | 450,420 | 34.79% | 162,188 | 12.53% | 85,995 | 6.64% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 1,010,064 | 425,149 | 42.09% | 291,384 | 28.85% | 80,599 | 7.98% | 76,661 | 7.59% | 136,271 | 13.49% | 0 | 0.00% | 0 | 0.00% |
| 1-3 times a month | 2,725,838 | 1,363,538 | 50.02% | 976,040 | 35.81% | 236,295 | 8.67% | 104,555 | 3.84% | 45,411 | 1.67% | 0 | 0.00% | 0 | 0.00% |
| Never | 7,013,931 | 3,676,147 | 52.41% | 2,204,120 | 31.42% | 602,311 | 8.59% | 354,363 | 5.05% | 176,990 | 2.52% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 466,082 | 75,241 | 16.14% | 129,711 | 27.83% | 129,491 | 27.78% | 79,080 | 16.97% | 52,558 | 11.28% | 0 | 0.00% | 0 | 0.00% |
| Never | 14,575,578 | 7,493,946 | 51.41% | 4,712,187 | 32.33% | 1,399,001 | 9.60% | 619,460 | 4.25% | 350,984 | 2.41% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on positive responses to KIQ.042 and KIQ.044 (stress urinary incontinence and urge urinary incontinence).