

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 10,266,409 | 7,469,647 | 72.76% | 1,651,809 | 16.09% | 638,454 | 6.22% | 294,310 | 2.87% | 106,897 | 1.04% | 10,036 | 0.10% | 95,257 | 0.93% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 30-39 | 1,245,096 | 1,132,677 | 90.97% | 24,717 | 1.99% | 57,189 | 4.59% | 0 | 0.00% | 30,513 | 2.45% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 1,282,331 | 984,022 | 76.74% | 176,661 | 13.78% | 26,391 | 2.06% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 95,257 | 7.43% |
| 50-59 | 2,024,486 | 1,508,492 | 74.51% | 252,104 | 12.45% | 108,956 | 5.38% | 140,066 | 6.92% | 14,868 | 0.73% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 929,881 | 637,633 | 68.57% | 236,734 | 25.46% | 53,203 | 5.72% | 2,311 | 0.25% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,080,369 | 894,871 | 82.83% | 87,432 | 8.09% | 32,347 | 2.99% | 65,719 | 6.08% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,373,568 | 892,810 | 65.00% | 336,134 | 24.47% | 72,790 | 5.30% | 47,668 | 3.47% | 14,128 | 1.03% | 10,036 | 0.73% | 0 | 0.00% |
| 75-79 | 758,372 | 430,076 | 56.71% | 182,814 | 24.11% | 112,961 | 14.90% | 32,521 | 4.29% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 624,458 | 420,713 | 67.37% | 131,013 | 20.98% | 69,432 | 11.12% | 3,301 | 0.53% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 85+ | 566,783 | 296,204 | 52.26% | 115,283 | 20.34% | 105,185 | 18.56% | 2,723 | 0.48% | 47,387 | 8.36% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 475,436 | 360,296 | 75.78% | 60,298 | 12.68% | 35,078 | 7.38% | 19,764 | 4.16% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 553,379 | 359,074 | 64.89% | 64,807 | 11.71% | 54,412 | 9.83% | 75,086 | 13.57% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 7,730,812 | 5,823,253 | 75.33% | 1,126,977 | 14.58% | 438,478 | 5.67% | 199,460 | 2.58% | 47,387 | 0.61% | 0 | 0.00% | 95,257 | 1.23% |
| Non-Hispanic Black | 1,311,346 | 794,605 | 60.59% | 336,708 | 25.68% | 110,486 | 8.43% | 0 | 0.00% | 59,510 | 4.54% | 10,036 | 0.77% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,274,348 | 706,613 | 55.45% | 357,420 | 28.05% | 160,804 | 12.62% | 24,664 | 1.94% | 14,812 | 1.16% | 10,036 | 0.79% | 0 | 0.00% |
| 9-11th Grade | 1,393,333 | 899,935 | 64.59% | 204,116 | 14.65% | 164,423 | 11.80% | 96,331 | 6.91% | 28,528 | 2.05% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 7,469,140 | 5,792,907 | 77.56% | 1,063,224 | 14.23% | 280,880 | 3.76% | 173,314 | 2.32% | 63,557 | 0.85% | 0 | 0.00% | 95,257 | 1.28% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 6,617,196 | 4,826,571 | 72.94% | 1,009,262 | 15.25% | 453,125 | 6.85% | 235,000 | 3.55% | 93,237 | 1.41% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 861,620 | 476,645 | 55.32% | 264,333 | 30.68% | 77,270 | 8.97% | 29,712 | 3.45% | 13,660 | 1.59% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,262,656 | 1,001,189 | 79.29% | 135,613 | 10.74% | 28,286 | 2.24% | 2,311 | 0.18% | 0 | 0.00% | 0 | 0.00% | 95,257 | 7.54% |
| Separated | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Never Married | 983,139 | 775,063 | 78.84% | 93,033 | 9.46% | 79,772 | 8.11% | 25,235 | 2.57% | 0 | 0.00% | 10,036 | 1.02% | 0 | 0.00% |
| Living with Partner | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 593,049 | 456,933 | 77.05% | 8,855 | 1.49% | 26,391 | 4.45% | 5,612 | 0.95% | 0 | 0.00% | 0 | 0.00% | 95,257 | 16.06% |
| Poverty income ratio less than 1 | 1,516,688 | 1,056,205 | 69.64% | 228,990 | 15.10% | 89,373 | 5.89% | 132,084 | 8.71% | 0 | 0.00% | 10,036 | 0.66% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 8,156,673 | 5,956,509 | 73.03% | 1,413,963 | 17.34% | 522,689 | 6.41% | 156,614 | 1.92% | 106,897 | 1.31% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 874,992 | 450,096 | 51.44% | 133,690 | 15.28% | 179,736 | 20.54% | 71,924 | 8.22% | 39,546 | 4.52% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 2,739,084 | 1,888,657 | 68.95% | 517,365 | 18.89% | 188,770 | 6.89% | 34,224 | 1.25% | 14,812 | 0.54% | 0 | 0.00% | 95,257 | 3.48% |
| Overweight: 25.0 - 29.9 | 3,551,109 | 2,648,817 | 74.59% | 621,264 | 17.49% | 209,385 | 5.90% | 16,966 | 0.48% | 44,642 | 1.26% | 10,036 | 0.28% | 0 | 0.00% |
| Obese: 30 or higher | 3,101,225 | 2,482,077 | 80.04% | 379,490 | 12.24% | 60,563 | 1.95% | 171,197 | 5.52% | 7,898 | 0.25% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 416,345 | 280,364 | 67.34% | 73,784 | 17.72% | 24,713 | 5.94% | 0 | 0.00% | 37,484 | 9.00% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 6,339,137 | 4,758,538 | 75.07% | 929,502 | 14.66% | 406,741 | 6.42% | 135,439 | 2.14% | 13,660 | 0.22% | 0 | 0.00% | 95,257 | 1.50% |
| Glycohemoglobin 5.7% - 6.4% | 1,757,161 | 1,144,537 | 65.14% | 332,192 | 18.91% | 173,996 | 9.90% | 64,811 | 3.69% | 41,625 | 2.37% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,753,766 | 1,286,207 | 73.34% | 316,330 | 18.04% | 33,004 | 1.88% | 94,060 | 5.36% | 14,128 | 0.81% | 10,036 | 0.57% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,695,100 | 1,181,622 | 69.71% | 339,008 | 20.00% | 82,708 | 4.88% | 81,726 | 4.82% | 0 | 0.00% | 10,036 | 0.59% | 0 | 0.00% |
| No | 8,571,309 | 6,288,024 | 73.36% | 1,312,801 | 15.32% | 555,746 | 6.48% | 212,584 | 2.48% | 106,897 | 1.25% | 0 | 0.00% | 95,257 | 1.11% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2001 - 2002

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 7,768,433 | 50.20% | 4,988,629 | 32.24% | 1,583,825 | 10.23% | 730,873 | 4.72% | 403,543 | 2.61% | 0 | 0.00% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 7,293,348 | 5,301,765 | 72.69% | 1,127,949 | 15.47% | 500,444 | 6.86% | 253,228 | 3.47% | 99,926 | 1.37% | 10,036 | 0.14% | 0 | 0.00% |
| No | 2,973,061 | 2,167,882 | 72.92% | 523,859 | 17.62% | 138,010 | 4.64% | 41,082 | 1.38% | 6,971 | 0.23% | 0 | 0.00% | 95,257 | 3.20% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Yes | 3,876,071 | 2,369,289 | 61.13% | 887,664 | 22.90% | 297,002 | 7.66% | 286,194 | 7.38% | 25,887 | 0.67% | 10,036 | 0.26% | 0 | 0.00% |
| No | 6,380,301 | 5,090,322 | 79.78% | 764,145 | 11.98% | 341,452 | 5.35% | 8,116 | 0.13% | 81,011 | 1.27% | 0 | 0.00% | 95,257 | 1.49% |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,480,096 | 833,080 | 56.29% | 360,840 | 24.38% | 31,999 | 2.16% | 211,577 | 14.29% | 42,600 | 2.88% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 2,270,983 | 1,414,949 | 62.31% | 412,361 | 18.16% | 329,208 | 14.50% | 80,681 | 3.55% | 33,784 | 1.49% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 3,383,951 | 2,669,419 | 78.88% | 484,979 | 14.33% | 217,466 | 6.43% | 2,051 | 0.06% | 0 | 0.00% | 10,036 | 0.30% | 0 | 0.00% |
| Refused/Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| N/A - Respondent not asked mobility limitation questions | 3,116,764 | 2,537,584 | 81.42% | 393,629 | 12.63% | 59,781 | 1.92% | 0 | 0.00% | 30,513 | 0.98% | 0 | 0.00% | 95,257 | 3.06% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 7,469,647 | 72.76% | 1,651,809 | 16.09% | 638,454 | 6.22% | 294,310 | 2.87% | 106,897 | 1.04% | 10,036 | 0.10% | 95,257 | 0.93% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 7,469,647 | 72.76% | 1,651,809 | 16.09% | 638,454 | 6.22% | 294,310 | 2.87% | 106,897 | 1.04% | 10,036 | 0.10% | 95,257 | 0.93% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 10,266,409 | 7,469,647 | 72.76% | 1,651,809 | 16.09% | 638,454 | 6.22% | 294,310 | 2.87% | 106,897 | 1.04% | 10,036 | 0.10% | 95,257 | 0.93% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 20-29 | 920,444 | 681,239 | 74.01% | 97,282 | 10.57% | 111,444 | 12.11% | 6,188 | 0.67% | 24,290 | 2.64% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,961,991 | 1,572,852 | 80.17% | 222,024 | 11.32% | 130,605 | 6.66% | 36,511 | 1.86% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 1,948,077 | 1,643,804 | 84.38% | 250,930 | 12.88% | 43,977 | 2.26% | 0 | 0.00% | 9,367 | 0.48% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 2,843,199 | 2,234,120 | 78.58% | 390,942 | 13.75% | 69,799 | 2.45% | 111,222 | 3.91% | 37,116 | 1.31% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,000,574 | 766,073 | 76.56% | 81,957 | 8.19% | 74,880 | 7.48% | 77,664 | 7.76% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,335,420 | 1,162,182 | 87.03% | 46,369 | 3.47% | 98,257 | 7.36% | 0 | 0.00% | 28,612 | 2.14% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,398,131 | 1,047,081 | 74.89% | 102,651 | 7.34% | 181,625 | 12.99% | 16,536 | 1.18% | 50,239 | 3.59% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 1,080,370 | 695,212 | 64.35% | 317,775 | 29.41% | 55,302 | 5.12% | 12,082 | 1.12% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 662,074 | 510,433 | 77.10% | 43,285 | 6.54% | 69,848 | 10.55% | 18,004 | 2.72% | 20,504 | 3.10% | 0 | 0.00% | 0 | 0.00% |
| 85+ | 552,328 | 417,178 | 75.53% | 53,256 | 9.64% | 20,507 | 3.71% | 31,206 | 5.65% | 30,181 | 5.46% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 987,200 | 686,173 | 69.51% | 163,566 | 16.57% | 94,246 | 9.55% | 13,917 | 1.41% | 29,297 | 2.97% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 501,938 | 434,856 | 86.64% | 0 | 0.00% | 15,782 | 3.14% | 0 | 0.00% | 51,300 | 10.22% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 9,809,316 | 7,755,260 | 79.06% | 1,177,188 | 12.00% | 531,490 | 5.42% | 237,519 | 2.42% | 107,858 | 1.10% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic Black | 2,024,497 | 1,598,651 | 78.97% | 191,865 | 9.48% | 164,151 | 8.11% | 57,976 | 2.86% | 11,853 | 0.59% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,425,778 | 936,363 | 65.67% | 167,551 | 11.75% | 198,840 | 13.95% | 99,691 | 6.99% | 23,334 | 1.64% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,149,203 | 798,965 | 69.52% | 209,830 | 18.26% | 47,748 | 4.15% | 57,199 | 4.98% | 35,459 | 3.09% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 10,919,458 | 8,924,574 | 81.73% | 1,216,241 | 11.14% | 508,896 | 4.66% | 152,522 | 1.40% | 117,226 | 1.07% | 0 | 0.00% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 8,855,567 | 7,018,439 | 79.25% | 1,145,660 | 12.94% | 360,499 | 4.07% | 160,531 | 1.81% | 170,438 | 1.92% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 869,715 | 601,667 | 69.18% | 76,019 | 8.74% | 128,267 | 14.75% | 43,258 | 4.97% | 20,504 | 2.36% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,217,057 | 897,070 | 73.71% | 148,610 | 12.21% | 171,376 | 14.08% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Separated | 334,445 | 256,127 | 76.58% | 0 | 0.00% | 28,887 | 8.64% | 40,063 | 11.98% | 9,367 | 2.80% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 1,587,594 | 1,301,008 | 81.95% | 157,900 | 9.95% | 69,314 | 4.37% | 59,371 | 3.74% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 622,715 | 460,833 | 74.00% | 63,981 | 10.27% | 97,901 | 15.72% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 572,832 | 344,704 | 60.18% | 94,398 | 16.48% | 54,681 | 9.55% | 0 | 0.00% | 79,049 | 13.80% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,153,158 | 702,369 | 60.91% | 277,677 | 24.08% | 79,992 | 6.94% | 68,831 | 5.97% | 24,290 | 2.11% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,976,617 | 9,683,100 | 80.85% | 1,234,395 | 10.31% | 721,571 | 6.02% | 240,581 | 2.01% | 96,970 | 0.81% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 268,019 | 74,444 | 27.78% | 0 | 0.00% | 108,106 | 40.34% | 35,230 | 13.14% | 50,239 | 18.74% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 3,529,345 | 2,614,589 | 74.08% | 571,076 | 16.18% | 206,647 | 5.86% | 31,758 | 0.90% | 105,276 | 2.98% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 4,301,128 | 3,646,133 | 84.77% | 415,598 | 9.66% | 219,248 | 5.10% | 20,149 | 0.47% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Obese: 30 or higher | 5,604,114 | 4,395,007 | 78.42% | 619,795 | 11.06% | 322,243 | 5.75% | 222,276 | 3.97% | 44,794 | 0.80% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 361,588 | 301,198 | 83.30% | 34,732 | 9.61% | 25,658 | 7.10% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 8,890,742 | 7,098,345 | 79.84% | 956,024 | 10.75% | 583,165 | 6.56% | 142,248 | 1.60% | 110,960 | 1.25% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 2,728,162 | 1,945,548 | 71.31% | 489,308 | 17.94% | 156,512 | 5.74% | 97,964 | 3.59% | 38,831 | 1.42% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,722,115 | 1,385,082 | 80.43% | 126,406 | 7.34% | 90,909 | 5.28% | 69,200 | 4.02% | 50,518 | 2.93% | 0 | 0.00% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,199,006 | 1,461,094 | 66.44% | 249,392 | 11.34% | 218,165 | 9.92% | 121,346 | 5.52% | 149,009 | 6.78% | 0 | 0.00% | 0 | 0.00% |
| No | 11,503,602 | 9,269,078 | 80.58% | 1,357,077 | 11.80% | 638,079 | 5.55% | 188,067 | 1.63% | 51,300 | 0.45% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2003 - 2004

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 8,509,424 | 6,601,720 | 77.58% | 892,756 | 10.49% | 603,764 | 7.10% | 215,881 | 2.54% | 195,302 | 2.30% | 0 | 0.00% | 0 | 0.00% |
| No | 5,181,352 | 4,128,453 | 79.68% | 713,713 | 13.77% | 252,480 | 4.87% | 81,700 | 1.58% | 5,007 | 0.10% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Yes | 6,058,883 | 4,417,929 | 72.92% | 891,075 | 14.71% | 462,477 | 7.63% | 137,333 | 2.27% | 150,070 | 2.48% | 0 | 0.00% | 0 | 0.00% |
| No | 7,560,067 | 6,249,092 | 82.66% | 694,890 | 9.19% | 393,767 | 5.21% | 172,080 | 2.28% | 50,239 | 0.66% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,964,807 | 1,819,508 | 61.37% | 347,734 | 11.73% | 461,811 | 15.58% | 201,173 | 6.79% | 134,581 | 4.54% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 1,853,409 | 1,454,402 | 78.47% | 220,150 | 11.88% | 55,670 | 3.00% | 62,466 | 3.37% | 60,721 | 3.28% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 3,470,630 | 2,958,077 | 85.23% | 315,610 | 9.09% | 177,090 | 5.10% | 14,847 | 0.43% | 5,007 | 0.14% | 0 | 0.00% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 5,413,761 | 4,498,187 | 83.09% | 722,975 | 13.35% | 161,674 | 2.99% | 30,926 | 0.57% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| N/A for these years of data | 13,702,607 | 10,730,173 | 78.31% | 1,606,469 | 11.72% | 856,244 | 6.25% | 309,412 | 2.26% | 200,309 | 1.46% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 12,078,665 | 85.53% | 1,314,584 | 9.31% | 627,453 | 4.44% | 38,185 | 0.27% | 63,861 | 0.45% | 0 | 0.00% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 899,897 | 801,411 | 89.06% | 9,413 | 1.05% | 25,212 | 2.80% | 0 | 0.00% | 63,861 | 7.10% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,031,971 | 888,163 | 86.06% | 59,970 | 5.81% | 83,838 | 8.12% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,978,059 | 2,603,755 | 87.43% | 165,536 | 5.56% | 208,768 | 7.01% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,614,824 | 3,136,489 | 86.77% | 364,218 | 10.08% | 114,117 | 3.16% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,085,089 | 942,184 | 86.83% | 35,708 | 3.29% | 84,417 | 7.78% | 22,780 | 2.10% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,466,777 | 1,307,043 | 89.11% | 144,329 | 9.84% | 0 | 0.00% | 15,405 | 1.05% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,031,394 | 816,021 | 79.12% | 180,421 | 17.49% | 34,951 | 3.39% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 959,797 | 754,635 | 78.62% | 160,682 | 16.74% | 44,480 | 4.63% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 80-84 | 733,159 | 631,811 | 86.18% | 90,790 | 12.38% | 10,558 | 1.44% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 85+ | 299,135 | 197,154 | 65.91% | 80,869 | 27.03% | 21,112 | 7.06% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,116,500 | 824,212 | 73.82% | 180,975 | 16.21% | 82,950 | 7.43% | 28,362 | 2.54% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Non-Hispanic White | 10,764,351 | 9,363,206 | 86.98% | 933,269 | 8.67% | 428,786 | 3.98% | 0 | 0.00% | 39,090 | 0.36% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic Black | 1,520,452 | 1,341,586 | 88.24% | 86,572 | 5.69% | 57,701 | 3.79% | 9,823 | 0.65% | 24,771 | 1.63% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,110,708 | 906,437 | 81.61% | 194,775 | 17.54% | 0 | 0.00% | 9,496 | 0.85% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,514,077 | 1,290,470 | 85.23% | 95,900 | 6.33% | 99,019 | 6.54% | 28,689 | 1.89% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 11,429,497 | 9,860,443 | 86.27% | 980,665 | 8.58% | 524,528 | 4.59% | 0 | 0.00% | 63,861 | 0.56% | 0 | 0.00% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 12,078,665 | 85.53% | 1,314,584 | 9.31% | 627,453 | 4.44% | 38,185 | 0.27% | 63,861 | 0.45% | 0 | 0.00% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 9,694,781 | 8,300,832 | 85.62% | 913,394 | 9.42% | 452,193 | 4.66% | 28,362 | 0.29% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 577,904 | 421,489 | 72.93% | 146,592 | 25.37% | 0 | 0.00% | 9,823 | 1.70% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,797,233 | 1,569,540 | 87.33% | 41,182 | 2.29% | 147,421 | 8.20% | 0 | 0.00% | 39,090 | 2.18% | 0 | 0.00% | 0 | 0.00% |
| Separated | 472,734 | 396,666 | 83.91% | 76,068 | 16.09% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 999,423 | 885,101 | 88.56% | 89,551 | 8.96% | 0 | 0.00% | 0 | 0.00% | 24,771 | 2.48% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 437,370 | 413,438 | 94.53% | 0 | 0.00% | 23,932 | 5.47% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 811,536 | 673,986 | 83.05% | 93,071 | 11.47% | 44,480 | 5.48% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,447,959 | 1,109,767 | 76.64% | 210,386 | 14.53% | 99,445 | 6.87% | 28,362 | 1.96% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,863,253 | 10,294,913 | 86.78% | 1,011,128 | 8.52% | 483,529 | 4.08% | 9,823 | 0.08% | 63,861 | 0.54% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Normal: less than 25 | 3,052,898 | 2,639,435 | 86.46% | 258,420 | 8.46% | 151,129 | 4.95% | 3,914 | 0.13% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 5,408,869 | 4,644,095 | 85.86% | 467,913 | 8.65% | 257,772 | 4.77% | 0 | 0.00% | 39,090 | 0.72% | 0 | 0.00% | 0 | 0.00% |
| Obese: 30 or higher | 5,534,849 | 4,724,229 | 85.35% | 554,136 | 10.01% | 197,441 | 3.57% | 34,271 | 0.62% | 24,771 | 0.45% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 300,823 | 219,033 | 72.81% | 31,728 | 10.55% | 44,480 | 14.79% | 5,582 | 1.86% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin < 5.7% | 9,038,263 | 7,751,230 | 85.76% | 862,907 | 9.55% | 395,441 | 4.38% | 3,914 | 0.04% | 24,771 | 0.27% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 3,182,425 | 2,798,452 | 87.93% | 322,104 | 10.12% | 22,779 | 0.72% | 0 | 0.00% | 39,090 | 1.23% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 1,601,238 | 1,309,950 | 81.81% | 97,845 | 6.11% | 164,754 | 10.29% | 28,689 | 1.79% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 12,078,665 | 85.53% | 1,314,584 | 9.31% | 627,453 | 4.44% | 38,185 | 0.27% | 63,861 | 0.45% | 0 | 0.00% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,702,435 | 1,341,645 | 78.81% | 145,449 | 8.54% | 160,847 | 9.45% | 15,405 | 0.90% | 39,090 | 2.30% | 0 | 0.00% | 0 | 0.00% |
| No | 12,420,313 | 10,737,020 | 86.45% | 1,169,135 | 9.41% | 466,606 | 3.76% | 22,780 | 0.18% | 24,771 | 0.20% | 0 | 0.00% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 9,608,565 | 8,047,395 | 83.75% | 942,626 | 9.81% | 541,270 | 5.63% | 38,185 | 0.40% | 39,090 | 0.41% | 0 | 0.00% | 0 | 0.00% |
| No | 4,514,183 | 4,031,270 | 89.30% | 371,958 | 8.24% | 86,183 | 1.91% | 0 | 0.00% | 24,771 | 0.55% | 0 | 0.00% | 0 | 0.00% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 6,581,507 | 5,504,244 | 83.63% | 718,251 | 10.91% | 300,603 | 4.57% | 19,319 | 0.29% | 39,090 | 0.59% | 0 | 0.00% | 0 | 0.00% |
| No | 7,484,663 | 6,574,421 | 87.84% | 596,333 | 7.97% | 270,271 | 3.61% | 18,866 | 0.25% | 24,771 | 0.33% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,617,259 | 980,439 | 60.62% | 293,177 | 18.13% | 276,191 | 17.08% | 28,362 | 1.75% | 39,090 | 2.42% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 2,018,270 | 1,646,199 | 81.56% | 257,180 | 12.74% | 105,069 | 5.21% | 9,823 | 0.49% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 4,434,930 | 4,005,850 | 90.32% | 419,941 | 9.47% | 9,138 | 0.21% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 6,052,290 | 5,446,178 | 89.99% | 344,285 | 5.69% | 237,055 | 3.92% | 0 | 0.00% | 24,771 | 0.41% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,079,229 | 1,700,493 | 81.78% | 153,563 | 7.39% | 221,259 | 10.64% | 3,914 | 0.19% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| No | 12,043,519 | 10,378,173 | 86.17% | 1,161,020 | 9.64% | 406,194 | 3.37% | 34,271 | 0.28% | 63,861 | 0.53% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2005 - 2006

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,122,748 | 12,078,665 | 85.53% | 1,314,584 | 9.31% | 627,453 | 4.44% | 38,185 | 0.27% | 63,861 | 0.45% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 2,904,617 | 2,456,545 | 84.57% | 192,592 | 6.63% | 212,476 | 7.32% | 3,914 | 0.13% | 39,090 | 1.35% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,026,938 | 873,951 | 85.10% | 147,755 | 14.39% | 5,231 | 0.51% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,451,505 | 1,110,655 | 76.52% | 217,242 | 14.97% | 123,608 | 8.52% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 1,065,210 | 915,148 | 85.91% | 93,097 | 8.74% | 47,142 | 4.43% | 9,823 | 0.92% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 1-3 times a month | 2,228,101 | 1,741,450 | 78.16% | 363,718 | 16.32% | 117,351 | 5.27% | 5,582 | 0.25% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 5,381,424 | 4,958,394 | 92.14% | 276,447 | 5.14% | 102,944 | 1.91% | 18,866 | 0.35% | 24,771 | 0.46% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 598,247 | 455,381 | 76.12% | 95,723 | 16.00% | 47,142 | 7.88% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 13,338,270 | 11,583,594 | 86.84% | 1,091,020 | 8.18% | 561,610 | 4.21% | 38,185 | 0.29% | 63,861 | 0.48% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 11,867,673 | 79.78% | 1,904,446 | 12.80% | 632,891 | 4.25% | 336,912 | 2.26% | 51,420 | 0.35% | 62,237 | 0.42% | 19,889 | 0.13% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 604,177 | 579,680 | 95.95% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 24,497 | 4.05% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,125,718 | 840,924 | 74.70% | 206,804 | 18.37% | 11,602 | 1.03% | 66,388 | 5.90% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,794,224 | 2,280,970 | 81.63% | 397,712 | 14.23% | 108,654 | 3.89% | 6,889 | 0.25% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,385,303 | 2,740,909 | 80.96% | 449,944 | 13.29% | 73,513 | 2.17% | 101,047 | 2.98% | 0 | 0.00% | 0 | 0.00% | 19,889 | 0.59% |
| 60-64 | 1,707,386 | 1,495,112 | 87.57% | 120,568 | 7.06% | 14,943 | 0.88% | 69,894 | 4.09% | 0 | 0.00% | 6,869 | 0.40% | 0 | 0.00% |
| 65-69 | 1,518,446 | 1,277,361 | 84.12% | 153,547 | 10.11% | 87,537 | 5.76% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,283,484 | 866,205 | 67.49% | 214,204 | 16.69% | 131,777 | 10.27% | 65,200 | 5.08% | 0 | 0.00% | 6,097 | 0.48% | 0 | 0.00% |
| 75-79 | 1,207,845 | 901,842 | 74.67% | 242,744 | 20.10% | 29,556 | 2.45% | 27,495 | 2.28% | 0 | 0.00% | 6,208 | 0.51% | 0 | 0.00% |
| 80+ | 1,233,178 | 884,670 | 71.74% | 103,216 | 8.37% | 175,307 | 14.22% | 0 | 0.00% | 26,923 | 2.18% | 43,062 | 3.49% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,208,868 | 936,270 | 77.45% | 193,976 | 16.05% | 31,853 | 2.63% | 16,064 | 1.33% | 24,497 | 2.03% | 6,208 | 0.51% | 0 | 0.00% |
| Other Hispanic | 624,985 | 416,034 | 66.57% | 105,860 | 16.94% | 8,188 | 1.31% | 88,806 | 14.21% | 0 | 0.00% | 6,097 | 0.98% | 0 | 0.00% |
| Non-Hispanic White | 10,135,298 | 8,366,996 | 82.55% | 1,093,608 | 10.79% | 442,280 | 4.36% | 175,329 | 1.73% | 26,923 | 0.27% | 30,163 | 0.30% | 0 | 0.00% |
| Non-Hispanic Black | 2,092,450 | 1,705,113 | 81.49% | 171,997 | 8.22% | 118,967 | 5.69% | 56,714 | 2.71% | 0 | 0.00% | 19,768 | 0.94% | 19,889 | 0.95% |
| Other Race - Including Multi-Racial | 813,867 | 443,260 | 54.46% | 339,005 | 41.65% | 31,602 | 3.88% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Less Than 9th Grade | 1,853,886 | 1,410,526 | 76.08% | 209,988 | 11.33% | 112,194 | 6.05% | 83,603 | 4.51% | 24,497 | 1.32% | 13,077 | 0.71% | 0 | 0.00% |
| 9-11th Grade | 2,135,076 | 1,739,054 | 81.45% | 249,955 | 11.71% | 75,053 | 3.52% | 51,125 | 2.39% | 0 | 0.00% | 0 | 0.00% | 19,889 | 0.93% |
| High School Grad/GED or Equivalent | 10,623,592 | 8,473,323 | 79.76% | 1,426,360 | 13.43% | 445,643 | 4.19% | 202,184 | 1.90% | 26,923 | 0.25% | 49,159 | 0.46% | 0 | 0.00% |
| Don't Know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 11,867,673 | 79.78% | 1,904,446 | 12.80% | 632,891 | 4.25% | 336,912 | 2.26% | 51,420 | 0.35% | 62,237 | 0.42% | 19,889 | 0.13% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 9,743,242 | 7,954,872 | 81.65% | 1,282,736 | 13.17% | 294,466 | 3.02% | 166,075 | 1.70% | 0 | 0.00% | 25,204 | 0.26% | 19,889 | 0.20% |
| Widowed | 884,467 | 620,009 | 70.10% | 87,579 | 9.90% | 155,494 | 17.58% | 14,516 | 1.64% | 0 | 0.00% | 6,869 | 0.78% | 0 | 0.00% |
| Divorced | 1,942,388 | 1,502,737 | 77.37% | 199,765 | 10.28% | 100,060 | 5.15% | 82,740 | 4.26% | 26,923 | 1.39% | 30,163 | 1.55% | 0 | 0.00% |
| Separated | 423,395 | 269,972 | 63.76% | 124,866 | 29.49% | 19,889 | 4.70% | 8,669 | 2.05% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 818,909 | 721,677 | 88.13% | 13,575 | 1.66% | 62,980 | 7.69% | 20,677 | 2.52% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 1,013,788 | 764,837 | 75.44% | 180,217 | 17.78% | 0 | 0.00% | 44,236 | 4.36% | 24,497 | 2.42% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,314,414 | 1,044,918 | 79.50% | 118,347 | 9.00% | 116,267 | 8.85% | 28,674 | 2.18% | 0 | 0.00% | 6,208 | 0.47% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,579,500 | 1,077,866 | 68.24% | 381,079 | 24.13% | 65,000 | 4.12% | 42,588 | 2.70% | 0 | 0.00% | 12,967 | 0.82% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 11,981,554 | 9,744,889 | 81.33% | 1,405,020 | 11.73% | 451,624 | 3.77% | 265,651 | 2.22% | 51,420 | 0.43% | 43,062 | 0.36% | 19,889 | 0.17% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 211,693 | 167,411 | 79.08% | 14,119 | 6.67% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 30,163 | 14.25% | 0 | 0.00% |
| Normal: less than 25 | 3,280,900 | 2,415,878 | 73.63% | 579,745 | 17.67% | 177,284 | 5.40% | 81,896 | 2.50% | 0 | 0.00% | 6,208 | 0.19% | 19,889 | 0.61% |
| Overweight: 25.0 - 29.9 | 5,264,464 | 4,353,735 | 82.70% | 637,448 | 12.11% | 172,268 | 3.27% | 94,916 | 1.80% | 0 | 0.00% | 6,097 | 0.12% | 0 | 0.00% |
| Obese: 30 or higher | 6,118,411 | 4,930,650 | 80.59% | 673,133 | 11.00% | 283,339 | 4.63% | 160,101 | 2.62% | 51,420 | 0.84% | 19,768 | 0.32% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 182,953 | 80,887 | 44.21% | 38,582 | 21.09% | 29,837 | 16.31% | 6,889 | 3.77% | 0 | 0.00% | 6,869 | 3.75% | 19,889 | 10.87% |
| Glycohemoglobin < 5.7% | 7,603,718 | 6,250,533 | 82.20% | 901,782 | 11.86% | 284,816 | 3.75% | 142,091 | 1.87% | 24,497 | 0.32% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 5,052,896 | 3,950,539 | 78.18% | 662,889 | 13.12% | 226,520 | 4.48% | 136,865 | 2.71% | 26,923 | 0.53% | 49,159 | 0.97% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 2,035,901 | 1,585,715 | 77.89% | 301,193 | 14.79% | 91,718 | 4.51% | 51,067 | 2.51% | 0 | 0.00% | 6,208 | 0.30% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,791,313 | 2,219,005 | 79.50% | 358,024 | 12.83% | 129,357 | 4.63% | 59,722 | 2.14% | 0 | 0.00% | 25,204 | 0.90% | 0 | 0.00% |
| No | 12,084,156 | 9,648,668 | 79.85% | 1,546,422 | 12.80% | 503,533 | 4.17% | 277,190 | 2.29% | 51,420 | 0.43% | 37,032 | 0.31% | 19,889 | 0.16% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 11,867,673 | 79.78% | 1,904,446 | 12.80% | 632,891 | 4.25% | 336,912 | 2.26% | 51,420 | 0.35% | 62,237 | 0.42% | 19,889 | 0.13% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 9,678,148 | 7,658,310 | 79.13% | 1,299,842 | 13.43% | 390,637 | 4.04% | 228,780 | 2.36% | 51,420 | 0.53% | 49,159 | 0.51% | 0 | 0.00% |
| No | 5,159,491 | 4,171,534 | 80.85% | 604,605 | 11.72% | 242,254 | 4.70% | 108,132 | 2.10% | 0 | 0.00% | 13,077 | 0.25% | 19,889 | 0.39% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 7,126,875 | 5,619,677 | 78.85% | 890,834 | 12.50% | 380,249 | 5.34% | 159,855 | 2.24% | 26,923 | 0.38% | 49,338 | 0.69% | 0 | 0.00% |
| No | 7,748,593 | 6,247,996 | 80.63% | 1,013,612 | 13.08% | 252,642 | 3.26% | 177,058 | 2.29% | 24,497 | 0.32% | 12,899 | 0.17% | 19,889 | 0.26% |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,826,339 | 1,958,633 | 69.30% | 480,226 | 16.99% | 125,577 | 4.44% | 185,822 | 6.57% | 26,923 | 0.95% | 49,159 | 1.74% | 0 | 0.00% |
| Mobility limitations | 2,377,127 | 1,962,229 | 82.55% | 100,947 | 4.25% | 201,858 | 8.49% | 85,334 | 3.59% | 0 | 0.00% | 6,869 | 0.29% | 19,889 | 0.84% |
| No mobility limitations | 3,828,510 | 3,026,437 | 79.05% | 603,008 | 15.75% | 192,857 | 5.04% | 0 | 0.00% | 0 | 0.00% | 6,208 | 0.16% | 0 | 0.00% |
| Refused/Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| N/A - Respondent not asked mobility limitation questions | 5,805,663 | 4,882,546 | 84.10% | 720,266 | 12.41% | 112,598 | 1.94% | 65,757 | 1.13% | 24,497 | 0.42% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,782,327 | 1,331,072 | 74.68% | 260,105 | 14.59% | 41,805 | 2.35% | 72,369 | 4.06% | 26,923 | 1.51% | 30,163 | 1.69% | 19,889 | 1.12% |
| No | 13,063,306 | 10,513,987 | 80.48% | 1,644,341 | 12.59% | 583,864 | 4.47% | 264,543 | 2.03% | 24,497 | 0.19% | 32,074 | 0.25% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2007 - 2008

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 14,875,469 | 11,867,673 | 79.78% | 1,904,446 | 12.80% | 632,891 | 4.25% | 336,912 | 2.26% | 51,420 | 0.35% | 62,237 | 0.42% | 19,889 | 0.13% |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 1,880,197 | 1,479,265 | 78.68% | 188,063 | 10.00% | 119,251 | 6.34% | 42,197 | 2.24% | 51,420 | 2.73% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,620,551 | 1,353,735 | 83.54% | 244,011 | 15.06% | 15,410 | 0.95% | 7,395 | 0.46% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,385,683 | 1,050,834 | 75.84% | 277,103 | 20.00% | 22,184 | 1.60% | 35,562 | 2.57% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 844,703 | 710,652 | 84.13% | 134,051 | 15.87% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 1-3 times a month | 2,186,284 | 1,435,409 | 65.66% | 547,789 | 25.06% | 138,696 | 6.34% | 51,491 | 2.36% | 0 | 0.00% | 12,899 | 0.59% | 0 | 0.00% |
| Never | 6,823,374 | 5,779,129 | 84.70% | 513,429 | 7.52% | 324,834 | 4.76% | 192,905 | 2.83% | 0 | 0.00% | 13,077 | 0.19% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 566,107 | 333,223 | 58.86% | 166,966 | 29.49% | 51,634 | 9.12% | 14,284 | 2.52% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 14,035,507 | 11,357,169 | 80.92% | 1,716,934 | 12.23% | 568,741 | 4.05% | 315,267 | 2.25% | 51,420 | 0.37% | 25,976 | 0.19% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-------------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 12,883,977 | 83.26% | 1,737,924 | 11.23% | 518,821 | 3.35% | 170,992 | 1.10% | 163,587 | 1.06% | 0 | 0.00% | 0 | 0.00% |
| Age at screening | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-19 | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 20-29 | 827,931 | 634,005 | 76.58% | 193,926 | 23.42% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 30-39 | 1,431,660 | 1,333,561 | 93.15% | 79,719 | 5.57% | 0 | 0.00% | 0 | 0.00% | 18,380 | 1.28% | 0 | 0.00% | 0 | 0.00% |
| 40-49 | 2,979,005 | 2,550,959 | 85.63% | 340,092 | 11.42% | 70,168 | 2.36% | 5,296 | 0.18% | 12,491 | 0.42% | 0 | 0.00% | 0 | 0.00% |
| 50-59 | 3,262,484 | 2,561,348 | 78.51% | 318,164 | 9.75% | 199,388 | 6.11% | 133,225 | 4.08% | 50,360 | 1.54% | 0 | 0.00% | 0 | 0.00% |
| 60-64 | 1,986,235 | 1,804,940 | 90.87% | 171,520 | 8.64% | 0 | 0.00% | 9,775 | 0.49% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 65-69 | 1,488,541 | 1,260,638 | 84.69% | 167,609 | 11.26% | 38,487 | 2.59% | 0 | 0.00% | 21,807 | 1.46% | 0 | 0.00% | 0 | 0.00% |
| 70-74 | 1,114,946 | 866,051 | 77.68% | 156,371 | 14.02% | 55,679 | 4.99% | 0 | 0.00% | 36,845 | 3.30% | 0 | 0.00% | 0 | 0.00% |
| 75-79 | 976,453 | 843,841 | 86.42% | 65,657 | 6.72% | 66,955 | 6.86% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 80+ | 1,366,355 | 1,028,635 | 75.28% | 203,175 | 14.87% | 88,145 | 6.45% | 22,696 | 1.66% | 23,705 | 1.73% | 0 | 0.00% | 0 | 0.00% |
| Race/ethnicity | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Mexican American | 1,243,153 | 1,018,891 | 81.96% | 157,459 | 12.67% | 11,147 | 0.90% | 5,296 | 0.43% | 50,360 | 4.05% | 0 | 0.00% | 0 | 0.00% |
| Other Hispanic | 611,745 | 488,195 | 79.80% | 89,677 | 14.66% | 21,382 | 3.50% | 0 | 0.00% | 12,491 | 2.04% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic White | 10,871,907 | 9,073,712 | 83.46% | 1,245,787 | 11.46% | 338,810 | 3.12% | 131,241 | 1.21% | 82,357 | 0.76% | 0 | 0.00% | 0 | 0.00% |
| Non-Hispanic Black | 2,146,928 | 1,786,846 | 83.23% | 159,765 | 7.44% | 147,482 | 6.87% | 34,456 | 1.60% | 18,380 | 0.86% | 0 | 0.00% | 0 | 0.00% |
| Other Race - Including Multi-Racial | 601,568 | 516,332 | 85.83% | 85,237 | 14.17% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Education | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 300,834 | 122,675 | 40.78% | 55,281 | 18.38% | 122,877 | 40.85% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Less Than 9th Grade | 1,222,885 | 801,308 | 65.53% | 355,698 | 29.09% | 38,151 | 3.12% | 0 | 0.00% | 27,726 | 2.27% | 0 | 0.00% | 0 | 0.00% |
| 9-11th Grade | 1,874,404 | 1,550,732 | 82.73% | 234,402 | 12.51% | 33,800 | 1.80% | 9,132 | 0.49% | 46,339 | 2.47% | 0 | 0.00% | 0 | 0.00% |
| High School Grad/GED or Equivalent | 12,077,179 | 10,409,261 | 86.19% | 1,092,542 | 9.05% | 323,993 | 2.68% | 161,861 | 1.34% | 89,522 | 0.74% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|-----------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 12,883,977 | 83.26% | 1,737,924 | 11.23% | 518,821 | 3.35% | 170,992 | 1.10% | 163,587 | 1.06% | 0 | 0.00% | 0 | 0.00% |
| Marital status | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Married | 10,682,851 | 8,841,788 | 82.77% | 1,225,128 | 11.47% | 411,641 | 3.85% | 140,372 | 1.31% | 63,922 | 0.60% | 0 | 0.00% | 0 | 0.00% |
| Widowed | 569,758 | 482,617 | 84.71% | 43,070 | 7.56% | 44,071 | 7.74% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Divorced | 1,496,763 | 1,176,435 | 78.60% | 203,775 | 13.61% | 35,267 | 2.36% | 0 | 0.00% | 81,285 | 5.43% | 0 | 0.00% | 0 | 0.00% |
| Separated | 210,011 | 179,158 | 85.31% | 9,775 | 4.65% | 21,078 | 10.04% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never Married | 1,602,517 | 1,405,486 | 87.70% | 187,256 | 11.69% | 0 | 0.00% | 9,775 | 0.61% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Living with Partner | 810,924 | 753,256 | 92.89% | 27,228 | 3.36% | 6,764 | 0.83% | 5,296 | 0.65% | 18,380 | 2.27% | 0 | 0.00% | 0 | 0.00% |
| Refused | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Poverty income ratio | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 1,455,121 | 1,216,141 | 83.58% | 141,997 | 9.76% | 84,492 | 5.81% | 0 | 0.00% | 12,491 | 0.86% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio less than 1 | 1,596,007 | 1,302,420 | 81.60% | 262,734 | 16.46% | 21,078 | 1.32% | 9,775 | 0.61% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Poverty income ratio 1 or greater | 12,424,174 | 10,365,415 | 83.43% | 1,333,193 | 10.73% | 413,252 | 3.33% | 161,217 | 1.30% | 151,097 | 1.22% | 0 | 0.00% | 0 | 0.00% |
| Body mass index | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 292,076 | 244,902 | 83.85% | 23,469 | 8.04% | 0 | 0.00% | 0 | 0.00% | 23,705 | 8.12% | 0 | 0.00% | 0 | 0.00% |
| Normal: less than 25 | 4,066,463 | 3,415,148 | 83.98% | 417,059 | 10.26% | 179,018 | 4.40% | 9,132 | 0.22% | 46,106 | 1.13% | 0 | 0.00% | 0 | 0.00% |
| Overweight: 25.0 - 29.9 | 4,788,091 | 4,107,673 | 85.79% | 556,222 | 11.62% | 69,918 | 1.46% | 32,471 | 0.68% | 21,807 | 0.46% | 0 | 0.00% | 0 | 0.00% |
| Obese: 30 or higher | 6,328,671 | 5,116,253 | 80.84% | 741,174 | 11.71% | 269,885 | 4.26% | 129,390 | 2.04% | 71,969 | 1.14% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Glycohemoglobin < 5.7% | 8,629,123 | 7,276,321 | 84.32% | 974,898 | 11.30% | 245,216 | 2.84% | 5,296 | 0.06% | 127,392 | 1.48% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 5.7% - 6.4% | 4,513,678 | 3,657,291 | 81.03% | 480,355 | 10.64% | 186,632 | 4.13% | 165,696 | 3.67% | 23,705 | 0.53% | 0 | 0.00% | 0 | 0.00% |
| Glycohemoglobin 6.5% or higher | 2,163,331 | 1,790,971 | 82.79% | 272,896 | 12.61% | 86,974 | 4.02% | 0 | 0.00% | 12,491 | 0.58% | 0 | 0.00% | 0 | 0.00% |
| Diabetes | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 2,480,826 | 2,155,750 | 86.90% | 250,720 | 10.11% | 74,355 | 3.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| No | 12,994,476 | 10,728,227 | 82.56% | 1,487,204 | 11.44% | 444,466 | 3.42% | 170,992 | 1.32% | 163,587 | 1.26% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|--|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 12,883,977 | 83.26% | 1,737,924 | 11.23% | 518,821 | 3.35% | 170,992 | 1.10% | 163,587 | 1.06% | 0 | 0.00% | 0 | 0.00% |
| Smoked at least 100 cigarettes in life | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 8,875,386 | 7,318,987 | 82.46% | 1,014,204 | 11.43% | 380,580 | 4.29% | 43,541 | 0.49% | 118,075 | 1.33% | 0 | 0.00% | 0 | 0.00% |
| No | 6,599,916 | 5,564,990 | 84.32% | 723,721 | 10.97% | 138,241 | 2.09% | 127,451 | 1.93% | 45,512 | 0.69% | 0 | 0.00% | 0 | 0.00% |
| High blood pressure | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 6,397,998 | 4,988,110 | 77.96% | 800,613 | 12.51% | 404,367 | 6.32% | 141,016 | 2.20% | 63,892 | 1.00% | 0 | 0.00% | 0 | 0.00% |
| No | 9,066,420 | 7,884,983 | 86.97% | 937,312 | 10.34% | 114,454 | 1.26% | 29,977 | 0.33% | 99,695 | 1.10% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Mobility limitation | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | 2,512,390 | 1,886,613 | 75.09% | 462,497 | 18.41% | 111,420 | 4.43% | 9,775 | 0.39% | 42,085 | 1.68% | 0 | 0.00% | 0 | 0.00% |
| Mobility limitations | 2,045,907 | 1,419,096 | 69.36% | 318,352 | 15.56% | 243,575 | 11.91% | 15,549 | 0.76% | 49,335 | 2.41% | 0 | 0.00% | 0 | 0.00% |
| No mobility limitations | 4,531,815 | 3,915,296 | 86.40% | 434,577 | 9.59% | 132,142 | 2.92% | 27,992 | 0.62% | 21,807 | 0.48% | 0 | 0.00% | 0 | 0.00% |
| N/A - Respondent not asked mobility limitation questions | 6,385,189 | 5,662,971 | 88.69% | 522,498 | 8.18% | 31,684 | 0.50% | 117,676 | 1.84% | 50,360 | 0.79% | 0 | 0.00% | 0 | 0.00% |
| Sleep disorder | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Yes | 1,963,555 | 1,205,154 | 61.38% | 375,568 | 19.13% | 274,288 | 13.97% | 108,545 | 5.53% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| No | 13,504,920 | 11,671,997 | 86.43% | 1,362,356 | 10.09% | 244,533 | 1.81% | 62,448 | 0.46% | 163,587 | 1.21% | 0 | 0.00% | 0 | 0.00% |
| Don't know | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Bowel leakage - gas | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more times a day | 1,691,877 | 1,471,498 | 86.97% | 106,547 | 6.30% | 52,800 | 3.12% | 20,845 | 1.23% | 40,187 | 2.38% | 0 | 0.00% | 0 | 0.00% |
| Once a day | 1,484,183 | 1,230,769 | 82.93% | 175,239 | 11.81% | 68,400 | 4.61% | 9,775 | 0.66% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| 2 or more time a week | 1,294,731 | 1,045,769 | 80.77% | 217,531 | 16.80% | 31,431 | 2.43% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Once a week | 1,010,064 | 660,250 | 65.37% | 153,510 | 15.20% | 50,902 | 5.04% | 117,676 | 11.65% | 27,726 | 2.74% | 0 | 0.00% | 0 | 0.00% |
| 1-3 times a month | 2,725,838 | 2,357,668 | 86.49% | 276,177 | 10.13% | 91,993 | 3.37% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% | 0 | 0.00% |
| Never | 7,013,931 | 5,918,677 | 84.38% | 808,921 | 11.53% | 167,962 | 2.39% | 22,696 | 0.32% | 95,674 | 1.36% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KI.Q.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?

Table M07 - Extent to which daily activities were affected among those men who responded 'yes' to experiencing any urinary incontinence, 2009 - 2010

| | Total | Not at all (N) | Not at all (%) | Only a little (N) | Only a little (%) | Somewhat (N) | Somewhat (%) | Very much (N) | Very much (%) | Greatly (N) | Greatly (%) | Refused to Answer or Don't Know (N) | Refused to Answer or Don't Know (%) | Missing (N) | Missing (%) |
|------------------------------|------------|----------------|----------------|-------------------|-------------------|--------------|--------------|---------------|---------------|-------------|-------------|-------------------------------------|-------------------------------------|-------------|-------------|
| Total | 15,475,302 | 12,883,977 | 83.26% | 1,737,924 | 11.23% | 518,821 | 3.35% | 170,992 | 1.10% | 163,587 | 1.06% | 0 | 0.00% | 0 | 0.00% |
| Bowel leakage - stool | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| Missing | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a day | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 2 or more time a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| Once a week | * | * | * | * | * | * | * | * | * | * | * | * | * | * | * |
| 1-3 times a month | 466,082 | 240,671 | 51.64% | 185,224 | 39.74% | 0 | 0.00% | 0 | 0.00% | 40,187 | 8.62% | 0 | 0.00% | 0 | 0.00% |
| Never | 14,575,578 | 12,307,104 | 84.44% | 1,510,592 | 10.36% | 463,489 | 3.18% | 170,992 | 1.17% | 123,400 | 0.85% | 0 | 0.00% | 0 | 0.00% |

Source: National Health and Nutrition Examination Survey 2001-2010

NOTE: CI stands for Confidence Interval.

NOTE: * Represents that the figure does not meet the standard for reliability of precision

NOTE: The data in this table are based on question KIQ.052: During the past 12 months, how much did your/his/her leakage of urine affect your/his/her day-to-day activities?