

Table M06 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Urinary Incontinence, 2002 - 2004

|               | 2002       |                           | 2003       |                           | 2004       |                           |
|---------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|
|               | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients |
| <b>Total</b>  | 1,128      | 40,474                    | 1,562      | 36,292                    | 2,315      | 39,337                    |
| <b>Age</b>    | -          | -                         | -          | -                         | -          | -                         |
| 18-24         | 32         | 30,189 *                  | 48         | 30,968 *                  | 78         | 43,094 *                  |
| 25-34         | 53         | 27,749 *                  | 111        | 34,365                    | 150        | 37,406 *                  |
| 35-39         | 62         | 44,604 <sup>R</sup>       | 98         | 39,200 <sup>R</sup>       | 92         | 28,049 <sup>R</sup>       |
| 40-44         | 86         | 40,566 *                  | 111        | 34,796 *                  | 140        | 30,973 *                  |
| 45-54         | 236        | 31,763 *                  | 398        | 31,339 *                  | 603        | 34,715 *                  |
| 55-64         | 659        | 47,206                    | 796        | 40,060 *                  | 1,252      | 44,939 *                  |
| <b>Region</b> | -          | -                         | -          | -                         | -          | -                         |
| Northeast     | 131        | 48,519                    | 187        | 39,203 *                  | 224        | 36,305 *                  |
| South         | 400        | 36,364 *                  | 605        | 35,012 *                  | 1,089      | 40,037 *                  |
| Midwest       | 472        | 46,365 <sup>R</sup>       | 527        | 43,880 <sup>R</sup>       | 671        | 44,823 <sup>R</sup>       |
| West          | 125        | 31,328 *                  | 243        | 27,060 *                  | 331        | 31,494 *                  |

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Urinary Incontinence, 2005 - 2007

|               | 2005       |                           | 2006       |                           | 2007       |                           |
|---------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|
|               | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients |
| <b>Total</b>  | 2,569      | 36,528                    | 3,081      | 31,720                    | 3,792      | 33,327                    |
| <b>Age</b>    | -          | -                         | -          | -                         | -          | -                         |
| 18-24         | 85         | 39,906 *                  | 111        | 35,691                    | 118        | 35,119 *                  |
| 25-34         | 138        | 28,337                    | 191        | 26,977 *                  | 194        | 24,129                    |
| 35-39         | 103        | 27,989 <sup>R</sup>       | 195        | 36,585 <sup>R</sup>       | 138        | 22,623 <sup>R</sup>       |
| 40-44         | 141        | 25,314 *                  | 161        | 22,025 *                  | 239        | 25,345 *                  |
| 45-54         | 737        | 34,026 *                  | 851        | 27,558 *                  | 1,250      | 32,869 *                  |
| 55-64         | 1,365      | 42,104 *                  | 1,572      | 36,205 *                  | 1,853      | 37,956 *                  |
| <b>Region</b> | -          | -                         | -          | -                         | -          | -                         |
| Northeast     | 314        | 37,028 *                  | 380        | 33,304                    | 486        | 38,787 *                  |
| South         | 1,185      | 36,904 *                  | 1,533      | 31,772                    | 1,936      | 33,722                    |
| Midwest       | 696        | 40,442 <sup>R</sup>       | 848        | 33,177 <sup>R</sup>       | 987        | 33,435 <sup>R</sup>       |
| West          | 374        | 29,848 *                  | 320        | 26,868 *                  | 383        | 26,746 *                  |

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Urinary Incontinence, 2008 - 2010

|               | 2008       |                           | 2009       |                           | 2010       |                           |
|---------------|------------|---------------------------|------------|---------------------------|------------|---------------------------|
|               | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients |
| <b>Total</b>  | 5,774      | 35,341                    | 8,029      | 37,921                    | 6,736      | 35,060                    |
| <b>Age</b>    | -          | -                         | -          | -                         | -          | -                         |
| 18-24         | 188        | 37,600 *                  | 243        | 41,538 *                  | 198        | 40,326 *                  |
| 25-34         | 361        | 32,203 *                  | 455        | 30,599                    | 373        | 27,266                    |
| 35-39         | 249        | 26,717 <sup>R</sup>       | 330        | 28,278 <sup>R</sup>       | 264        | 25,832 <sup>R</sup>       |
| 40-44         | 317        | 24,980 *                  | 457        | 27,849 *                  | 402        | 25,869 *                  |
| 45-54         | 1,702      | 32,500 *                  | 2,546      | 34,853 *                  | 2,390      | 34,683 *                  |
| 55-64         | 2,957      | 40,624 *                  | 3,998      | 44,482 *                  | 3,109      | 39,419 *                  |
| <b>Region</b> | -          | -                         | -          | -                         | -          | -                         |
| Northeast     | 1,103      | 31,632 *                  | 1,441      | 39,807 *                  | 1,356      | 33,399 *                  |
| South         | 2,309      | 34,806 *                  | 3,113      | 34,497 *                  | 2,514      | 33,524 *                  |
| Midwest       | 1,588      | 39,258 <sup>R</sup>       | 2,341      | 44,633 <sup>R</sup>       | 1,844      | 43,884 <sup>R</sup>       |
| West          | 774        | 35,635 *                  | 1,134      | 34,531 *                  | 1,022      | 29,606 *                  |

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Urinary Incontinence, 2011

|               | 2011       |                           |
|---------------|------------|---------------------------|
|               | Count of X | Rate per 100,000 Patients |
| <b>Total</b>  | 8,343      | 36,239                    |
| <b>Age</b>    | -          | -                         |
| 18-24         | 280        | 42,169 *                  |
| 25-34         | 480        | 30,829                    |
| 35-39         | 392        | 31,335 <sup>R</sup>       |
| 40-44         | 531        | 28,969 *                  |
| 45-54         | 2,807      | 35,428 *                  |
| 55-64         | 3,853      | 39,340 *                  |
| <b>Region</b> | -          | -                         |
| Northeast     | 1,953      | 32,458 *                  |
| South         | 2,987      | 35,585 *                  |
| Midwest       | 2,124      | 43,453 <sup>R</sup>       |
| West          | 1,279      | 34,354 *                  |

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of urinary incontinence.