

Table M07 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002		2003		2004	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	945	33,907	1,287	29,902	1,930	32,795
Age	-	-	-	-	-	-
18-24	28	26,415	35	22,581	67	37,017 *
25-34	45	23,560	89	27,554	121	30,175 *
35-39	37	26,619 ^R	69	27,600 ^R	74	22,561 ^R
40-44	72	33,962 *	79	24,765 *	111	24,558 *
45-54	183	24,630 *	317	24,961 *	494	28,440 *
55-64	580	41,547 *	698	35,128 *	1,063	38,155 *
Region	-	-	-	-	-	-
Northeast	100	37,037	144	30,189 *	195	31,605 *
South	344	31,273 *	495	28,646 *	891	32,757 *
Midwest	392	38,507 ^R	441	36,719 ^R	555	37,074 ^R
West	109	27,318 *	207	23,051 *	289	27,498 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005		2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	2,170	30,855	2,574	26,501	3,213	28,239
Age	-	-	-	-	-	-
18-24	74	34,742 *	101	32,476	100	29,762 *
25-34	104	21,355	153	21,610 *	164	20,398
35-39	90	24,457 ^R	147	27,580 ^R	105	17,213 ^R
40-44	117	21,005 *	131	17,921 *	192	20,361 *
45-54	606	27,978 *	707	22,895 *	1,052	27,662 *
55-64	1,179	36,366 *	1,335	30,746 *	1,600	32,773 *
Region	-	-	-	-	-	-
Northeast	261	30,778 *	318	27,870	420	33,520 *
South	996	31,018 *	1,252	25,948 *	1,587	27,643
Midwest	609	35,386 ^R	731	28,599 ^R	864	29,268 ^R
West	304	24,262 *	273	22,922 *	342	23,883 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008		2009		2010	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	4,862	29,759	6,715	31,715	5,739	29,870
Age	-	-	-	-	-	-
18-24	169	33,800 *	216	36,923 *	175	35,642 *
25-34	299	26,673 *	373	25,084 *	294	21,491
35-39	212	22,747 ^R	255	21,851 ^R	236	23,092 ^R
40-44	245	19,307 *	352	21,450 *	334	21,493 *
45-54	1,404	26,809 *	2,124	29,076 *	2,047	29,705 *
55-64	2,533	34,799 *	3,395	37,773 *	2,653	33,638 *
Region	-	-	-	-	-	-
Northeast	913	26,183 *	1,211	33,453 *	1,128	27,783 *
South	1,911	28,806 *	2,540	28,147 *	2,141	28,550 *
Midwest	1,345	33,251 ^R	1,967	37,502 ^R	1,589	37,815 ^R
West	693	31,906 *	997	30,359 *	881	25,521 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Stress Urinary Incontinence, 2011

	2011	
	Count of X	Rate per 100,000 Patients
Total	7,144	31,031
Age	-	-
18-24	247	37,199 *
25-34	395	25,369
35-39	341	27,258 ^R
40-44	413	22,531 *
45-54	2,396	30,241 *
55-64	3,352	34,225 *
Region	-	-
Northeast	1,642	27,289 *
South	2,586	30,808 *
Midwest	1,828	37,398 ^R
West	1,088	29,224 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.