Table M09 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Other Urinary Incontinence, 2002 - 2004

| | 200 | 2002 | | 2003 | | 2004 | |
|-----------|------------|---------------------------------|------------|---------------------------------|------------|---------------------------------|--|
| | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | |
| Total | 36 | 1,292 | 54 | 1,255 | 83 | 1,410 | |
| Age | - | - | - | _ | - | - | |
| 18-24 | 1 | 943 * | 2 | 1,290 * | 3 | 1,658 | |
| 25-34 | 0 | 0 * | 4 | 1,238 * | 6 | 1,496 | |
| 35-39 | 16 | 11,511 ^R | 17 | 6,800 R | 3 | 915 ^R | |
| 40-44 | 0 | 0 * | 0 | 0 * | 9 | 1,991 * | |
| 45-54 | 6 | 808 * | 13 | 1,024 * | 19 | 1,094 * | |
| 55-64 | 13 | 931 * | 18 | 906 * | 43 | 1,543 * | |
| Region | - | - | - | - | - | - | |
| Northeast | 16 | 5,926 * | 18 | 3,774 * | 9 | 1,459 | |
| South | 10 | 909 | 16 | 926 | 39 | 1,434 | |
| Midwest | 7 | 688 ^R | 11 | 916 R | 29 | 1,937 R | |
| West | 3 | 752 * | 9 | 1,002 * | 6 | 571 * | |

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of other urinary incontinence.

 $^{^{\}mbox{\tiny R}}$ Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

Table M09 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Other Urinary Incontinence, 2005 - 2007

| | 2005 | | 2006 | | 2007 | |
|-----------|------------|---------------------------------|------------|---------------------------------|------------|---------------------------------|
| | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients |
| Total | 80 | 1,138 | 104 | 1,071 | 91 | 800 |
| Age | - | - | - | - | - | - |
| 18-24 | 2 | 939 * | 2 | 643 | 1 | 298 |
| 25-34 | 7 | 1,437 * | 6 | 848 | 7 | 871 |
| 35-39 | 0 | 0 R | 7 | 1,313 R | 5 | 820 R |
| 40-44 | 5 | 898 * | 5 | 684 * | 3 | 318 * |
| 45-54 | 14 | 646 * | 34 | 1,101 * | 34 | 894 * |
| 55-64 | 52 | 1,604 * | 50 | 1,152 * | 41 | 840 * |
| Region | - | - | - | - | - | - |
| Northeast | 11 | 1,297 | 16 | 1,402 | 16 | 1,277 * |
| South | 26 | 810 | 56 | 1,161 | 49 | 854 * |
| Midwest | 17 | 988 ^R | 24 | 939 R | 15 | 508 R |
| West | 26 | 2,075 * | 8 | 672 * | 11 | 768 * |

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of other urinary incontinence.

 $^{^{\}mbox{\tiny R}}$ Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

Table M09 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Other Urinary Incontinence, 2008 - 2010

| | 200 | 2008 | | 2009 | | 2010 | |
|-----------|------------|---------------------------------|------------|---------------------------------|------------|---------------------------------|--|
| | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | Count of X | Rate per 100,000 Patients | |
| Total | 214 | 1,310 | 278 | 1,313 | 194 | 1,010 | |
| Age | - | - | - | - | - | - | |
| 18-24 | 5 | 1,000 | 10 | 1,709 * | 15 | 3,055 * | |
| 25-34 | 29 | 2,587 * | 14 | 942 | 19 | 1,389 * | |
| 35-39 | 6 | 644 ^R | 6 | 514 ^R | 3 | 294 R | |
| 40-44 | 12 | 946 * | 22 | 1,341 * | 12 | 772 * | |
| 45-54 | 59 | 1,127 * | 85 | 1,164 * | 63 | 914 * | |
| 55-64 | 103 | 1,415 * | 141 | 1,569 * | 82 | 1,040 * | |
| Region | - | - | - | - | - | - | |
| Northeast | 59 | 1,692 * | 57 | 1,575 | 56 | 1,379 * | |
| South | 90 | 1,357 | 105 | 1,164 * | 74 | 987 | |
| Midwest | 50 | 1,236 R | 82 | 1,563 R | 35 | 833 R | |
| West | 15 | 691 * | 34 | 1,035 * | 29 | 840 * | |

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of other urinary incontinence.

R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

Table M09 - Ambulatory Visits (Other than Physician Visits) by Privately Insured Males with Any Diagnosis of Other Urinary Incontinence, 2011

| | 201 | 2011 | | |
|-----------|------------|---------------------------------|--|--|
| | Count of X | Rate per 100,000 Patients | | |
| Total | 274 | 1,190 | | |
| Age | - | - | | |
| 18-24 | 10 | 1,506 | | |
| 25-34 | 21 | 1,349 | | |
| 35-39 | 12 | 959 R | | |
| 40-44 | 29 | 1,582 * | | |
| 45-54 | 82 | 1,035 * | | |
| 55-64 | 120 | 1,225 * | | |
| Region | - | - | | |
| Northeast | 83 | 1,379 | | |
| South | 90 | 1,072 | | |
| Midwest | 52 | 1,064 R | | |
| West | 49 | 1,316 * | | |

NOTE: X represents the number of ambulatory visits (other than physician visits) by privately insured male adults with any diagnosis of other urinary incontinence.

R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group