

Table M10 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Urinary Incontinence,
2002 - 2004

	2002		2003		2004	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	23	825	44	1,022	42	714
Age	-	-	-	-	-	-
18-24	1	943	0	0	1	553
25-34	1	524	1	310	3	748
35-39	0	0 ^R	1	400 ^R	2	610 ^R
40-44	1	472 [*]	1	314 [*]	2	443 [*]
45-54	6	808 [*]	16	1,260 [*]	12	691 [*]
55-64	14	1,003 [*]	25	1,258 [*]	22	790 [*]
Region	-	-	-	-	-	-
Northeast	2	741	11	2,306 [*]	3	486
South	9	818	15	868	15	552
Midwest	8	786 ^R	5	416 ^R	9	601 ^R
West	4	1,003 [*]	13	1,448 [*]	15	1,427 [*]

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Urinary Incontinence,
2005 - 2007

	2005		2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	43	611	50	515	45	396
Age	-	-	-	-	-	-
18-24	1	470	0	0	2	595 *
25-34	1	205	0	0	2	249
35-39	3	815 ^R	2	375 ^R	0	0 ^R
40-44	0	0 *	0	0 *	2	212 *
45-54	8	369 *	10	324 *	13	342 *
55-64	30	925 *	38	875 *	26	533 *
Region	-	-	-	-	-	-
Northeast	4	472	7	614	3	239
South	18	561	29	601	20	348
Midwest	13	755 ^R	11	430 ^R	16	542 ^R
West	8	639 *	3	252 *	6	419 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Urinary Incontinence,
2008 - 2010

	2008		2009		2010	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	87	533	91	430	63	328
Age	-	-	-	-	-	-
18-24	2	400 *	0	0 *	1	204
25-34	2	178	5	336 *	0	0
35-39	0	0 ^R	0	0 ^R	1	98 ^R
40-44	3	236 *	5	305 *	1	64 *
45-54	23	439 *	21	288 *	22	319 *
55-64	57	783 *	60	668 *	38	482 *
Region	-	-	-	-	-	-
Northeast	16	459	21	580	10	246 *
South	37	558	34	377	22	293 *
Midwest	16	396 ^R	20	381 ^R	21	500 ^R
West	18	829 *	16	487 *	10	290 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Urinary Incontinence, 2011

	2011	
	Count of X	Rate per 100,000 Patients
Total	64	278
Age	-	-
18-24	1	151
25-34	0	0 *
35-39	0	0 ^R
40-44	4	218 *
45-54	29	366 *
55-64	30	306 *
Region	-	-
Northeast	12	199 *
South	18	214 *
Midwest	20	409 ^R
West	14	376 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of urinary incontinence.