

Table M11 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Stress Urinary Incontinence,
2002 - 2004

	2002		2003		2004	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	18	646	29	674	30	510
Age	-	-	-	-	-	-
18-24	1	943	0	0 *	0	0
25-34	1	524	1	310	2	499
35-39	0	0 ^R	0	0 ^R	1	305 ^R
40-44	0	0 *	0	0 *	2	443 *
45-54	5	673 *	10	787 *	8	461 *
55-64	11	788 *	18	906 *	17	610 *
Region	-	-	-	-	-	-
Northeast	2	741	7	1,468 *	2	324
South	7	636	10	579	10	368
Midwest	6	589 ^R	4	333 ^R	6	401 ^R
West	3	752 *	8	891 *	12	1,142 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Stress Urinary Incontinence,
2005 - 2007

	2005		2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	33	469	37	381	35	308
Age	-	-	-	-	-	-
18-24	1	470	0	0	1	298
25-34	1	205	0	0	0	0 *
35-39	1	272 ^R	1	188 ^R	0	0 ^R
40-44	0	0 *	0	0 *	1	106 *
45-54	7	323 *	7	227 *	10	263 *
55-64	23	709 *	29	668 *	23	471 *
Region	-	-	-	-	-	-
Northeast	4	472	5	438	3	239
South	16	498	23	477	15	261
Midwest	9	523 ^R	7	274 ^R	11	373 ^R
West	4	319 *	2	168 *	6	419 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Stress Urinary Incontinence,
2008 - 2010

	2008		2009		2010	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	67	410	74	350	45	234
Age	-	-	-	-	-	-
18-24	0	0 *	0	0 *	0	0 *
25-34	0	0 *	3	202	0	0 *
35-39	0	0 ^R	0	0 ^R	0	0 ^R
40-44	1	79 *	5	305 *	0	0 *
45-54	20	382 *	16	219 *	13	189 *
55-64	46	632 *	50	556 *	32	406 *
Region	-	-	-	-	-	-
Northeast	11	316	17	470	8	197
South	27	407	28	310	18	240
Midwest	16	396 ^R	17	324 ^R	12	286 ^R
West	13	599 *	12	365 *	7	203 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Stress Urinary Incontinence, 2011

	2011	
	Count of X	Rate per 100,000 Patients
Total	47	204
Age	-	-
18-24	0	0 *
25-34	0	0 *
35-39	0	0 ^R
40-44	1	55 *
45-54	23	290 *
55-64	23	235 *
Region	-	-
Northeast	8	133 *
South	11	131 *
Midwest	17	348 ^R
West	11	296 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of stress urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of stress urinary incontinence.