

Table M13 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Other Urinary Incontinence,  
2002 - 2004

	2002		2003		2004	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
<b>Total</b>	3	108	5	116	8	136
<b>Age</b>	-	-	-	-	-	-
18-24	0	0 *	0	0 *	0	0 *
25-34	0	0 *	0	0 *	1	249
35-39	0	0 <sup>R</sup>	0	0 <sup>R</sup>	0	0 <sup>R</sup>
40-44	0	0 *	0	0 *	0	0 *
45-54	0	0 *	3	236 *	3	173 *
55-64	3	215 *	2	101 *	4	144 *
<b>Region</b>	-	-	-	-	-	-
Northeast	0	0	2	419 *	0	0
South	0	0	1	58	4	147
Midwest	2	197 <sup>R</sup>	0	0 <sup>R</sup>	1	67 <sup>R</sup>
West	1	251 *	2	223 *	3	285 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of other urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Other Urinary Incontinence,  
2005 - 2007

	2005		2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
<b>Total</b>	6	85	6	62	3	26
<b>Age</b>	-	-	-	-	-	-
18-24	0	0 *	0	0	0	0 *
25-34	0	0 *	0	0	0	0 *
35-39	0	0 <sup>R</sup>	1	188 <sup>R</sup>	0	0 <sup>R</sup>
40-44	0	0 *	0	0 *	0	0 *
45-54	1	46 *	2	65 *	2	53 *
55-64	5	154 *	3	69 *	1	21 *
<b>Region</b>	-	-	-	-	-	-
Northeast	0	0	2	175	0	0 *
South	2	62	3	62	3	52
Midwest	1	58 <sup>R</sup>	1	39 <sup>R</sup>	0	0 <sup>R</sup>
West	3	239 *	0	0 *	0	0 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of other urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Other Urinary Incontinence, 2008 - 2010

	2008		2009		2010	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
<b>Total</b>	10	61	7	33	5	26
<b>Age</b>	-	-	-	-	-	-
18-24	0	0 *	0	0 *	0	0 *
25-34	0	0 *	1	67	0	0 *
35-39	0	0 <sup>R</sup>	0	0 <sup>R</sup>	0	0 <sup>R</sup>
40-44	1	79 *	0	0 *	1	64 *
45-54	1	19 *	2	27 *	1	15 *
55-64	8	110 *	4	45 *	3	38 *
<b>Region</b>	-	-	-	-	-	-
Northeast	3	86 *	2	55	2	49
South	5	75 *	2	22	1	13
Midwest	0	0 <sup>R</sup>	1	19 <sup>R</sup>	0	0 <sup>R</sup>
West	2	92 *	2	61 *	2	58 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of other urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Privately Insured Males with a Primary Diagnosis of Other Urinary Incontinence, 2011

	2011	
	Count of X	Rate per 100,000 Patients
<b>Total</b>	3	13
<b>Age</b>	-	-
18-24	0	0 *
25-34	0	0 *
35-39	0	0 <sup>R</sup>
40-44	0	0 *
45-54	1	13 *
55-64	2	20 *
<b>Region</b>	-	-
Northeast	2	33
South	1	12
Midwest	0	0 <sup>R</sup>
West	0	0 *

Source: Truven Health MarketScan Research Database, 2002 - 2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by privately insured male adults with a primary diagnosis of other urinary incontinence.

NOTE: Rate is per 100,000 privately insured male adults who were hospitalized for a primary diagnosis of other urinary incontinence.