Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2002 - 2003

	20	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	
Total	27	968.8	30	697.0	
All Procedures	27	968.8	30	697.0	
Age	-	-	-	-	
18-24	2	1,886.8	0	0.0	
25-34	0	0.0	2	619.2	
35-39	2	1,438.8	0	0.0	
40-44	0	0.0	0	0.0	
45-54	5	672.9	8	629.9	
55-64	18	1,289.4	20	1,006.5	
Region	-	-	-	-	
Northeast	3	1,111.1	6	1,257.9	
South	12	1,090.9	9	520.8	
Midwest	6	589.4	10	832.6	
West	6	1,503.8	5	556.8	

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2002 - 2003

	200	2002		03
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	27	968.8	30	697.0
Operation for correction of incontinence	21	753.5	25	580.9
Age	_	-	-	-
18-24	1	943.4	0	0.0
25-34	0	0.0	1	309.6
35-39	1	719.4	0	0.0
40-44	0	0.0	0	0.0
45-54	5	672.9	7	551.2
55-64	14	1,002.9	17	855.6
Region	_	-	-	-
Northeast	2	740.7	6	1,257.9
South	9	818.2	7	405.1
Midwest	6	589.4	8	666.1
West	4	1,002.5	4	445.4
Revision or repair of prosthetic	6	215.3	5	116.2
Age	-	-	-	-
18-24	1	943.4	0	0.0
25-34	0	0.0	1	309.6
35-39	1	719.4	0	0.0
40-44	0	0.0	0	0.0
45-54	0	0.0	1	78.7
55-64	4	286.5	3	151.0
Region	-	-	-	-
Northeast	1	370.4	0	0.0
South	3	272.7	2	115.7
Midwest	0	0.0	2	166.5
West	2	501.3	1	111.4

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2004 - 2005

	20	2004		05
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	34	577.7	50	710.9
All Procedures	34	577.7	50	710.9
Age	-	-	-	-
18-24	0	0.0	0	0.0
25-34	1	249.4	2	410.7
35-39	3	914.6	3	815.2
40-44	0	0.0	0	0.0
45-54	6	345.4	5	230.8
55-64	24	861.5	40	1,233.8
Region	-	-	-	-
Northeast	1	162.1	3	353.8
South	6	220.6	16	498.3
Midwest	13	868.4	22	1,278.3
West	14	1,332.1	9	718.3

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2004 - 2005

	20	2004		05
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	34	577.7	50	710.9
Operation for correction of incontinence	29	492.8	41	583.0
Age	-	-	-	-
18-24	0	0.0	0	0.0
25-34	1	249.4	1	205.3
35-39	2	609.8	2	543.5
40-44	0	0.0	0	0.0
45-54	5	287.9	5	230.8
55-64	21	753.8	33	1,017.9
Region	-	_	-	_
Northeast	0	0.0	2	235.8
South	5	183.8	12	373.7
Midwest	12	801.6	18	1,045.9
West	12	1,141.8	9	718.3
Revision or repair of prosthetic	5	85.0	9	128.0
Age	-	-	-	<u>-</u>
18-24	0	0.0	0	0.0
25-34	0	0.0	1	205.3
35-39	1	304.9	1	271.7
40-44	0	0.0	0	0.0
45-54	1	57.6	0	0.0
55-64	3	107.7	7	215.9
Region	-	-	-	
Northeast	1	162.1	1	117.9
South	1	36.8	4	124.6
Midwest	1	66.8	4	232.4
West	2	190.3	0	0.0

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2006 - 2007

	20	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients	
Total	56	576.5	48	421.9	
All Procedures	56	576.5	48	421.9	
Age	-	-	-	-	
18-24	2	643.1	2	595.2	
25-34	0	0.0	0	0.0	
35-39	1	187.6	0	0.0	
40-44	1	136.8	1	106.0	
45-54	7	226.7	8	210.4	
55-64	45	1,036.4	37	757.9	
Region	-	_	-	-	
Northeast	8	701.1	10	798.1	
South	10	207.3	9	156.8	
Midwest	35	1,369.3	25	846.9	
West	3	251.9	4	279.3	

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2006 - 2007

	200	2006		07
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	56	576.5	48	421.9
Operation for correction of incontinence	45	463.3	38	334.0
Age	_	-	-	-
18-24	2	643.1	1	297.6
25-34	0	0.0	0	0.0
35-39	1	187.6	0	0.0
40-44	1	136.8	1	106.0
45-54	6	194.3	8	210.4
55-64	35	806.1	28	573.5
Region	_	-	-	_
Northeast	8	701.1	7	558.7
South	9	186.5	8	139.3
Midwest	27	1,056.3	20	677.5
West	1	84.0	3	209.5
Revision or repair of prosthetic	11	113.3	10	87.9
Age	-	-	-	
18-24	0	0.0	1	297.6
25-34	0	0.0	0	0.0
35-39	0	0.0	0	0.0
40-44	0	0.0	0	0.0
45-54	1	32.4	0	0.0
55-64	10	230.3	9	184.4
Region	-	-	-	_
Northeast	0	0.0	3	239.4
South	1	20.7	1	17.4
Midwest	8	313.0	5	169.4
West	2	167.9	1	69.8

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2008 - 2009

	20	2008		09
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	85	520.3	77	363.7
All Procedures	85	520.3	77	363.7
Age	-	-	-	-
18-24	1	200.0	0	0.0
25-34	0	0.0	2	134.5
35-39	2	214.6	2	171.4
40-44	0	0.0	3	182.8
45-54	12	229.1	11	150.6
55-64	70	961.7	59	656.4
Region	-	-	-	-
Northeast	14	401.5	11	303.9
South	20	301.5	23	254.9
Midwest	31	766.4	25	476.6
West	20	920.8	18	548.1

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2008 - 2009

	200	2008		09
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	85	520.3	77	363.7
Operation for correction of incontinence	73	446.8	64	302.3
Age	_	-	_	-
18-24	1	200.0	0	0.0
25-34	0	0.0	1	67.2
35-39	1	107.3	1	85.7
40-44	0	0.0	1	60.9
45-54	12	229.1	9	123.2
55-64	59	810.6	52	578.5
Region	_	_	_	_
Northeast	11	315.5	8	221.0
South	17	256.3	19	210.5
Midwest	28	692.2	23	438.5
West	17	782.7	14	426.3
Revision or repair of prosthetic	12	73.4	13	61.4
Age	-	-	_	_
18-24	0	0.0	0	0.0
25-34	0	0.0	1	67.2
35-39	1	107.3	1	85.7
40-44	0	0.0	2	121.9
45-54	0	0.0	2	27.4
55-64	11	151.1	7	77.9
Region	-	-	-	-
Northeast	3	86.0	3	82.9
South	3	45.2	4	44.3
Midwest	3	74.2	2	38.1
West	3	138.1	4	121.8

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2010 - 2011

	20	2010		11
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	45	234.2	45	195.5
All Procedures	45	234.2	45	195.5
Age	-	-	-	-
18-24	1	203.7	0	0.0
25-34	0	0.0	1	64.2
35-39	0	0.0	1	79.9
40-44	0	0.0	0	0.0
45-54	10	145.1	7	88.4
55-64	34	431.1	36	367.6
Region	-	-	-	-
Northeast	8	197.0	10	166.2
South	10	133.4	15	178.7
Midwest	17	404.6	10	204.6
West	10	289.7	10	268.6

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.

Table M16b - Surgical Procedures for Privately Insured Males with a Primary Diagnosis of Urinary Incontinence by Age and Geographic Region, 2010 - 2011

	20	2010		11
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	45	234.2	45	195.5
Operation for correction of incontinence	36	187.4	42	182.4
Age	_	-	-	-
18-24	0	0.0	0	0.0
25-34	0	0.0	0	0.0
35-39	0	0.0	1	79.9
40-44	0	0.0	0	0.0
45-54	8	116.1	7	88.4
55-64	28	355.0	34	347.2
Region	_	-	-	_
Northeast	7	172.4	9	149.6
South	8	106.7	14	166.8
Midwest	13	309.4	9	184.1
West	8	231.7	10	268.6
Revision or repair of prosthetic	9	46.8	3	13.0
Age	-	-	-	<u>-</u>
18-24	1	203.7	0	0.0
25-34	0	0.0	1	64.2
35-39	0	0.0	0	0.0
40-44	0	0.0	0	0.0
45-54	2	29.0	0	0.0
55-64	6	76.1	2	20.4
Region	-	-	-	-
Northeast	1	24.6	1	16.6
South	2	26.7	1	11.9
Midwest	4	95.2	1	20.5
West	2	57.9	0	0.0

NOTE: X represents the number of surgical procedures for privately insured male adults with a primary

diagnosis of urinary incontinence.