

Table F03 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
<b>Total</b>	766,040	92,492	764,880	91,344	758,800	90,348
<b>Age</b>	-	-	-	-	-	-
65-69	153,780	100,300 <sup>R</sup>	152,720	97,822 <sup>R</sup>	154,860	95,912 <sup>R</sup>
70-74	175,000	97,776 <sup>*</sup>	165,340	95,783 <sup>*</sup>	161,040	95,346 <sup>*</sup>
75-79	180,940	95,786 <sup>*</sup>	181,040	94,538 <sup>*</sup>	180,000	95,379 <sup>*</sup>
80-84	142,620	91,236 <sup>*</sup>	146,400	90,516 <sup>*</sup>	144,500	88,945 <sup>*</sup>
85+	113,700	75,448 <sup>*</sup>	119,380	76,831 <sup>*</sup>	118,400	74,785 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	696,660	93,334 <sup>R</sup>	693,980	92,479 <sup>R</sup>	678,940	91,459 <sup>R</sup>
Black	33,920	74,879 <sup>*</sup>	33,660	72,481 <sup>*</sup>	35,300	71,026 <sup>*</sup>
Hispanic	9,540	93,346 <sup>*</sup>	12,000	105,079 <sup>*</sup>	13,660	98,699 <sup>*</sup>
Asian	6,340	99,686 <sup>*</sup>	6,620	89,218 <sup>*</sup>	9,500	91,699 <sup>*</sup>
Other	16,020	98,767 <sup>*</sup>	15,920	86,147 <sup>*</sup>	18,460	91,386 <sup>*</sup>
Unknown	3,560	96,216 <sup>*</sup>	2,700	84,906 <sup>*</sup>	2,940	85,965 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	131,100	89,378	130,460	88,149 <sup>*</sup>	132,740	88,024 <sup>*</sup>
South	304,460	93,998 <sup>*</sup>	313,240	95,002 <sup>*</sup>	303,480	93,580 <sup>*</sup>
Midwest	188,120	87,441 <sup>R</sup>	177,720	84,839 <sup>R</sup>	178,820	83,600 <sup>R</sup>
West	142,360	99,902 <sup>*</sup>	143,460	95,538 <sup>*</sup>	143,760	95,294 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table F03 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
<b>Total</b>	770,820	89,127	763,600	88,269	747,660	89,190
<b>Age</b>	-	-	-	-	-	-
65-69	157,080	95,350 <sup>R</sup>	158,040	93,063 <sup>R</sup>	153,300	95,017 <sup>R</sup>
70-74	158,640	91,837 <sup>*</sup>	158,860	93,723 <sup>*</sup>	156,500	94,551 <sup>*</sup>
75-79	177,600	92,994 <sup>*</sup>	171,100	91,654 <sup>*</sup>	161,240	92,486 <sup>*</sup>
80-84	151,920	88,676 <sup>*</sup>	148,940	88,507 <sup>*</sup>	146,540	89,660 <sup>*</sup>
85+	125,580	76,072 <sup>*</sup>	126,660	74,157 <sup>*</sup>	130,080	74,914 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	694,900	90,595 <sup>R</sup>	690,300	89,350 <sup>R</sup>	678,740	90,090 <sup>R</sup>
Black	32,720	65,941 <sup>*</sup>	32,580	68,188 <sup>*</sup>	30,380	70,849 <sup>*</sup>
Hispanic	14,980	95,051 <sup>*</sup>	14,440	93,161 <sup>*</sup>	14,380	99,723 <sup>*</sup>
Asian	9,480	94,800 <sup>*</sup>	8,920	94,692 <sup>*</sup>	9,220	95,445 <sup>*</sup>
Other	16,240	85,926 <sup>*</sup>	14,740	86,604 <sup>*</sup>	12,620	82,376 <sup>*</sup>
Unknown	2,500	70,621 <sup>*</sup>	2,620	94,245 <sup>*</sup>	2,320	89,231 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	132,120	87,242 <sup>*</sup>	132,860	86,284 <sup>*</sup>	132,000	86,842 <sup>*</sup>
South	310,220	91,939 <sup>*</sup>	314,320	92,507 <sup>*</sup>	305,620	92,635 <sup>*</sup>
Midwest	177,260	81,830 <sup>R</sup>	176,760	80,757 <sup>R</sup>	170,580	81,719 <sup>R</sup>
West	151,220	94,880 <sup>*</sup>	139,660	91,616 <sup>*</sup>	139,460	94,472 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table F03 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
<b>Total</b>	755,120	88,902	792,240	89,236	824,120	90,545
<b>Age</b>	-	-	-	-	-	-
65-69	157,120	93,680 <sup>R</sup>	166,440	94,023 <sup>R</sup>	172,500	95,251 <sup>R</sup>
70-74	156,820	91,901 <sup>*</sup>	163,640	92,661 <sup>*</sup>	173,980	95,762 <sup>*</sup>
75-79	155,360	91,777 <sup>*</sup>	167,400	93,269 <sup>*</sup>	171,520	94,314 <sup>*</sup>
80-84	150,760	91,603 <sup>*</sup>	150,180	88,927 <sup>*</sup>	149,240	88,235 <sup>*</sup>
85+	135,060	76,236 <sup>*</sup>	144,580	77,806 <sup>*</sup>	156,880	79,878 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	680,840	89,398 <sup>R</sup>	714,560	90,034 <sup>R</sup>	741,520	91,318 <sup>R</sup>
Black	31,680	71,159 <sup>*</sup>	34,020	69,231 <sup>*</sup>	35,060	69,757 <sup>*</sup>
Hispanic	14,740	105,436 <sup>*</sup>	14,900	96,253 <sup>*</sup>	16,440	101,232 <sup>*</sup>
Asian	11,740	107,117 <sup>*</sup>	12,400	105,442 <sup>*</sup>	13,260	99,251 <sup>*</sup>
Other	14,340	88,738 <sup>*</sup>	14,000	89,974 <sup>*</sup>	15,560	95,813 <sup>*</sup>
Unknown	1,780	81,651 <sup>*</sup>	2,360	107,273 <sup>*</sup>	2,280	110,680 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	134,580	88,890 <sup>*</sup>	139,980	89,696 <sup>*</sup>	148,120	91,500 <sup>*</sup>
South	309,140	91,031 <sup>*</sup>	328,040	92,421 <sup>*</sup>	345,080	95,579 <sup>*</sup>
Midwest	170,880	81,949 <sup>R</sup>	173,520	80,015 <sup>R</sup>	175,720	79,793 <sup>R</sup>
West	140,520	93,768 <sup>*</sup>	150,700	94,223 <sup>*</sup>	155,200	92,912 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table F03 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
<b>Total</b>	811,340	88,902
<b>Age</b>	-	-
65-69	167,640	91,307 <sup>R</sup>
70-74	174,700	92,995 <sup>*</sup>
75-79	164,520	93,234 <sup>*</sup>
80-84	147,420	87,531 <sup>*</sup>
85+	157,060	80,018 <sup>*</sup>
<b>Race</b>	-	-
White	724,760	89,446 <sup>R</sup>
Black	37,460	72,011 <sup>*</sup>
Hispanic	16,260	98,545 <sup>*</sup>
Asian	15,080	103,288 <sup>*</sup>
Other	15,120	90,756 <sup>*</sup>
Unknown	2,660	103,906 <sup>*</sup>
<b>Region</b>	-	-
Northeast	142,160	88,573 <sup>*</sup>
South	337,380	92,825 <sup>*</sup>
Midwest	169,660	77,719 <sup>R</sup>
West	162,140	95,175 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.