

Table F05 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Fistula,
2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	6,380	770	6,820	815	6,700	798
Age	-	-	-	-	-	-
65-69	1,660	1,083 ^R	1,320	846 ^R	1,520	941 ^R
70-74	1,960	1,095 [*]	1,840	1,066 [*]	1,640	971 [*]
75-79	1,340	709 [*]	1,500	783 [*]	1,420	752 [*]
80-84	580	371 [*]	1,240	767 [*]	1,000	616 [*]
85+	840	557 [*]	920	592 [*]	1,120	707 [*]
Race	-	-	-	-	-	-
White	5,860	785 ^R	6,080	810 ^R	5,920	798 ^R
Black	440	971 [*]	340	732 [*]	600	1,207 [*]
Hispanic	0	0 [*]	320	2,802 [*]	40	289 [*]
Asian	0	0 [*]	0	0 [*]	20	193 [*]
Other	80	493 [*]	80	433 [*]	120	594 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	1,260	859 [*]	1,680	1,135 [*]	1,380	915
South	2,760	852 [*]	2,760	837	3,040	937
Midwest	1,160	539 ^R	1,480	707 ^R	1,460	683 ^R
West	1,200	842 [*]	900	599 [*]	820	544 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of fistula.

Table F05 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Fistula,
2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	7,340	849	5,980	691	5,540	661
Age	-	-	-	-	-	-
65-69	2,280	1,384 ^R	1,500	883 ^R	1,460	905 ^R
70-74	1,260	729 [*]	1,140	673 [*]	1,260	761 [*]
75-79	1,580	827 [*]	1,400	750 [*]	1,440	826 [*]
80-84	1,140	665 [*]	1,200	713 [*]	860	526 [*]
85+	1,080	654 [*]	740	433 [*]	520	300 [*]
Race	-	-	-	-	-	-
White	6,460	842 ^R	5,560	720 ^R	4,960	658 ^R
Black	500	1,008 [*]	220	460 [*]	300	700 [*]
Hispanic	160	1,015 [*]	140	903 [*]	60	416 [*]
Asian	60	600 [*]	20	212 [*]	120	1,242 [*]
Other	60	318 [*]	40	235 [*]	80	522 [*]
Unknown	100	2,825 [*]	0	0 [*]	20	769 [*]
Region	-	-	-	-	-	-
Northeast	1,280	845	1,560	1,013	1,000	658
South	3,080	913	2,080	612	2,440	740
Midwest	1,880	868 ^R	1,360	621 ^R	1,420	680 ^R
West	1,100	690 [*]	980	643 [*]	680	461 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of fistula.

Table F05 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Fistula,
2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	5,740	676	7,000	789	7,160	787
Age	-	-	-	-	-	-
65-69	1,440	859 ^R	1,860	1,051 ^R	1,920	1,060 ^R
70-74	920	539 [*]	1,460	827 [*]	1,520	837 [*]
75-79	1,660	981 [*]	1,400	780 [*]	1,440	792 [*]
80-84	820	498 [*]	900	533 [*]	980	579 [*]
85+	900	508 [*]	1,380	743 [*]	1,300	662 [*]
Race	-	-	-	-	-	-
White	5,020	659 ^R	6,380	804 ^R	6,300	776 ^R
Black	380	854 [*]	240	488 [*]	600	1,194 [*]
Hispanic	240	1,717 [*]	240	1,550 [*]	140	862 [*]
Asian	40	365 [*]	60	510 [*]	100	749 [*]
Other	40	248 [*]	60	386 [*]	20	123 [*]
Unknown	20	917 [*]	20	909 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	820	542	1,060	679	1,300	803
South	2,480	730 [*]	3,480	980 [*]	3,220	892
Midwest	1,060	508 ^R	1,460	673 ^R	1,940	881 ^R
West	1,380	921 [*]	1,000	625 [*]	700	419 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of fistula.

Table F05 - Physician Office Visits by Female Medicare Beneficiaries with Any Diagnosis of Fistula, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	7,560	828
Age	-	-
65-69	1,940	1,057 ^R
70-74	1,560	830 [*]
75-79	1,700	963 [*]
80-84	1,460	867 [*]
85+	900	459 [*]
Race	-	-
White	6,640	820 ^R
Black	500	961 [*]
Hispanic	60	364 [*]
Asian	140	959 [*]
Other	220	1,321 [*]
Unknown	0	0 [*]
Region	-	-
Northeast	1,220	760
South	3,260	897
Midwest	1,920	880 ^R
West	1,160	681 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of physician office visits by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had a physician office visit for any diagnosis of fistula.