	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	152,720	18,440	155,200	18,534	160,460	19,106
Age	-	-	-	-	-	-
65-69	31,000	20,219 ^R	30,560	19,575 ^R	31,220	19,336 ^R
70-74	32,740	18,293 *	32,640	18,909 *	34,100	20,189 *
75-79	34,540	18,285 *	37,220	19,436 *	35,240	18,673 *
80-84	27,680	17,707 *	28,560	17,658 *	31,820	19,586 *
85+	26,760	17,757 *	26,220	16,875 *	28,080	17,736 *
Race	-	-	-	-	-	-
White	131,820	17,660 ^R	132,280	17,627 ^R	136,260	18,355 ^R
Black	12,700	28,035 *	13,560	29,199 *	14,600	29,376 *
Hispanic	2,340	22,896 *	2,440	21,366 *	2,800	20,231 *
Asian	700	11,006 *	960	12,938 *	1,100	10,618 *
Other	4,640	28,607 *	5,480	29,654 *	5,360	26,535 *
Unknown	520	14,054 *	480	15,094 *	340	9,942 *
Region	-	-	-	-	-	-
Northeast	26,300	17,930 *	27,600	18,649 *	28,240	18,727 *
South	53,100	16,394 *	53,420	16,202 *	51,260	15,806 *
Midwest	48,780	22,674 ^R	48,000	22,914 ^R	52,720	24,647 ^R
West	24,540	17,221 *	26,180	17,435 *	28,240	18,719 *

Table F07 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2002 - 2004

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	170,820	19,751	179,520	20,752	173,640	20,714
Age	-	-	-	-	-	-
65-69	34,960	21,221 ^R	36,280	21,364 ^R	35,180	21,805 ^R
70-74	34,260	19,833 *	37,940	22,383 *	35,240	21,290 *
75-79	37,760	19,772 *	35,980	19,274 *	35,660	20,454 *
80-84	34,180	19,951 *	35,140	20,882 *	32,660	19,983 *
85+	29,660	17,967 *	34,180	20,012 *	34,900	20,099 *
Race	-	-	-	-	-	-
White	146,460	19,094 ^R	157,020	20,324 ^R	152,520	20,244 ^R
Black	14,500	29,222 *	12,900	26,999 *	11,620	27,099 *
Hispanic	3,160	20,051 *	2,600	16,774 *	2,940	20,388 *
Asian	800	8,000 *	1,080	11,465 *	1,380	14,286 *
Other	5,180	27,407 *	5,300	31,140 *	4,680	30,548 *
Unknown	720	20,339 *	620	22,302 *	500	19,231 *
Region	-	-	-	-	-	-
Northeast	30,480	20,127 *	31,620	20,535 *	33,020	21,724 *
South	56,140	16,638 *	55,920	16,458 *	52,240	15,834 *
Midwest	55,600	25,667 ^R	59,060	26,983 ^R	55,600	26,636 ^R
West	28,600	17,945 *	32,920	21,595 *	32,780	22,206 *

Table F07 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries withAny Diagnosis of Urinary Incontinence, 2005 - 2007

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	178,020	20,959	192,880	21,726	203,260	22,332
Age	-	-	-	-	-	-
65-69	37,960	22,633 ^R	41,680	23,545 ^R	44,280	24,451 ^R
70-74	37,880	22,199 *	40,940	23,182 *	40,960	22,545 *
75-79	36,660	21,656 *	40,340	22,476 *	41,420	22,776 *
80-84	31,720	19,273 *	33,740	19,979 *	37,300	22,053 *
85+	33,800	19,079 *	36,180	19,470 *	39,300	20,010 *
Race	-	-	-	-	-	-
White	156,140	20,502 ^R	168,360	21,213 ^R	179,300	22,081 ^R
Black	12,160	27,314 *	14,620	29,752 *	13,440	26,741 *
Hispanic	3,000	21,459 *	2,940	18,992 *	3,260	20,074 *
Asian	1,320	12,044 *	1,320	11,224 *	1,740	13,024 *
Other	5,040	31,188 *	5,220	33,548 *	5,200	32,020 *
Unknown	360	16,514 *	420	19,091 *	320	15,534 *
Region	-	-	-	-	-	-
Northeast	33,080	21,849 *	36,280	23,247 *	37,260	23,017 *
South	56,160	16,537 *	59,060	16,639 *	63,080	17,472 *
Midwest	56,640	27,163 ^R	60,800	28,037 ^R	62,260	28,272 ^R
West	32,140	21,447 *	36,740	22,971 *	40,660	24,341 *

Table F07 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with
Any Diagnosis of Urinary Incontinence, 2008 - 2010

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

 $^{\scriptscriptstyle \mathsf{R}}$ Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

Table F07 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	220,960	24,212
Age	-	-
65-69	49,300	26,852 ^R
70-74	47,620	25,349 *
75-79	42,840	24,277 *
80-84	38,940	23,121 *
85+	42,260	21,530 *
Race	-	-
White	192,640	23,774 ^R
Black	16,180	31,103 *
Hispanic	4,280	25,939 *
Asian	2,040	13,973 *
Other	5,320	31,933 *
Unknown	500	19,531 *
Region	-	-
Northeast	38,420	23,938 *
South	68,720	18,907 *
Midwest	66,980	30,683 ^R
West	46,840	27,495 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.