

Table F08 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
<b>Total</b>	127,480	15,392	129,780	15,499	134,780	16,048
<b>Age</b>	-	-	-	-	-	-
65-69	26,260	17,128 <sup>R</sup>	25,680	16,449 <sup>R</sup>	26,440	16,376 <sup>R</sup>
70-74	27,360	15,287 <sup>*</sup>	27,160	15,734 <sup>*</sup>	28,500	16,874 <sup>*</sup>
75-79	28,280	14,971 <sup>*</sup>	31,160	16,272 <sup>*</sup>	29,200	15,473 <sup>*</sup>
80-84	22,980	14,701 <sup>*</sup>	23,580	14,579 <sup>*</sup>	26,800	16,496 <sup>*</sup>
85+	22,600	14,997 <sup>*</sup>	22,200	14,288 <sup>*</sup>	23,840	15,058 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	109,880	14,721 <sup>R</sup>	111,060	14,800 <sup>R</sup>	114,980	15,489 <sup>R</sup>
Black	10,680	23,576 <sup>*</sup>	10,700	23,040 <sup>*</sup>	11,800	23,742 <sup>*</sup>
Hispanic	1,940	18,982 <sup>*</sup>	2,100	18,389 <sup>*</sup>	2,500	18,064 <sup>*</sup>
Asian	540	8,491 <sup>*</sup>	740	9,973 <sup>*</sup>	940	9,073 <sup>*</sup>
Other	3,980	24,538 <sup>*</sup>	4,760	25,758 <sup>*</sup>	4,260	21,089 <sup>*</sup>
Unknown	460	12,432 <sup>*</sup>	420	13,208 <sup>*</sup>	300	8,772 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	20,100	13,703 <sup>*</sup>	21,900	14,797 <sup>*</sup>	22,640	15,013 <sup>*</sup>
South	45,600	14,078 <sup>*</sup>	45,340	13,751 <sup>*</sup>	44,080	13,592 <sup>*</sup>
Midwest	41,140	19,122 <sup>R</sup>	40,500	19,334 <sup>R</sup>	44,900	20,991 <sup>R</sup>
West	20,640	14,484 <sup>*</sup>	22,040	14,678 <sup>*</sup>	23,160	15,352 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table F08 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
<b>Total</b>	141,900	16,407	148,540	17,171	143,400	17,106
<b>Age</b>	-	-	-	-	-	-
65-69	29,560	17,943 <sup>R</sup>	30,780	18,125 <sup>R</sup>	30,060	18,631 <sup>R</sup>
70-74	27,900	16,151 <sup>*</sup>	31,280	18,454 <sup>*</sup>	28,540	17,243 <sup>*</sup>
75-79	30,880	16,169 <sup>*</sup>	29,040	15,556 <sup>*</sup>	29,320	16,818 <sup>*</sup>
80-84	28,560	16,671 <sup>*</sup>	28,780	17,102 <sup>*</sup>	27,100	16,581 <sup>*</sup>
85+	25,000	15,144 <sup>*</sup>	28,660	16,780 <sup>*</sup>	28,380	16,344 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	122,000	15,905 <sup>R</sup>	130,280	16,863 <sup>R</sup>	125,860	16,706 <sup>R</sup>
Black	11,680	23,539 <sup>*</sup>	10,020	20,971 <sup>*</sup>	9,660	22,528 <sup>*</sup>
Hispanic	2,660	16,878 <sup>*</sup>	2,300	14,839 <sup>*</sup>	2,400	16,644 <sup>*</sup>
Asian	600	6,000 <sup>*</sup>	920	9,767 <sup>*</sup>	1,120	11,594 <sup>*</sup>
Other	4,340	22,963 <sup>*</sup>	4,520	26,557 <sup>*</sup>	4,020	26,240 <sup>*</sup>
Unknown	620	17,514 <sup>*</sup>	500	17,986 <sup>*</sup>	340	13,077 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	24,140	15,940 <sup>*</sup>	25,000	16,236 <sup>*</sup>	25,820	16,987 <sup>*</sup>
South	47,900	14,196 <sup>*</sup>	47,500	13,980 <sup>*</sup>	43,860	13,294 <sup>*</sup>
Midwest	46,500	21,466 <sup>R</sup>	49,220	22,487 <sup>R</sup>	46,160	22,114 <sup>R</sup>
West	23,360	14,657 <sup>*</sup>	26,820	17,594 <sup>*</sup>	27,560	18,670 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table F08 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
<b>Total</b>	146,920	17,297	156,960	17,680	166,120	18,251
<b>Age</b>	-	-	-	-	-	-
65-69	31,280	18,650 <sup>R</sup>	34,980	19,760 <sup>R</sup>	36,400	20,099 <sup>R</sup>
70-74	30,740	18,015 *	33,240	18,822 *	33,480	18,428 *
75-79	29,940	17,687 *	32,900	18,331 *	33,920	18,652 *
80-84	26,100	15,859 *	26,660	15,786 *	29,540	17,465 *
85+	28,860	16,290 *	29,180	15,703 *	32,780	16,690 *
<b>Race</b>	-	-	-	-	-	-
White	128,620	16,889 <sup>R</sup>	136,920	17,252 <sup>R</sup>	147,340	18,145 <sup>R</sup>
Black	10,140	22,776 *	11,520	23,443 *	9,960	19,817 *
Hispanic	2,460	17,597 *	2,400	15,504 *	2,700	16,626 *
Asian	1,020	9,307 *	1,140	9,694 *	1,400	10,479 *
Other	4,400	27,228 *	4,600	29,563 *	4,460	27,463 *
Unknown	280	12,844 *	380	17,273 *	260	12,621 *
<b>Region</b>	-	-	-	-	-	-
Northeast	26,040	17,199 *	27,760	17,788 *	30,020	18,545 *
South	47,080	13,863 *	49,540	13,957 *	52,620	14,575 *
Midwest	46,900	22,492 <sup>R</sup>	49,900	23,010 <sup>R</sup>	50,700	23,022 <sup>R</sup>
West	26,900	17,950 *	29,760	18,607 *	32,780	19,624 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.

Table F08 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
<b>Total</b>	178,280	19,535
<b>Age</b>	-	-
65-69	40,620	22,124 <sup>R</sup>
70-74	38,180	20,324 *
75-79	33,720	19,109 *
80-84	31,720	18,834 *
85+	34,040	17,343 *
<b>Race</b>	-	-
White	155,880	19,238 <sup>R</sup>
Black	12,360	23,760 *
Hispanic	3,580	21,697 *
Asian	1,540	10,548 *
Other	4,560	27,371 *
Unknown	360	14,063 *
<b>Region</b>	-	-
Northeast	30,140	18,779 *
South	55,940	15,391 *
Midwest	53,960	24,718 <sup>R</sup>
West	38,240	22,447 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of stress urinary incontinence.