

Table F10 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Fistula, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	1,600	193	1,880	225	1,740	207
Age	-	-	-	-	-	-
65-69	340	222 ^R	520	333 ^R	560	347 ^R
70-74	700	391 [*]	380	220 [*]	320	190 [*]
75-79	320	169 [*]	280	146 [*]	460	244 [*]
80-84	140	90 [*]	420	260 [*]	220	135 [*]
85+	100	66 [*]	280	180 [*]	180	114 [*]
Race	-	-	-	-	-	-
White	1,360	182 ^R	1,440	192 ^R	1,300	175 ^R
Black	220	486 [*]	420	904 [*]	400	805 [*]
Hispanic	0	0 [*]	20	175 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	20	123 [*]	0	0 [*]	40	198 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	180	123 [*]	380	257	360	239 [*]
South	520	161 [*]	500	152 [*]	460	142
Midwest	500	232 ^R	580	277 ^R	320	150 ^R
West	400	281 [*]	420	280 [*]	600	398 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of fistula.

Table F10 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Fistula, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	2,140	247	1,700	197	1,780	212
Age	-	-	-	-	-	-
65-69	740	449 ^R	380	224 ^R	460	285 ^R
70-74	520	301 [*]	520	307 [*]	540	326 [*]
75-79	280	147 [*]	280	150 [*]	300	172 [*]
80-84	320	187 [*]	320	190 [*]	280	171 [*]
85+	280	170 [*]	200	117 [*]	200	115 [*]
Race	-	-	-	-	-	-
White	1,780	232 ^R	1,280	166 ^R	1,420	189 ^R
Black	240	484 [*]	400	837 [*]	240	560 [*]
Hispanic	80	508 [*]	20	129 [*]	40	277 [*]
Asian	0	0 [*]	0	0 [*]	20	207 [*]
Other	20	106 [*]	0	0 [*]	60	392 [*]
Unknown	20	565 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	500	330 [*]	340	221 [*]	520	342 [*]
South	680	202 [*]	500	147 [*]	600	182 [*]
Midwest	580	268 ^R	620	283 ^R	340	163 ^R
West	380	238 [*]	240	157 [*]	320	217 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of fistula.

Table F10 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Fistula, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	2,160	254	2,140	241	2,020	222
Age	-	-	-	-	-	-
65-69	400	239 ^R	780	441 ^R	560	309 ^R
70-74	760	445 [*]	560	317 [*]	580	319 [*]
75-79	500	295 [*]	300	167 [*]	280	154 [*]
80-84	240	146 [*]	240	142 [*]	360	213 [*]
85+	260	147 [*]	260	140 [*]	240	122 [*]
Race	-	-	-	-	-	-
White	1,980	260 ^R	1,820	229 ^R	1,620	200 ^R
Black	160	359 [*]	240	488 [*]	260	517 [*]
Hispanic	0	0 [*]	60	388 [*]	40	246 [*]
Asian	0	0 [*]	0	0 [*]	100	749 [*]
Other	0	0 [*]	20	129 [*]	0	0 [*]
Unknown	20	917 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	440	291 [*]	620	397 [*]	280	173 [*]
South	540	159 [*]	660	186 [*]	480	133 [*]
Midwest	780	374 ^R	640	295 ^R	860	391 ^R
West	400	267 [*]	220	138 [*]	400	240 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of fistula.

Table F10 - Ambulatory Visits (Other than Physician Visits) by Female Medicare Beneficiaries with Any Diagnosis of Fistula, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	2,360	259
Age	-	-
65-69	760	414 ^R
70-74	480	256 *
75-79	620	351 *
80-84	240	143 *
85+	260	133 *
Race	-	-
White	2,000	247 ^R
Black	180	346 *
Hispanic	60	364 *
Asian	60	411 *
Other	0	0 *
Unknown	60	2,344 *
Region	-	-
Northeast	340	212 *
South	800	220 *
Midwest	820	376 ^R
West	400	235 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of ambulatory visits (other than physician visits) by female adult Medicare beneficiaries with any diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who had an ambulatory visit (other than a physician visit) for any diagnosis of fistula.