

Table F12 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
<b>Total</b>	41,880	5,057	39,400	4,705	37,680	4,487
<b>Age</b>	-	-	-	-	-	-
65-69	11,080	7,227 <sup>R</sup>	10,320	6,610 <sup>R</sup>	10,400	6,441 <sup>R</sup>
70-74	11,220	6,269 <sup>*</sup>	10,700	6,199 <sup>*</sup>	9,220	5,459 <sup>*</sup>
75-79	9,460	5,008 <sup>*</sup>	9,420	4,919 <sup>*</sup>	8,940	4,737 <sup>*</sup>
80-84	6,200	3,966 <sup>*</sup>	5,660	3,499 <sup>*</sup>	5,620	3,459 <sup>*</sup>
85+	3,920	2,601 <sup>*</sup>	3,300	2,124 <sup>*</sup>	3,500	2,211 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	38,600	5,171 <sup>R</sup>	36,140	4,816 <sup>R</sup>	34,660	4,669 <sup>R</sup>
Black	1,780	3,929 <sup>*</sup>	1,580	3,402 <sup>*</sup>	1,480	2,978 <sup>*</sup>
Hispanic	440	4,305 <sup>*</sup>	560	4,904 <sup>*</sup>	380	2,746 <sup>*</sup>
Asian	220	3,459 <sup>*</sup>	200	2,695 <sup>*</sup>	100	965 <sup>*</sup>
Other	520	3,206 <sup>*</sup>	760	4,113 <sup>*</sup>	900	4,455 <sup>*</sup>
Unknown	320	8,649 <sup>*</sup>	160	5,031 <sup>*</sup>	160	4,678 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	6,720	4,581 <sup>*</sup>	6,280	4,243 <sup>*</sup>	5,400	3,581 <sup>*</sup>
South	17,300	5,341 <sup>*</sup>	16,460	4,992 <sup>*</sup>	15,940	4,915 <sup>*</sup>
Midwest	10,960	5,094 <sup>R</sup>	9,780	4,669 <sup>R</sup>	10,160	4,750 <sup>R</sup>
West	6,900	4,842 <sup>*</sup>	6,880	4,582 <sup>*</sup>	6,180	4,097 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table F12 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
<b>Total</b>	36,560	4,227	36,380	4,205	31,340	3,739
<b>Age</b>	-	-	-	-	-	-
65-69	9,900	6,010 <sup>R</sup>	9,720	5,724 <sup>R</sup>	8,660	5,368 <sup>R</sup>
70-74	9,420	5,453 <sup>*</sup>	9,720	5,735 <sup>*</sup>	8,600	5,196 <sup>*</sup>
75-79	8,360	4,377 <sup>*</sup>	8,360	4,478 <sup>*</sup>	6,720	3,855 <sup>*</sup>
80-84	5,340	3,117 <sup>*</sup>	5,680	3,375 <sup>*</sup>	4,540	2,778 <sup>*</sup>
85+	3,540	2,144 <sup>*</sup>	2,900	1,698 <sup>*</sup>	2,820	1,624 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	33,840	4,412 <sup>R</sup>	33,200	4,297 <sup>R</sup>	28,720	3,812 <sup>R</sup>
Black	1,260	2,539 <sup>*</sup>	1,520	3,181 <sup>*</sup>	1,200	2,799 <sup>*</sup>
Hispanic	340	2,157 <sup>*</sup>	700	4,516 <sup>*</sup>	420	2,913 <sup>*</sup>
Asian	180	1,800 <sup>*</sup>	100	1,062 <sup>*</sup>	220	2,277 <sup>*</sup>
Other	640	3,386 <sup>*</sup>	700	4,113 <sup>*</sup>	580	3,786 <sup>*</sup>
Unknown	300	8,475 <sup>*</sup>	160	5,755 <sup>*</sup>	200	7,692 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	5,540	3,658 <sup>*</sup>	5,660	3,676 <sup>*</sup>	4,900	3,224 <sup>*</sup>
South	14,780	4,380 <sup>*</sup>	14,360	4,226 <sup>*</sup>	13,240	4,013 <sup>*</sup>
Midwest	9,740	4,496 <sup>R</sup>	9,940	4,541 <sup>R</sup>	7,940	3,804 <sup>R</sup>
West	6,500	4,078 <sup>*</sup>	6,420	4,212 <sup>*</sup>	5,260	3,563 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table F12 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
<b>Total</b>	30,800	3,626	27,640	3,113	24,600	2,703
<b>Age</b>	-	-	-	-	-	-
65-69	9,740	5,807 <sup>R</sup>	8,980	5,073 <sup>R</sup>	7,740	4,274 <sup>R</sup>
70-74	7,720	4,524 <sup>*</sup>	6,540	3,703 <sup>*</sup>	6,620	3,644 <sup>*</sup>
75-79	6,700	3,958 <sup>*</sup>	5,840	3,254 <sup>*</sup>	4,740	2,606 <sup>*</sup>
80-84	3,900	2,370 <sup>*</sup>	4,000	2,369 <sup>*</sup>	3,280	1,939 <sup>*</sup>
85+	2,740	1,547 <sup>*</sup>	2,280	1,227 <sup>*</sup>	2,220	1,130 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	28,020	3,679 <sup>R</sup>	25,120	3,165 <sup>R</sup>	22,560	2,778 <sup>R</sup>
Black	1,240	2,785 <sup>*</sup>	1,380	2,808 <sup>*</sup>	1,160	2,308 <sup>*</sup>
Hispanic	440	3,147 <sup>*</sup>	460	2,972 <sup>*</sup>	300	1,847 <sup>*</sup>
Asian	160	1,460 <sup>*</sup>	240	2,041 <sup>v</sup>	180	1,347 <sup>*</sup>
Other	820	5,074 <sup>*</sup>	360	2,314 <sup>*</sup>	380	2,340 <sup>*</sup>
Unknown	120	5,505 <sup>*</sup>	80	3,636 <sup>*</sup>	20	971 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	5,120	3,382 <sup>*</sup>	4,960	3,178 <sup>*</sup>	3,760	2,323 <sup>*</sup>
South	12,800	3,769 <sup>*</sup>	12,260	3,454 <sup>*</sup>	10,380	2,875 <sup>*</sup>
Midwest	7,320	3,511 <sup>R</sup>	6,040	2,785 <sup>R</sup>	6,160	2,797 <sup>R</sup>
West	5,560	3,710 <sup>*</sup>	4,380	2,739 <sup>*</sup>	4,300	2,574 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table F12 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
<b>Total</b>	18,840	2,064
<b>Age</b>	-	-
65-69	5,280	2,876 <sup>R</sup>
70-74	4,980	2,651 *
75-79	3,860	2,188 *
80-84	2,640	1,568 *
85+	2,080	1,060 *
<b>Race</b>	-	-
White	17,160	2,118 <sup>R</sup>
Black	1,040	1,999 *
Hispanic	200	1,212 *
Asian	160	1,096 *
Other	200	1,201 *
Unknown	80	3,125 *
<b>Region</b>	-	-
Northeast	3,240	2,019 *
South	7,060	1,942 *
Midwest	5,060	2,318 <sup>R</sup>
West	3,480	2,043 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of hospitalizations by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.