

Table F13 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	38,020	4,591	35,840	4,280	34,280	4,082
Age	-	-	-	-	-	-
65-69	10,340	6,744 ^R	9,680	6,200 ^R	9,680	5,995 ^R
70-74	10,460	5,844 *	10,060	5,828 *	8,680	5,139 *
75-79	8,620	4,563 *	8,480	4,428 *	8,200	4,345 *
80-84	5,320	3,403 *	4,980	3,079 *	5,020	3,090 *
85+	3,280	2,177 *	2,640	1,699 *	2,700	1,705 *
Race	-	-	-	-	-	-
White	35,180	4,713 ^R	32,980	4,395 ^R	31,740	4,276 ^R
Black	1,360	3,002 *	1,280	2,756 *	1,180	2,374 *
Hispanic	440	4,305 *	500	4,378 *	320	2,312 *
Asian	220	3,459 *	200	2,695 *	80	772 *
Other	520	3,206 *	720	3,896 *	820	4,059 *
Unknown	300	8,108 *	160	5,031 *	140	4,094 *
Region	-	-	-	-	-	-
Northeast	5,780	3,941 *	5,340	3,608 *	4,700	3,117 *
South	15,980	4,934 *	15,200	4,610 *	14,480	4,465 *
Midwest	9,880	4,592 ^R	8,840	4,220 ^R	9,240	4,320 ^R
West	6,380	4,477 *	6,460	4,302 *	5,860	3,884 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table F13 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	33,180	3,837	33,600	3,884	28,660	3,419
Age	-	-	-	-	-	-
65-69	9,140	5,548 ^R	9,020	5,312 ^R	8,000	4,959 ^R
70-74	8,740	5,060 [*]	9,200	5,428 [*]	7,960	4,809 [*]
75-79	7,620	3,990 [*]	7,760	4,157 [*]	6,240	3,579 [*]
80-84	4,720	2,755 [*]	5,100	3,031 [*]	4,020	2,460 [*]
85+	2,960	1,793 [*]	2,520	1,475 [*]	2,440	1,405 [*]
Race	-	-	-	-	-	-
White	30,840	4,021 ^R	30,760	3,982 ^R	26,340	3,496 ^R
Black	1,040	2,096 [*]	1,380	2,888 [*]	1,000	2,332 [*]
Hispanic	280	1,777 [*]	660	4,258 [*]	400	2,774 [*]
Asian	160	1,600 [*]	80	849 [*]	160	1,656 [*]
Other	560	2,963 [*]	580	3,408 [*]	560	3,655 [*]
Unknown	300	8,475 [*]	140	5,036 [*]	200	7,692 [*]
Region	-	-	-	-	-	-
Northeast	4,780	3,156 [*]	5,020	3,260 [*]	4,420	2,908 [*]
South	13,480	3,995 [*]	13,360	3,932 [*]	12,180	3,692 [*]
Midwest	9,000	4,155 ^R	9,280	4,240 ^R	7,360	3,526 ^R
West	5,920	3,714 [*]	5,940	3,897 [*]	4,700	3,184 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table F13 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	27,200	3,202	24,600	2,771	21,560	2,369
Age	-	-	-	-	-	-
65-69	9,020	5,378 ^R	8,000	4,519 ^R	6,960	3,843 ^R
70-74	7,060	4,137 [*]	6,080	3,443 [*]	5,840	3,214 [*]
75-79	5,780	3,415 [*]	5,280	2,942 [*]	4,220	2,321 [*]
80-84	3,180	1,932 [*]	3,440	2,037 [*]	2,800	1,655 [*]
85+	2,160	1,219 [*]	1,800	969 [*]	1,740	886 [*]
Race	-	-	-	-	-	-
White	24,700	3,243 ^R	22,460	2,830 ^R	19,920	2,453 ^R
Black	1,040	2,336 [*]	1,180	2,401 [*]	900	1,791 [*]
Hispanic	400	2,861 [*]	320	2,067 [*]	240	1,478 [*]
Asian	160	1,460 [*]	220	1,871 [*]	160	1,198 [*]
Other	800	4,951 [*]	340	2,185 [*]	340	2,094 [*]
Unknown	100	4,587 [*]	80	3,636 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	4,320	2,853 [*]	4,360	2,794 [*]	3,240	2,002 [*]
South	11,420	3,363 [*]	10,940	3,082 [*]	9,120	2,526 [*]
Midwest	6,480	3,108 ^R	5,320	2,453 ^R	5,220	2,370 ^R
West	4,980	3,323 [*]	3,980	2,488 [*]	3,980	2,383 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table F13 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	15,440	1,692
Age	-	-
65-69	4,400	2,397 ^R
70-74	4,300	2,289 [*]
75-79	3,280	1,859 [*]
80-84	1,980	1,176 [*]
85+	1,480	754 [*]
Race	-	-
White	14,220	1,755 ^R
Black	720	1,384 [*]
Hispanic	160	970 [*]
Asian	120	822 [*]
Other	180	1,080 [*]
Unknown	40	1,563 [*]
Region	-	-
Northeast	2,460	1,533 [*]
South	5,740	1,579 [*]
Midwest	4,200	1,924 ^R
West	3,040	1,785 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.