

Table F14 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urge Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	680	82	560	67	520	62
Age	-	-	-	-	-	-
65-69	20	13 ^R	40	26 ^R	60	37 ^R
70-74	80	45 [*]	100	58 [*]	120	71 [*]
75-79	180	95 [*]	220	115 [*]	100	53 [*]
80-84	220	141 [*]	120	74 [*]	80	49 [*]
85+	180	119 [*]	80	52 [*]	160	101 [*]
Race	-	-	-	-	-	-
White	620	83 ^R	520	69 ^R	520	70 ^R
Black	60	133 [*]	40	86 [*]	0	0 [*]
Hispanic	0	0 [*]	0	0 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	120	82 [*]	160	108	100	66
South	160	49 [*]	160	49 [*]	240	74
Midwest	360	167 ^R	220	105 ^R	140	66 ^R
West	40	28 [*]	20	13 [*]	40	27 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

Table F14 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urge Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	520	60	440	51	420	50
Age	-	-	-	-	-	-
65-69	20	12 ^R	40	24 ^R	60	37 ^R
70-74	60	35 [*]	40	24 [*]	80	48 [*]
75-79	120	63 [*]	140	75 [*]	100	57 [*]
80-84	200	117 [*]	80	48 [*]	100	61 [*]
85+	120	73 [*]	140	82 [*]	80	46 [*]
Race	-	-	-	-	-	-
White	480	63 ^R	400	52 ^R	400	53 ^R
Black	20	40 [*]	20	42 [*]	0	0 [*]
Hispanic	20	127 [*]	0	0 [*]	20	139 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	20	719 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	120	79 [*]	80	52 [*]	120	79 [*]
South	160	47	100	29 [*]	140	42
Midwest	120	55 ^R	160	73 ^R	80	38 ^R
West	120	75 [*]	100	66 [*]	80	54 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

Table F14 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urge Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	520	61	420	47	400	44
Age	-	-	-	-	-	-
65-69	120	72 ^R	100	57 ^R	60	33 ^R
70-74	80	47 [*]	80	45 [*]	120	66 [*]
75-79	40	24 [*]	60	33 [*]	80	44 [*]
80-84	160	97 [*]	120	71 [*]	40	24 [*]
85+	120	68 [*]	60	32 [*]	100	51 [*]
Race	-	-	-	-	-	-
White	440	58 ^R	360	45 ^R	380	47 ^R
Black	60	135 [*]	40	81 [*]	20	40 [*]
Hispanic	20	143 [*]	20	129 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	120	79 [*]	60	38 [*]	100	62 [*]
South	280	82 [*]	120	34 [*]	180	50 [*]
Midwest	80	38 ^R	180	83 ^R	80	36 ^R
West	40	27 [*]	60	38 [*]	40	24 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

Table F14 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Urge Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	540	59
Age	-	-
65-69	100	55 ^R
70-74	40	21 [*]
75-79	140	79 [*]
80-84	140	83 [*]
85+	120	61 [*]
Race	-	-
White	420	52 ^R
Black	80	154 [*]
Hispanic	40	242 [*]
Asian	0	0 [*]
Other	0	0 [*]
Unknown	0	0 [*]
Region	-	-
Northeast	180	112 [*]
South	140	39 [*]
Midwest	180	83 ^R
West	40	24 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urge urinary incontinence.