

Table F15 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Fistula,
2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	2,980	360	2,900	346	2,780	331
Age	-	-	-	-	-	-
65-69	700	457 ^R	600	384 ^R	660	409 ^R
70-74	680	380 [*]	540	313 [*]	400	237 [*]
75-79	620	328 [*]	720	376 [*]	640	339 [*]
80-84	620	397 [*]	500	309 [*]	500	308 [*]
85+	360	239 [*]	540	348 [*]	580	366 [*]
Race	-	-	-	-	-	-
White	2,620	351 ^R	2,580	344 ^R	2,340	315 ^R
Black	340	751 [*]	220	474 [*]	260	523 [*]
Hispanic	0	0 [*]	60	525 [*]	60	434 [*]
Asian	0	0 [*]	0	0 [*]	20	193 [*]
Other	0	0 [*]	40	217 [*]	80	396 [*]
Unknown	20	541 [*]	0	0 [*]	20	585 [*]
Region	-	-	-	-	-	-
Northeast	760	518 [*]	720	487 [*]	540	358
South	1,120	346 [*]	1,080	328	1,200	370
Midwest	680	316 ^R	700	334 ^R	760	355 ^R
West	420	295 [*]	400	266 [*]	280	186 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

Table F15 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Fistula,
2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	2,820	326	2,240	259	2,180	260
Age	-	-	-	-	-	-
65-69	740	449 ^R	620	365 ^R	600	372 ^R
70-74	620	359 [*]	440	260 [*]	540	326 [*]
75-79	620	325 [*]	460	246 [*]	380	218 [*]
80-84	420	245 [*]	500	297 [*]	380	233 [*]
85+	420	254 [*]	220	129 [*]	280	161 [*]
Race	-	-	-	-	-	-
White	2,480	323 ^R	1,960	254 ^R	1,900	252 ^R
Black	200	403 [*]	120	251 [*]	200	466 [*]
Hispanic	40	254 [*]	40	258 [*]	0	0 [*]
Asian	20	200 [*]	20	212 [*]	60	621 [*]
Other	80	423 [*]	100	588 [*]	20	131 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	640	423 [*]	560	364 [*]	340	224
South	1,120	332 [*]	840	247 [*]	900	273 [*]
Midwest	600	277 ^R	480	219 ^R	460	220 ^R
West	460	289 [*]	360	236 [*]	480	325 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

Table F15 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Fistula,
2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	3,000	353	2,560	288	2,520	277
Age	-	-	-	-	-	-
65-69	580	346 ^R	880	497 ^R	700	387 ^R
70-74	580	340 [*]	380	215 [*]	600	330 [*]
75-79	840	496 [*]	480	267 [*]	440	242 [*]
80-84	560	340 [*]	400	237 [*]	400	237 [*]
85+	440	248 [*]	420	226 [*]	380	194 [*]
Race	-	-	-	-	-	-
White	2,800	368 ^R	2,240	282 ^R	2,160	266 ^R
Black	140	315 [*]	160	326 [*]	240	478 [*]
Hispanic	20	143 [*]	120	775 [*]	60	370 [*]
Asian	0	0 [*]	20	170 [*]	20	150 [*]
Other	20	124 [*]	20	129 [*]	20	123 [*]
Unknown	20	917 [*]	0	0 [*]	20	971 [*]
Region	-	-	-	-	-	-
Northeast	660	436 [*]	520	333 [*]	420	260 [*]
South	1,060	312 [*]	1,180	333 [*]	1,040	288 [*]
Midwest	760	365 ^R	540	249 ^R	820	372 ^R
West	520	347 [*]	320	200 [*]	240	144 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

Table F15 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Fistula, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	2,800	307
Age	-	-
65-69	780	425 ^R
70-74	580	309 [*]
75-79	440	249 [*]
80-84	520	309 [*]
85+	480	245 [*]
Race	-	-
White	2,500	309 ^R
Black	220	423 [*]
Hispanic	0	0 [*]
Asian	20	137 [*]
Other	20	120 [*]
Unknown	40	1,563 ^v
Region	-	-
Northeast	600	374 [*]
South	1,160	319
Midwest	660	302 ^R
West	380	223 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of fistula.