

Table F16 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
<b>Total</b>	200	24	100	12	100	12
<b>Age</b>	-	-	-	-	-	-
65-69	20	13 <sup>R</sup>	0	0 <sup>R</sup>	0	0 <sup>R</sup>
70-74	0	0 <sup>*</sup>	0	0 <sup>*</sup>	20	12 <sup>*</sup>
75-79	40	21 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
80-84	40	26 <sup>*</sup>	60	37 <sup>*</sup>	20	12 <sup>*</sup>
85+	100	66 <sup>*</sup>	40	26 <sup>*</sup>	60	38 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	180	24 <sup>R</sup>	60	8 <sup>R</sup>	60	8 <sup>R</sup>
Black	20	44 <sup>*</sup>	40	86 <sup>*</sup>	40	81 <sup>*</sup>
Hispanic	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Asian	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Other	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Unknown	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	60	41 <sup>*</sup>	60	41 <sup>*</sup>	60	40 <sup>*</sup>
South	40	12 <sup>*</sup>	20	6	20	6
Midwest	40	19 <sup>R</sup>	20	10 <sup>R</sup>	20	9 <sup>R</sup>
West	60	42 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table F16 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
<b>Total</b>	40	5	100	12	80	10
<b>Age</b>	-	-	-	-	-	-
65-69	0	0 <sup>R</sup>	40	24 <sup>R</sup>	0	0 <sup>R</sup>
70-74	0	0 <sup>*</sup>	40	24 <sup>*</sup>	20	12 <sup>*</sup>
75-79	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
80-84	0	0 <sup>*</sup>	0	0 <sup>*</sup>	40	25 <sup>*</sup>
85+	40	24 <sup>*</sup>	20	12 <sup>*</sup>	20	12 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	40	5 <sup>R</sup>	80	10 <sup>R</sup>	80	11 <sup>R</sup>
Black	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Hispanic	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Asian	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Other	0	0 <sup>*</sup>	20	118 <sup>*</sup>	0	0 <sup>*</sup>
Unknown	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	0	0 <sup>*</sup>	0	0 <sup>*</sup>	20	13
South	20	6	60	18 <sup>*</sup>	20	6 <sup>*</sup>
Midwest	20	9 <sup>R</sup>	20	9 <sup>R</sup>	40	19 <sup>R</sup>
West	0	0 <sup>*</sup>	20	13 <sup>*</sup>	0	0 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table F16 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
<b>Total</b>	80	9	60	7	120	13
<b>Age</b>	-	-	-	-	-	-
65-69	20	12 <sup>R</sup>	0	0 <sup>R</sup>	20	11 <sup>R</sup>
70-74	0	0 <sup>*</sup>	0	0 <sup>*</sup>	60	33 <sup>*</sup>
75-79	40	24 <sup>*</sup>	20	11 <sup>*</sup>	0	0 <sup>*</sup>
80-84	0	0 <sup>*</sup>	40	24 <sup>*</sup>	40	24 <sup>*</sup>
85+	20	11 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	80	11 <sup>R</sup>	60	8 <sup>R</sup>	100	12 <sup>R</sup>
Black	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Hispanic	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Asian	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
Other	0	0 <sup>*</sup>	0	0 <sup>*</sup>	20	123 <sup>*</sup>
Unknown	0	0 <sup>*</sup>	0	0 <sup>*</sup>	0	0 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	20	13 <sup>*</sup>	20	13 <sup>*</sup>	0	0 <sup>*</sup>
South	40	12 <sup>*</sup>	20	6 <sup>*</sup>	40	11 <sup>*</sup>
Midwest	0	0 <sup>R</sup>	0	0 <sup>R</sup>	40	18 <sup>R</sup>
West	20	13 <sup>*</sup>	20	13 <sup>*</sup>	40	24 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table F16 - Hospitalizations by Female Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
<b>Total</b>	60	7
<b>Age</b>	-	-
65-69	0	0 <sup>R</sup>
70-74	60	32 <sup>*</sup>
75-79	0	0 <sup>*</sup>
80-84	0	0 <sup>*</sup>
85+	0	0 <sup>*</sup>
<b>Race</b>	-	-
White	20	3 <sup>R</sup>
Black	20	38 <sup>*</sup>
Hispanic	0	0 <sup>*</sup>
Asian	20	137 <sup>*</sup>
Other	0	0 <sup>*</sup>
Unknown	0	0 <sup>*</sup>
<b>Region</b>	-	-
Northeast	0	0 <sup>*</sup>
South	20	6
Midwest	20	9 <sup>R</sup>
West	20	12 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.