

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2002 - 2003

	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	62,500	7,546.3	60,320	7,203.6
All Procedures	62,500	7,546.3	60,320	7,203.6
Age	-	-	-	-
65-69	17,260	11,257.5	16,740	10,722.5
70-74	18,000	10,057.0	16,640	9,639.7
75-79	14,780	7,824.2	15,860	8,282.0
80-84	8,280	5,296.8	7,940	4,909.1
85+	4,180	2,773.7	3,140	2,020.9
Race	-	-	-	-
White	58,800	7,877.6	56,920	7,585.1
Black	1,340	2,958.1	1,020	2,196.4
Hispanic	740	7,240.7	1,060	9,282.0
Asian	440	6,918.2	340	4,582.2
Other	660	4,069.1	680	3,679.7
Unknown	520	14,054.1	300	9,434.0
Region	-	-	-	-
Northeast	8,900	6,067.6	7,980	5,391.9
South	26,340	8,132.1	26,280	7,970.4
Midwest	16,300	7,576.5	15,340	7,322.9
West	10,960	7,691.2	10,720	7,139.1

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2002 - 2003

	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	62,500	7,546.3	60,320	7,203.6
Pubovaginal Sling	31,020	3,745.4	34,360	4,103.4
Age	-	-	-	-
65-69	9,220	6,013.6	10,820	6,930.6
70-74	9,280	5,184.9	9,500	5,503.4
75-79	7,840	4,150.3	8,820	4,605.7
80-84	3,440	2,200.6	4,060	2,510.2
85+	1,240	822.8	1,160	746.6
Race	-	-	-	-
White	29,220	3,914.7	32,340	4,309.6
Black	700	1,545.3	660	1,421.2
Hispanic	340	3,326.8	660	5,779.3
Asian	200	3,144.7	140	1,886.8
Other	260	1,603.0	420	2,272.7
Unknown	300	8,108.1	140	4,402.5
Region	-	-	-	-
Northeast	4,180	2,849.7	4,120	2,783.8
South	13,400	4,137.1	15,160	4,597.8
Midwest	8,060	3,746.4	8,800	4,200.9
West	5,380	3,775.4	6,280	4,182.2

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2002 - 2003

	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	62,500	7,546.3	60,320	7,203.6
Collagen Injection	14,940	1,803.9	12,900	1,540.6
Age	-	-	-	-
65-69	2,780	1,813.2	1,960	1,255.4
70-74	3,740	2,089.6	3,240	1,877.0
75-79	3,460	1,831.7	3,860	2,015.7
80-84	2,820	1,804.0	2,380	1,471.5
85+	2,140	1,420.0	1,460	939.6
Race	-	-	-	-
White	14,300	1,915.8	12,500	1,665.7
Black	220	485.7	100	215.3
Hispanic	80	782.8	120	1,050.8
Asian	0	0.0	0	0.0
Other	200	1,233.0	60	324.7
Unknown	140	3,783.8	120	3,773.6
Region	-	-	-	-
Northeast	2,840	1,936.2	2,220	1,500.0
South	5,800	1,790.7	5,420	1,643.8
Midwest	4,220	1,961.5	3,280	1,565.8
West	2,080	1,459.6	1,980	1,318.6

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2002 - 2003

	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	62,500	7,546.3	60,320	7,203.6
Sacral Nerve Stimulation	940	113.5	1,240	148.1
Age	-	-	-	-
65-69	360	234.8	340	217.8
70-74	200	111.7	240	139.0
75-79	280	148.2	300	156.7
80-84	20	12.8	260	160.8
85+	80	53.1	100	64.4
Race	-	-	-	-
White	940	125.9	1,140	151.9
Black	0	0.0	0	0.0
Hispanic	0	0.0	40	350.3
Asian	0	0.0	40	539.1
Other	0	0.0	0	0.0
Unknown	0	0.0	20	628.9
Region	-	-	-	-
Northeast	100	68.2	200	135.1
South	360	111.1	420	127.4
Midwest	380	176.6	400	190.9
West	100	70.2	220	146.5

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2002 - 2003

	2002		2003	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	62,500	7,546.3	60,320	7,203.6
Other Procedures	15,600	1,883.6	11,820	1,411.6
Age	-	-	-	-
65-69	4,900	3,195.9	3,620	2,318.7
70-74	4,780	2,670.7	3,660	2,120.3
75-79	3,200	1,694.0	2,880	1,503.9
80-84	2,000	1,279.4	1,240	766.7
85+	720	477.8	420	270.3
Race	-	-	-	-
White	14,340	1,921.2	10,940	1,457.9
Black	420	927.2	260	559.9
Hispanic	320	3,131.1	240	2,101.6
Asian	240	3,773.6	160	2,156.3
Other	200	1,233.0	200	1,082.3
Unknown	80	2,162.2	20	628.9
Region	-	-	-	-
Northeast	1,780	1,213.5	1,440	973.0
South	6,780	2,093.2	5,280	1,601.4
Midwest	3,640	1,691.9	2,860	1,365.3
West	3,400	2,386.0	2,240	1,491.7

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2004 - 2005

	2004		2005	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	64,540	7,684.6	63,800	7,376.9
All Procedures	64,540	7,684.6	63,800	7,376.9
Age	-	-	-	-
65-69	18,720	11,594.2	19,440	11,800.4
70-74	17,740	10,503.3	16,480	9,540.3
75-79	15,680	8,308.6	14,720	7,707.6
80-84	8,580	5,281.3	8,700	5,078.2
85+	3,820	2,412.8	4,460	2,701.7
Race	-	-	-	-
White	61,300	8,257.7	60,360	7,869.2
Black	1,000	2,012.1	1,060	2,136.2
Hispanic	1,160	8,381.5	800	5,076.1
Asian	240	2,316.6	200	2,000.0
Other	460	2,277.2	840	4,444.4
Unknown	380	11,111.1	540	15,254.2
Region	-	-	-	-
Northeast	8,540	5,663.1	8,380	5,533.5
South	27,740	8,553.8	27,240	8,073.0
Midwest	17,440	8,153.3	17,040	7,866.3
West	10,820	7,172.2	11,140	6,989.6

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2004 - 2005

	2004		2005	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	64,540	7,684.6	63,800	7,376.9
Pubovaginal Sling	37,860	4,507.9	38,940	4,502.5
Age	-	-	-	-
65-69	12,380	7,667.5	12,800	7,769.8
70-74	10,660	6,311.4	10,820	6,263.7
75-79	8,680	4,599.4	8,980	4,702.1
80-84	4,620	2,843.8	4,640	2,708.4
85+	1,520	960.1	1,700	1,029.8
Race	-	-	-	-
White	35,860	4,830.7	36,660	4,779.4
Black	480	965.8	700	1,410.7
Hispanic	860	6,213.9	480	3,045.7
Asian	200	1,930.5	120	1,200.0
Other	280	1,386.1	620	3,280.4
Unknown	180	5,263.2	360	10,169.5
Region	-	-	-	-
Northeast	4,520	2,997.3	4,680	3,090.3
South	16,260	5,013.9	16,760	4,967.1
Midwest	10,260	4,796.6	10,320	4,764.1
West	6,820	4,520.7	7,180	4,505.0

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2004 - 2005

	2004		2005	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	64,540	7,684.6	63,800	7,376.9
Collagen Injection	13,580	1,616.9	12,400	1,433.8
Age	-	-	-	-
65-69	2,260	1,399.7	2,420	1,469.0
70-74	3,300	1,953.8	2,280	1,319.9
75-79	3,880	2,056.0	3,000	1,570.8
80-84	2,460	1,514.2	2,540	1,482.6
85+	1,680	1,061.1	2,160	1,308.5
Race	-	-	-	-
White	13,040	1,756.6	12,100	1,577.5
Black	180	362.2	140	282.1
Hispanic	160	1,156.1	60	380.7
Asian	0	0.0	20	200.0
Other	80	396.0	80	423.3
Unknown	120	3,508.8	0	0.0
Region	-	-	-	-
Northeast	2,260	1,498.7	2,160	1,426.3
South	5,620	1,733.0	5,180	1,535.2
Midwest	4,060	1,898.1	3,320	1,532.6
West	1,640	1,087.1	1,740	1,091.7

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2004 - 2005

	2004		2005	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	64,540	7,684.6	63,800	7,376.9
Sacral Nerve Stimulation	2,560	304.8	2,460	284.4
Age	-	-	-	-
65-69	460	284.9	720	437.1
70-74	900	532.9	620	358.9
75-79	680	360.3	420	219.9
80-84	300	184.7	560	326.9
85+	220	139.0	140	84.8
Race	-	-	-	-
White	2,560	344.9	2,320	302.5
Black	0	0.0	20	40.3
Hispanic	0	0.0	20	126.9
Asian	0	0.0	0	0.0
Other	0	0.0	100	529.1
Unknown	0	0.0	0	0.0
Region	-	-	-	-
Northeast	520	344.8	400	264.1
South	980	302.2	1,120	331.9
Midwest	480	224.4	640	295.4
West	580	384.5	300	188.2

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2004 - 2005

	2004		2005	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	64,540	7,684.6	63,800	7,376.9
Other Procedures	10,540	1,255.0	10,000	1,156.3
Age	-	-	-	-
65-69	3,620	2,242.0	3,500	2,124.6
70-74	2,880	1,705.2	2,760	1,597.8
75-79	2,440	1,292.9	2,320	1,214.8
80-84	1,200	738.6	960	560.4
85+	400	252.7	460	278.7
Race	-	-	-	-
White	9,840	1,325.5	9,280	1,209.8
Black	340	684.1	200	403.1
Hispanic	140	1,011.6	240	1,522.8
Asian	40	386.1	60	600.0
Other	100	495.0	40	211.6
Unknown	80	2,339.2	180	5,084.7
Region	-	-	-	-
Northeast	1,240	822.3	1,140	752.8
South	4,880	1,504.8	4,180	1,238.8
Midwest	2,640	1,234.2	2,760	1,274.1
West	1,780	1,179.9	1,920	1,204.7

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2006 - 2007

	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	69,860	8,075.6	71,260	8,500.7
All Procedures	69,860	8,075.6	71,260	8,500.7
Age	-	-	-	-
65-69	20,540	12,095.2	20,600	12,768.1
70-74	19,780	11,669.6	19,200	11,599.8
75-79	15,640	8,378.0	15,840	9,085.7
80-84	9,980	5,930.6	10,380	6,351.0
85+	3,920	2,295.1	5,240	3,017.7
Race	-	-	-	-
White	65,920	8,532.4	67,320	8,935.5
Black	1,440	3,013.8	1,500	3,498.1
Hispanic	860	5,548.4	680	4,715.7
Asian	460	4,883.2	520	5,383.0
Other	640	3,760.3	720	4,699.7
Unknown	540	19,424.5	520	20,000.0
Region	-	-	-	-
Northeast	9,740	6,325.5	11,120	7,315.8
South	30,180	8,882.2	31,880	9,662.9
Midwest	18,420	8,415.6	16,980	8,134.5
West	11,520	7,557.1	11,280	7,641.2

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2006 - 2007

	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	69,860	8,075.6	71,260	8,500.7
Pubovaginal Sling	43,780	5,060.8	43,360	5,172.5
Age	-	-	-	-
65-69	14,400	8,479.6	14,640	9,074.0
70-74	13,180	7,775.8	12,360	7,467.4
75-79	9,420	5,046.1	9,060	5,196.7
80-84	5,060	3,006.9	5,320	3,255.0
85+	1,720	1,007.0	1,980	1,140.3
Race	-	-	-	-
White	41,060	5,314.7	40,540	5,380.9
Black	1,020	2,134.8	1,020	2,378.7
Hispanic	600	3,871.0	440	3,051.3
Asian	220	2,335.5	380	3,933.7
Other	500	2,937.7	560	3,655.4
Unknown	380	13,669.1	420	16,153.8
Region	-	-	-	-
Northeast	5,820	3,779.7	6,120	4,026.3
South	19,220	5,656.6	19,660	5,959.0
Midwest	10,960	5,007.3	10,280	4,924.8
West	7,780	5,103.6	7,300	4,945.1

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2006 - 2007

	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	69,860	8,075.6	71,260	8,500.7
Collagen Injection	13,880	1,604.5	14,060	1,677.2
Age	-	-	-	-
65-69	2,640	1,554.6	2,460	1,524.7
70-74	2,940	1,734.5	2,640	1,595.0
75-79	3,380	1,810.6	3,440	1,973.2
80-84	3,120	1,854.1	3,240	1,982.4
85+	1,800	1,053.9	2,280	1,313.1
Race	-	-	-	-
White	13,500	1,747.4	13,640	1,810.5
Black	180	376.7	160	373.1
Hispanic	60	387.1	120	832.2
Asian	80	849.3	80	828.2
Other	0	0.0	40	261.1
Unknown	60	2,158.3	20	769.2
Region	-	-	-	-
Northeast	2,100	1,363.8	2,280	1,500.0
South	5,900	1,736.4	6,220	1,885.3
Midwest	4,500	2,055.9	3,600	1,724.6
West	1,380	905.3	1,960	1,327.7

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2006 - 2007

	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	69,860	8,075.6	71,260	8,500.7
Sacral Nerve Stimulation	3,300	381.5	5,380	641.8
Age	-	-	-	-
65-69	680	400.4	1,000	619.8
70-74	900	531.0	1,500	906.2
75-79	700	375.0	1,320	757.1
80-84	820	487.3	900	550.7
85+	200	117.1	660	380.1
Race	-	-	-	-
White	3,100	401.3	5,220	692.9
Black	40	83.7	100	233.2
Hispanic	40	258.1	40	277.4
Asian	60	636.9	0	0.0
Other	60	352.5	0	0.0
Unknown	0	0.0	20	769.2
Region	-	-	-	-
Northeast	360	233.8	1,240	815.8
South	1,400	412.0	2,260	685.0
Midwest	980	447.7	1,260	603.6
West	560	367.4	620	420.0

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2006 - 2007

	2006		2007	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	69,860	8,075.6	71,260	8,500.7
Other Procedures	8,900	1,028.8	8,460	1,009.2
Age	-	-	-	-
65-69	2,820	1,660.6	2,500	1,549.5
70-74	2,760	1,628.3	2,700	1,631.2
75-79	2,140	1,146.3	2,020	1,158.7
80-84	980	582.4	920	562.9
85+	200	117.1	320	184.3
Race	-	-	-	-
White	8,260	1,069.1	7,920	1,051.2
Black	200	418.6	220	513.1
Hispanic	160	1,032.3	80	554.8
Asian	100	1,061.6	60	621.1
Other	80	470.0	120	783.3
Unknown	100	3,597.1	60	2,307.7
Region	-	-	-	-
Northeast	1,460	948.2	1,480	973.7
South	3,660	1,077.2	3,740	1,133.6
Midwest	1,980	904.6	1,840	881.5
West	1,800	1,180.8	1,400	948.4

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2008 - 2009

	2008		2009	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	70,680	8,321.4	74,300	8,369.0
All Procedures	70,680	8,321.4	74,300	8,369.0
Age	-	-	-	-
65-69	22,160	13,212.5	24,340	13,749.9
70-74	18,560	10,876.7	19,560	11,075.9
75-79	15,240	9,002.8	15,280	8,513.5
80-84	9,820	5,966.7	9,720	5,755.6
85+	4,900	2,765.9	5,400	2,906.0
Race	-	-	-	-
White	66,640	8,750.2	70,480	8,880.4
Black	1,460	3,279.4	1,580	3,215.3
Hispanic	1,040	7,439.2	860	5,555.6
Asian	400	3,649.6	380	3,231.3
Other	660	4,084.2	580	3,727.5
Unknown	480	22,018.3	420	19,090.9
Region	-	-	-	-
Northeast	10,600	7,001.3	10,620	6,805.1
South	32,220	9,487.6	34,360	9,680.5
Midwest	17,140	8,219.8	17,680	8,152.7
West	10,720	7,153.3	11,640	7,277.7

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2008 - 2009

	2008		2009	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	70,680	8,321.4	74,300	8,369.0
Pubovaginal Sling	44,080	5,189.7	45,460	5,120.5
Age	-	-	-	-
65-69	15,420	9,193.9	16,360	9,241.9
70-74	12,760	7,477.7	12,900	7,304.6
75-79	9,340	5,517.5	9,000	5,014.5
80-84	4,580	2,782.8	4,900	2,901.5
85+	1,980	1,117.6	2,300	1,237.8
Race	-	-	-	-
White	41,320	5,425.6	43,100	5,430.5
Black	920	2,066.5	960	1,953.6
Hispanic	780	5,579.4	560	3,617.6
Asian	260	2,372.3	260	2,210.9
Other	420	2,599.0	320	2,056.6
Unknown	380	17,431.2	260	11,818.2
Region	-	-	-	-
Northeast	6,380	4,214.0	6,220	3,985.6
South	20,180	5,942.3	21,280	5,995.4
Midwest	10,640	5,102.6	10,420	4,804.9
West	6,880	4,591.0	7,540	4,714.3

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2008 - 2009

	2008		2009	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	70,680	8,321.4	74,300	8,369.0
Collagen Injection	12,320	1,450.5	10,820	1,218.7
Age	-	-	-	-
65-69	2,520	1,502.5	2,320	1,310.6
70-74	2,260	1,324.4	2,240	1,268.4
75-79	2,520	1,488.7	2,580	1,437.5
80-84	3,200	1,944.3	1,840	1,089.5
85+	1,820	1,027.3	1,840	990.2
Race	-	-	-	-
White	11,920	1,565.2	10,540	1,328.0
Black	220	494.2	120	244.2
Hispanic	60	429.2	60	387.6
Asian	80	729.9	40	340.1
Other	20	123.8	40	257.1
Unknown	20	917.4	20	909.1
Region	-	-	-	-
Northeast	1,980	1,307.8	1,500	961.2
South	5,180	1,525.3	4,680	1,318.5
Midwest	3,380	1,620.9	3,240	1,494.1
West	1,780	1,187.8	1,400	875.3

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2008 - 2009

	2008		2009	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	70,680	8,321.4	74,300	8,369.0
Sacral Nerve Stimulation	6,640	781.7	9,180	1,034.0
Age	-	-	-	-
65-69	1,600	954.0	2,100	1,186.3
70-74	1,500	879.0	2,280	1,291.1
75-79	1,600	945.2	1,780	991.8
80-84	1,200	729.1	2,080	1,231.6
85+	740	417.7	940	505.9
Race	-	-	-	-
White	6,240	819.3	8,720	1,098.7
Black	180	404.3	140	284.9
Hispanic	0	0.0	80	516.8
Asian	40	365.0	20	170.1
Other	100	618.8	140	899.7
Unknown	80	3,669.7	80	3,636.4
Region	-	-	-	-
Northeast	960	634.1	1,460	935.5
South	3,280	965.8	4,720	1,329.8
Midwest	1,680	805.7	1,780	820.8
West	720	480.4	1,220	762.8

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2008 - 2009

	2008		2009	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	70,680	8,321.4	74,300	8,369.0
Other Procedures	7,640	899.5	8,840	995.7
Age	-	-	-	-
65-69	2,620	1,562.1	3,560	2,011.1
70-74	2,040	1,195.5	2,140	1,211.8
75-79	1,780	1,051.5	1,920	1,069.8
80-84	840	510.4	900	532.9
85+	360	203.2	320	172.2
Race	-	-	-	-
White	7,160	940.2	8,120	1,023.1
Black	140	314.5	360	732.6
Hispanic	200	1,430.6	160	1,033.6
Asian	20	182.5	60	510.2
Other	120	742.6	80	514.1
Unknown	0	0.0	60	2,727.3
Region	-	-	-	-
Northeast	1,280	845.4	1,440	922.7
South	3,580	1,054.2	3,680	1,036.8
Midwest	1,440	690.6	2,240	1,032.9
West	1,340	894.2	1,480	925.3

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2010 - 2011

	2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	79,480	8,732.3	72,280	7,920.1
All Procedures	79,480	8,732.3	72,280	7,920.1
Age	-	-	-	-
65-69	24,140	13,329.7	21,700	11,819.2
70-74	21,200	11,668.9	19,380	10,316.2
75-79	17,260	9,490.8	16,060	9,101.2
80-84	10,940	6,468.0	9,260	5,498.2
85+	5,940	3,024.4	5,880	2,995.7
Race	-	-	-	-
White	74,960	9,231.3	67,320	8,308.2
Black	2,060	4,098.7	2,100	4,036.9
Hispanic	640	3,940.9	1,060	6,424.2
Asian	480	3,592.8	480	3,287.7
Other	820	5,049.3	1,080	6,482.6
Unknown	520	25,242.7	240	9,375.0
Region	-	-	-	-
Northeast	10,920	6,745.7	9,200	5,732.1
South	37,420	10,364.5	34,780	9,569.1
Midwest	18,140	8,237.2	15,240	6,981.2
West	13,000	7,782.6	13,060	7,666.1

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2010 - 2011

	2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	79,480	8,732.3	72,280	7,920.1
Pubovaginal Sling	46,820	5,144.0	42,280	4,632.8
Age	-	-	-	-
65-69	16,220	8,956.4	14,480	7,886.7
70-74	13,520	7,441.7	12,180	6,483.6
75-79	9,540	5,245.8	8,860	5,021.0
80-84	5,020	2,968.0	4,620	2,743.1
85+	2,520	1,283.1	2,140	1,090.3
Race	-	-	-	-
White	43,880	5,403.8	39,040	4,818.1
Black	1,220	2,427.4	1,260	2,422.1
Hispanic	480	2,955.7	700	4,242.4
Asian	380	2,844.3	420	2,876.7
Other	560	3,448.3	740	4,441.8
Unknown	300	14,563.1	120	4,687.5
Region	-	-	-	-
Northeast	6,000	3,706.4	4,860	3,028.0
South	22,360	6,193.2	20,340	5,596.2
Midwest	10,440	4,740.7	9,300	4,260.2
West	8,020	4,801.2	7,780	4,566.8

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2010 - 2011

	2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	79,480	8,732.3	72,280	7,920.1
Collagen Injection	11,480	1,261.3	10,220	1,119.9
Age	-	-	-	-
65-69	2,260	1,247.9	2,080	1,132.9
70-74	2,140	1,177.9	2,280	1,213.7
75-79	3,240	1,781.6	2,200	1,246.7
80-84	2,160	1,277.0	1,780	1,056.9
85+	1,680	855.4	1,880	957.8
Race	-	-	-	-
White	11,120	1,369.4	10,000	1,234.1
Black	160	318.3	160	307.6
Hispanic	80	492.6	20	121.2
Asian	40	299.4	0	0.0
Other	40	246.3	0	0.0
Unknown	40	1,941.7	40	1,562.5
Region	-	-	-	-
Northeast	1,900	1,173.7	1,540	959.5
South	4,520	1,251.9	4,700	1,293.1
Midwest	3,200	1,453.1	1,760	806.2
West	1,860	1,113.5	2,220	1,303.1

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2010 - 2011

	2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	79,480	8,732.3	72,280	7,920.1
Sacral Nerve Stimulation	12,480	1,371.2	12,700	1,391.6
Age	-	-	-	-
65-69	2,440	1,347.3	2,840	1,546.8
70-74	3,180	1,750.3	2,740	1,458.5
75-79	2,820	1,550.6	3,520	1,994.8
80-84	2,680	1,584.5	2,020	1,199.4
85+	1,360	692.5	1,580	805.0
Race	-	-	-	-
White	11,840	1,458.1	11,860	1,463.7
Black	400	795.9	500	961.2
Hispanic	0	0.0	160	969.7
Asian	0	0.0	0	0.0
Other	140	862.1	120	720.3
Unknown	100	4,854.4	60	2,343.8
Region	-	-	-	-
Northeast	1,500	926.6	1,880	1,171.3
South	6,900	1,911.1	6,480	1,782.9
Midwest	2,700	1,226.0	2,980	1,365.1
West	1,380	826.1	1,360	798.3

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.

Table F19b - Surgical Procedures for Female Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence by Age, Race/Ethnicity, and Geographic Region, 2010 - 2011

	2010		2011	
	Count of X	Rate per 100,000 Patients	Count of X	Rate per 100,000 Patients
Total	79,480	8,732.3	72,280	7,920.1
Other Procedures	8,700	955.9	7,080	775.8
Age	-	-	-	-
65-69	3,220	1,778.0	2,300	1,252.7
70-74	2,360	1,299.0	2,180	1,160.4
75-79	1,660	912.8	1,480	838.7
80-84	1,080	638.5	840	498.8
85+	380	193.5	280	142.7
Race	-	-	-	-
White	8,120	1,000.0	6,420	792.3
Black	280	557.1	180	346.0
Hispanic	80	492.6	180	1,090.9
Asian	60	449.1	60	411.0
Other	80	492.6	220	1,320.5
Unknown	80	3,883.5	20	781.3
Region	-	-	-	-
Northeast	1,520	939.0	920	573.2
South	3,640	1,008.2	3,260	896.9
Midwest	1,800	817.4	1,200	549.7
West	1,740	1,041.7	1,700	997.9

Source: Centers for Medicare and Medicaid Services, 5% Carrier and Outpatient Files, 2002-2011

NOTE: X represents the number of surgical procedures by female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 female adult Medicare beneficiaries with a primary diagnosis of urinary incontinence.

NOTE: The designation "Other Procedures" includes Anterior Urethropexy, e.g. MMK (Marshall-Marchetti-Krantz procedure), Raz-Type Suspension, Laparoscopic Repair, Hysterectomy with Colpo Urethropexy, Pereyra Procedure, Kelly Plication, and Botox Injection.