

Table M03 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	190,600	72,339	205,020	72,246	211,260	70,453
Age	-	-	-	-	-	-
65-69	30,340	75,360 ^R	30,040	70,649 ^R	34,580	71,832 ^R
70-74	40,300	74,327 [*]	44,040	75,179 [*]	44,360	72,201 [*]
75-79	48,100	73,977 [*]	49,580	70,768 [*]	50,300	70,036 [*]
80-84	41,240	74,306 [*]	44,460	72,552 [*]	44,080	70,754 [*]
85+	30,620	63,160 [*]	36,900	71,874 [*]	37,940	67,557 [*]
Race	-	-	-	-	-	-
White	165,680	71,935 ^R	177,120	72,153 ^R	180,040	70,065 ^R
Black	13,960	70,791 [*]	14,900	71,292 [*]	16,560	73,797 [*]
Hispanic	3,480	78,027 [*]	3,920	72,325 [*]	4,420	72,222 [*]
Asian	3,300	94,828 [*]	3,620	77,350 [*]	4,200	67,308 [*]
Other	3,660	77,542 [*]	4,480	69,782 [*]	5,180	75,072 [*]
Unknown	520	66,667	980	111,364 [*]	860	71,667
Region	-	-	-	-	-	-
Northeast	37,520	69,507	40,760	71,811	41,340	69,107
South	73,100	75,005 [*]	74,300	71,899	79,400	72,525 [*]
Midwest	46,220	68,923 ^R	49,680	70,011 ^R	48,200	66,009 ^R
West	33,760	75,056 [*]	40,280	76,404 [*]	42,320	73,549 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table M03 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	225,960	71,670	231,200	71,922	218,780	72,602
Age	-	-	-	-	-	-
65-69	37,060	73,068 ^R	38,500	74,067 ^R	35,920	76,785 ^R
70-74	46,200	73,101 *	46,700	75,787 *	45,240	75,350 *
75-79	55,200	73,639 *	55,060	74,973 *	52,140	76,295 *
80-84	46,220	69,692 *	47,800	68,169 *	45,800	70,310 *
85+	41,280	68,708 *	43,140	67,092 *	39,680	65,007 *
Race	-	-	-	-	-	-
White	194,900	72,233 ^R	201,240	72,399 ^R	190,660	72,456 ^R
Black	16,440	66,130 *	16,920	66,405 *	16,220	73,195 *
Hispanic	5,140	74,277 *	4,560	74,754 *	3,460	62,909
Asian	4,280	67,085 *	3,740	76,016 *	3,940	86,784 *
Other	4,520	70,186 *	4,140	67,427 *	4,000	74,349 *
Unknown	680	79,070	600	69,767	500	80,645
Region	-	-	-	-	-	-
Northeast	43,900	70,944	47,220	74,315 *	43,780	72,652 *
South	85,340	73,760 *	92,080	73,865 *	87,180	74,360 *
Midwest	51,480	67,790 ^R	51,800	66,684 ^R	47,700	68,123 ^R
West	45,240	73,251 *	40,100	72,148 *	40,120	74,545 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table M03 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	227,480	72,850	237,300	73,087	245,220	73,942
Age	-	-	-	-	-	-
65-69	39,820	78,914 ^R	41,020	76,444 ^R	43,420	78,460 ^R
70-74	50,060	77,087 *	51,380	79,560 *	52,780	80,310 *
75-79	50,180	74,165 *	52,820	74,880 *	52,480	74,166 *
80-84	46,780	69,925 *	48,100	69,650 *	50,460	72,024 *
85+	40,640	65,233 *	43,980	65,799 *	46,080	66,055 *
Race	-	-	-	-	-	-
White	198,060	72,465 ^R	206,740	73,162 ^R	210,780	73,371 ^R
Black	15,560	69,217 *	15,960	67,742 *	18,380	75,266 *
Hispanic	4,020	79,447 *	4,300	70,033 *	5,000	76,220 *
Asian	4,140	87,342 *	4,360	75,958 *	5,540	86,834 *
Other	5,100	85,859 *	4,900	84,775 *	4,960	82,943 *
Unknown	600	83,333	1,040	118,182	560	54,902
Region	-	-	-	-	-	-
Northeast	47,380	73,824 *	46,540	71,821 *	46,220	70,371 *
South	89,220	74,029 *	96,440	77,103 *	101,800	79,631 *
Midwest	49,300	67,627 ^R	48,660	64,759 ^R	49,900	64,704 ^R
West	41,580	76,070 *	45,660	76,534 *	47,300	77,541 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

Table M03 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	253,140	74,960
Age	-	-
65-69	44,660	79,693 ^R
70-74	56,480	80,456 *
75-79	54,660	76,921 *
80-84	47,480	70,154 *
85+	49,860	68,564 *
Race	-	-
White	217,700	74,888 ^R
Black	19,020	73,607 *
Hispanic	4,920	75,229 *
Asian	5,120	74,636 *
Other	5,420	84,688 *
Unknown	960	70,588
Region	-	-
Northeast	46,560	70,163 *
South	108,960	82,696 *
Midwest	47,560	63,481 ^R
West	50,060	77,420 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of stress urinary incontinence.