

Table M05 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Other Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	6,360	2,414	6,600	2,326	7,020	2,341
Age	-	-	-	-	-	-
65-69	900	2,236 ^R	1,020	2,399 ^R	920	1,911 ^R
70-74	880	1,623 [*]	1,100	1,878 [*]	1,240	2,018 [*]
75-79	1,960	3,015 [*]	1,540	2,198 [*]	1,600	2,228 [*]
80-84	1,400	2,523 [*]	1,980	3,231 [*]	1,620	2,600 [*]
85+	1,220	2,517 [*]	960	1,870 [*]	1,640	2,920 [*]
Race	-	-	-	-	-	-
White	5,420	2,353 ^R	5,720	2,330 ^R	6,240	2,428 ^R
Black	480	2,434 [*]	340	1,627 [*]	300	1,337 [*]
Hispanic	160	3,587 [*]	340	6,273 [*]	140	2,288 [*]
Asian	60	1,724 [*]	60	1,282 [*]	140	2,244 [*]
Other	200	4,237 [*]	80	1,246 [*]	140	2,029 [*]
Unknown	40	5,128 [*]	60	6,818 [*]	60	5,000 [*]
Region	-	-	-	-	-	-
Northeast	1,580	2,927	1,840	3,242	2,080	3,477
South	2,240	2,298	2,080	2,013	1,780	1,626 [*]
Midwest	1,480	2,207 ^R	1,620	2,283 ^R	1,900	2,602 ^R
West	1,060	2,357 [*]	1,060	2,011 [*]	1,260	2,190 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

Table M05 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Other Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	6,480	2,055	6,380	1,985	5,820	1,931
Age	-	-	-	-	-	-
65-69	1,020	2,011 ^R	1,040	2,001 ^R	980	2,095 ^R
70-74	1,040	1,646 [*]	1,080	1,753 [*]	920	1,532 [*]
75-79	1,880	2,508 [*]	1,200	1,634 [*]	1,560	2,283 [*]
80-84	1,380	2,081 [*]	1,680	2,396 [*]	1,340	2,057 [*]
85+	1,160	1,931 [*]	1,380	2,146 [*]	1,020	1,671 [*]
Race	-	-	-	-	-	-
White	5,580	2,068 ^R	5,200	1,871 ^R	5,020	1,908 ^R
Black	300	1,207 [*]	580	2,276 [*]	280	1,264 [*]
Hispanic	140	2,023 [*]	120	1,967 [*]	120	2,182 [*]
Asian	220	3,448 [*]	160	3,252 [*]	180	3,965 [*]
Other	180	2,795 [*]	300	4,886 [*]	160	2,974 [*]
Unknown	60	6,977 [*]	20	2,326 [*]	60	9,677 [*]
Region	-	-	-	-	-	-
Northeast	1,760	2,844	1,220	1,920	1,000	1,660
South	1,880	1,625 [*]	2,360	1,893	2,180	1,859
Midwest	1,600	2,107 ^R	1,500	1,931 ^R	1,320	1,885 ^R
West	1,240	2,008 [*]	1,300	2,339 [*]	1,320	2,453 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

Table M05 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Other Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	6,160	1,973	5,740	1,768	6,320	1,906
Age	-	-	-	-	-	-
65-69	1,000	1,982 ^R	940	1,752 ^R	920	1,663 ^R
70-74	1,020	1,571 [*]	900	1,394 [*]	1,100	1,674 [*]
75-79	1,220	1,803 [*]	1,120	1,588 [*]	1,520	2,148 [*]
80-84	1,600	2,392 [*]	1,400	2,027 [*]	980	1,399 [*]
85+	1,320	2,119 [*]	1,380	2,065 [*]	1,800	2,580 [*]
Race	-	-	-	-	-	-
White	5,400	1,976 ^R	5,080	1,798 ^R	5,620	1,956 ^R
Black	400	1,779 [*]	360	1,528 [*]	380	1,556 [*]
Hispanic	160	3,162 [*]	200	3,257 [*]	40	610 [*]
Asian	0	0 [*]	40	697 [*]	160	2,508 [*]
Other	180	3,030 [*]	60	1,038 [*]	100	1,672 [*]
Unknown	20	2,778 [*]	0	0 [*]	20	1,961 [*]
Region	-	-	-	-	-	-
Northeast	1,020	1,589	840	1,296 [*]	1,340	2,040
South	2,420	2,008	2,000	1,599 [*]	2,060	1,611 [*]
Midwest	1,260	1,728 ^R	1,700	2,262 ^R	1,580	2,049 ^R
West	1,460	2,671 [*]	1,200	2,011 [*]	1,340	2,197 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

Table M05 - Physician Office Visits by Male Medicare Beneficiaries with Any Diagnosis of Other Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	7,060	2,091
Age	-	-
65-69	1,080	1,927 ^R
70-74	1,240	1,766 [*]
75-79	1,220	1,717 [*]
80-84	1,540	2,275 [*]
85+	1,980	2,723 [*]
Race	-	-
White	6,360	2,188 ^R
Black	280	1,084 [*]
Hispanic	160	2,447 [*]
Asian	140	2,041 [*]
Other	120	1,875 [*]
Unknown	0	0 [*]
Region	-	-
Northeast	1,020	1,537 [*]
South	2,420	1,837 [*]
Midwest	1,800	2,403 ^R
West	1,820	2,815 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had a physician office visit for any diagnosis of other urinary incontinence.