

Table M06 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
<b>Total</b>	40,940	15,538	40,420	14,243	44,140	14,720
<b>Age</b>	-	-	-	-	-	-
65-69	7,400	18,381 <sup>R</sup>	7,240	17,027 <sup>R</sup>	7,740	16,078 <sup>R</sup>
70-74	9,420	17,374 <sup>*</sup>	8,400	14,339 <sup>*</sup>	9,760	15,885 <sup>*</sup>
75-79	9,340	14,365 <sup>*</sup>	9,660	13,788 <sup>*</sup>	10,480	14,592 <sup>*</sup>
80-84	8,020	14,450 <sup>*</sup>	8,000	13,055 <sup>*</sup>	8,340	13,387 <sup>*</sup>
85+	6,760	13,944 <sup>*</sup>	7,120	13,868 <sup>*</sup>	7,820	13,925 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	34,640	15,040 <sup>R</sup>	33,940	13,826 <sup>R</sup>	37,200	14,477 <sup>R</sup>
Black	4,440	22,515 <sup>*</sup>	4,360	20,861 <sup>*</sup>	4,340	19,340 <sup>*</sup>
Hispanic	640	14,350 <sup>*</sup>	600	11,070 <sup>*</sup>	780	12,745 <sup>*</sup>
Asian	300	8,621 <sup>*</sup>	280	5,983 <sup>*</sup>	440	7,051 <sup>*</sup>
Other	760	16,102 <sup>*</sup>	1,140	17,757 <sup>*</sup>	1,200	17,391 <sup>*</sup>
Unknown	160	20,513 <sup>*</sup>	100	11,364 <sup>*</sup>	180	15,000 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	6,940	12,857 <sup>*</sup>	7,940	13,989 <sup>*</sup>	8,180	13,674 <sup>*</sup>
South	14,060	14,426 <sup>*</sup>	13,500	13,064 <sup>*</sup>	13,620	12,441 <sup>*</sup>
Midwest	13,580	20,251 <sup>R</sup>	11,620	16,375 <sup>R</sup>	14,100	19,310 <sup>R</sup>
West	6,360	14,140 <sup>*</sup>	7,360	13,961 <sup>*</sup>	8,240	14,320 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
<b>Total</b>	48,960	15,529	52,900	16,456	49,840	16,539
<b>Age</b>	-	-	-	-	-	-
65-69	8,740	17,232 <sup>R</sup>	8,960	17,237 <sup>R</sup>	9,900	21,163 <sup>R</sup>
70-74	10,180	16,108 *	11,280	18,306 *	9,840	16,389 *
75-79	11,780	15,715 *	11,820	16,095 *	10,500	15,364 *
80-84	9,640	14,536 *	10,760	15,345 *	10,660	16,365 *
85+	8,620	14,348 *	10,080	15,677 *	8,940	14,646 *
<b>Race</b>	-	-	-	-	-	-
White	42,020	15,573 <sup>R</sup>	45,340	16,312 <sup>R</sup>	42,440	16,128 <sup>R</sup>
Black	4,400	17,699 *	4,960	19,466 *	4,480	20,217 *
Hispanic	840	12,139 *	880	14,426 *	780	14,182 *
Asian	580	9,091 *	600	12,195 *	740	16,300 *
Other	1,020	15,839 *	1,040	16,938 *	1,320	24,535 *
Unknown	100	11,628 *	80	9,302 *	80	12,903 *
<b>Region</b>	-	-	-	-	-	-
Northeast	9,180	14,835 *	10,600	16,682 *	10,300	17,093 *
South	14,940	12,913 *	15,520	12,450 *	15,740	13,425 *
Midwest	15,780	20,780 <sup>R</sup>	17,660	22,734 <sup>R</sup>	14,540	20,765 <sup>R</sup>
West	9,060	14,670 *	9,120	16,409 *	9,260	17,205 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

\* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
<b>Total</b>	54,400	17,421	57,800	17,802	59,440	17,923
<b>Age</b>	-	-	-	-	-	-
65-69	9,880	19,580 <sup>R</sup>	10,300	19,195 <sup>R</sup>	12,240	22,118 <sup>R</sup>
70-74	12,080	18,602 <sup>*</sup>	12,380	19,170 <sup>*</sup>	12,820	19,507 <sup>*</sup>
75-79	11,560	17,085 <sup>*</sup>	12,980	18,401 <sup>*</sup>	11,840	16,733 <sup>*</sup>
80-84	10,940	16,353 <sup>*</sup>	11,220	16,247 <sup>*</sup>	11,780	16,814 <sup>*</sup>
85+	9,940	15,955 <sup>*</sup>	10,920	16,338 <sup>*</sup>	10,760	15,424 <sup>*</sup>
<b>Race</b>	-	-	-	-	-	-
White	47,720	17,459 <sup>R</sup>	49,900	17,659 <sup>R</sup>	51,180	17,815 <sup>R</sup>
Black	3,840	17,082 <sup>*</sup>	4,920	20,883 <sup>*</sup>	5,160	21,130 <sup>*</sup>
Hispanic	1,060	20,949 <sup>*</sup>	860	14,007 <sup>*</sup>	1,200	18,293 <sup>*</sup>
Asian	540	11,392 <sup>*</sup>	740	12,892 <sup>*</sup>	600	9,404 <sup>*</sup>
Other	1,200	20,202 <sup>*</sup>	1,160	20,069 <sup>*</sup>	1,060	17,726 <sup>*</sup>
Unknown	40	5,556 <sup>*</sup>	220	25,000 <sup>*</sup>	240	23,529 <sup>*</sup>
<b>Region</b>	-	-	-	-	-	-
Northeast	10,780	16,797 <sup>*</sup>	11,520	17,778 <sup>*</sup>	11,620	17,692 <sup>*</sup>
South	16,820	13,956 <sup>*</sup>	17,820	14,247 <sup>*</sup>	18,480	14,456 <sup>*</sup>
Midwest	15,900	21,811 <sup>R</sup>	17,080	22,731 <sup>R</sup>	18,340	23,781 <sup>R</sup>
West	10,900	19,941 <sup>*</sup>	11,380	19,075 <sup>*</sup>	11,000	18,033 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

Table M06 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
<b>Total</b>	67,520	19,994
<b>Age</b>	-	-
65-69	12,820	22,877 <sup>R</sup>
70-74	15,260	21,738 <sup>*</sup>
75-79	14,080	19,814 <sup>*</sup>
80-84	12,940	19,119 <sup>*</sup>
85+	12,420	17,079 <sup>*</sup>
<b>Race</b>	-	-
White	57,360	19,732 <sup>R</sup>
Black	5,980	23,142 <sup>*</sup>
Hispanic	1,260	19,266 <sup>*</sup>
Asian	960	13,994 <sup>*</sup>
Other	1,780	27,813 <sup>*</sup>
Unknown	180	13,235 <sup>*</sup>
<b>Region</b>	-	-
Northeast	13,060	19,681 <sup>*</sup>
South	20,560	15,604 <sup>*</sup>
Midwest	20,740	27,683 <sup>R</sup>
West	13,160	20,353 <sup>*</sup>

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

<sup>R</sup> Indicates the reference group

<sup>\*</sup> Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of urinary incontinence.