

Table M07 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	33,700	12,790	33,280	11,727	36,340	12,119
Age	-	-	-	-	-	-
65-69	5,900	14,655 ^R	5,940	13,970 ^R	6,440	13,378 ^R
70-74	7,920	14,607 *	7,060	12,052 *	7,820	12,728 *
75-79	7,700	11,843 *	7,840	11,190 *	8,960	12,476 *
80-84	6,380	11,495 *	6,660	10,868 *	6,740	10,819 *
85+	5,800	11,964 *	5,780	11,258 *	6,380	11,360 *
Race	-	-	-	-	-	-
White	28,580	12,409 ^R	27,780	11,317 ^R	30,520	11,877 ^R
Black	3,640	18,458 *	3,720	17,799 *	3,700	16,488 *
Hispanic	540	12,108 *	540	9,963 *	640	10,458 *
Asian	220	6,322 *	200	4,274 *	380	6,090 *
Other	580	12,288 *	940	14,642 *	920	13,333 *
Unknown	140	17,949 *	100	11,364 *	180	15,000 *
Region	-	-	-	-	-	-
Northeast	5,320	9,856 *	6,000	10,571 *	6,620	11,067 *
South	11,580	11,882 *	11,480	11,109 *	11,100	10,139 *
Midwest	11,380	16,970 ^R	9,600	13,529 ^R	11,840	16,215 ^R
West	5,420	12,050 *	6,200	11,760 *	6,780	11,783 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	40,340	12,795	43,160	13,426	40,580	13,467
Age	-	-	-	-	-	-
65-69	7,000	13,801 ^R	7,360	14,159 ^R	8,280	17,700 ^R
70-74	8,340	13,196 [*]	9,320	15,125 [*]	7,940	13,225 [*]
75-79	10,020	13,367 [*]	9,460	12,881 [*]	8,400	12,291 [*]
80-84	7,820	11,791 [*]	8,680	12,379 [*]	8,500	13,049 [*]
85+	7,160	11,917 [*]	8,340	12,970 [*]	7,460	12,221 [*]
Race	-	-	-	-	-	-
White	34,560	12,809 ^R	36,880	13,268 ^R	34,500	13,111 ^R
Black	3,640	14,642 [*]	4,000	15,699 [*]	3,600	16,245 [*]
Hispanic	720	10,405 [*]	760	12,459 [*]	680	12,364 [*]
Asian	520	8,151 [*]	460	9,350 [*]	600	13,216 [*]
Other	800	12,422 [*]	980	15,961 [*]	1,140	21,190 [*]
Unknown	100	11,628 [*]	80	9,302 [*]	60	9,677 [*]
Region	-	-	-	-	-	-
Northeast	7,040	11,377 [*]	7,900	12,433 [*]	7,800	12,944 [*]
South	12,520	10,821 [*]	13,100	10,509 [*]	13,040	11,122 [*]
Midwest	13,080	17,224 ^R	14,540	18,718 ^R	12,020	17,167 ^R
West	7,700	12,468 [*]	7,620	13,710 [*]	7,720	14,344 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	44,040	14,104	46,800	14,414	47,960	14,461
Age	-	-	-	-	-	-
65-69	8,240	16,330 ^R	8,820	16,437 ^R	10,380	18,757 ^R
70-74	9,780	15,060 [*]	9,680	14,989 [*]	10,120	15,399 [*]
75-79	9,660	14,277 [*]	10,520	14,914 [*]	9,480	13,397 [*]
80-84	8,280	12,377 [*]	8,640	12,511 [*]	9,360	13,360 [*]
85+	8,080	12,970 [*]	9,140	13,674 [*]	8,620	12,357 [*]
Race	-	-	-	-	-	-
White	38,400	14,049 ^R	40,720	14,410 ^R	41,060	14,293 ^R
Black	3,200	14,235 [*]	3,840	16,299 [*]	4,420	18,100 [*]
Hispanic	940	18,577 [*]	660	10,749 [*]	920	14,024 [*]
Asian	420	8,861 [*]	500	8,711 [*]	440	6,897 [*]
Other	1,040	17,508 [*]	960	16,609 [*]	900	15,050 [*]
Unknown	40	5,556 [*]	120	13,636 [*]	220	21,569 [*]
Region	-	-	-	-	-	-
Northeast	8,340	12,995 [*]	8,740	13,488 [*]	8,940	13,611 [*]
South	14,540	12,064 [*]	15,140	12,104 [*]	15,200	11,890 [*]
Midwest	12,620	17,311 ^R	13,820	18,392 ^R	15,120	19,606 ^R
West	8,540	15,624 [*]	9,100	15,253 [*]	8,700	14,262 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

Table M07 - Ambulatory Visits (Other than Physician Visits) by Male Medicare Beneficiaries with Any Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	52,740	15,617
Age	-	-
65-69	10,380	18,522 ^R
70-74	11,920	16,980 *
75-79	10,320	14,523 *
80-84	10,060	14,864 *
85+	10,060	13,834 *
Race	-	-
White	44,280	15,232 ^R
Black	4,840	18,731 *
Hispanic	1,140	17,431 *
Asian	820	11,953 *
Other	1,500	23,438 *
Unknown	160	11,765 *
Region	-	-
Northeast	9,320	14,045 *
South	16,080	12,204 *
Midwest	16,780	22,397 ^R
West	10,560	16,332 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who had an ambulatory visit (other than physician visits) for any diagnosis of stress urinary incontinence.