

Table M10 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence,
2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	4,600	1,746	4,280	1,508	3,880	1,294
Age	-	-	-	-	-	-
65-69	780	1,937 ^R	600	1,411 ^R	780	1,620 ^R
70-74	1,120	2,066 [*]	660	1,127 [*]	760	1,237 [*]
75-79	1,180	1,815 [*]	1,220	1,741 [*]	1,200	1,671 [*]
80-84	780	1,405 [*]	980	1,599 [*]	780	1,252 [*]
85+	740	1,526 [*]	820	1,597 [*]	360	641 [*]
Race	-	-	-	-	-	-
White	4,100	1,780 ^R	3,680	1,499 ^R	3,380	1,315 ^R
Black	440	2,231 [*]	440	2,105 [*]	380	1,693 [*]
Hispanic	40	897 [*]	140	2,583 [*]	20	327 [*]
Asian	0	0 [*]	0	0 [*]	20	321 [*]
Other	0	0 [*]	20	312 [*]	60	870 [*]
Unknown	20	2,564 [*]	0	0 [*]	20	1,667 [*]
Region	-	-	-	-	-	-
Northeast	740	1,371 [*]	780	1,374 [*]	580	970 [*]
South	2,160	2,216 [*]	1,560	1,510 [*]	1,520	1,388 [*]
Midwest	1,060	1,581 ^R	1,480	2,086 ^R	1,180	1,616 ^R
West	640	1,423 [*]	460	873 [*]	600	1,043 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	3,720	1,180	3,360	1,045	3,500	1,162
Age	-	-	-	-	-	-
65-69	560	1,104 ^R	380	731 ^R	680	1,454 ^R
70-74	1,040	1,646 [*]	820	1,331 [*]	840	1,399 [*]
75-79	800	1,067 [*]	800	1,089 [*]	820	1,200 [*]
80-84	540	814 [*]	680	970 [*]	660	1,013 [*]
85+	780	1,298 [*]	680	1,058 [*]	500	819 [*]
Race	-	-	-	-	-	-
White	3,240	1,201 ^R	3,040	1,094 ^R	2,980	1,133 ^R
Black	400	1,609 [*]	240	942 [*]	380	1,715 [*]
Hispanic	0	0 [*]	20	328 [*]	0	0 [*]
Asian	0	0 [*]	20	407 [*]	0	0 [*]
Other	60	932 [*]	20	326 [*]	80	1,487 [*]
Unknown	20	2,326 [*]	20	2,326 [*]	60	9,677 [*]
Region	-	-	-	-	-	-
Northeast	780	1,261 [*]	580	913 [*]	600	996 [*]
South	1,200	1,037 [*]	1,180	947 [*]	1,420	1,211 [*]
Midwest	1,200	1,580 ^R	1,040	1,339 ^R	1,060	1,514 ^R
West	540	874 [*]	560	1,008 [*]	420	780 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	2,880	922	2,760	850	2,660	802
Age	-	-	-	-	-	-
65-69	520	1,031 ^R	400	745 ^R	500	904 ^R
70-74	660	1,016 [*]	800	1,239 [*]	600	913 [*]
75-79	740	1,094 [*]	660	936 [*]	560	791 [*]
80-84	540	807 [*]	460	666 [*]	520	742 [*]
85+	420	674 [*]	440	658 [*]	480	688 [*]
Race	-	-	-	-	-	-
White	2,520	922 ^R	2,500	885 ^R	2,300	801 ^R
Black	280	1,246 [*]	180	764 [*]	200	819 [*]
Hispanic	0	0 [*]	20	326 [*]	20	305 [*]
Asian	20	422 [*]	40	697 [*]	40	627 [*]
Other	60	1,010 [*]	20	346 [*]	100	1,672 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	580	904 [*]	600	926 [*]	620	944
South	980	813 [*]	980	784 [*]	780	610 [*]
Midwest	940	1,289 ^R	840	1,118 ^R	720	934 ^R
West	380	695 [*]	340	570 [*]	540	885 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

Table M10 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	2,960	877
Age	-	-
65-69	480	857 ^R
70-74	640	912 [*]
75-79	780	1,098 [*]
80-84	420	621 [*]
85+	640	880 [*]
Race	-	-
White	2,440	839 ^R
Black	420	1,625 [*]
Hispanic	40	612 [*]
Asian	40	583 [*]
Other	20	313 [*]
Unknown	0	0 [*]
Region	-	-
Northeast	840	1,266
South	900	683 [*]
Midwest	880	1,175 ^R
West	340	526 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of urinary incontinence.