

Table M11 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	3,960	1,503	3,780	1,332	3,480	1,161
Age	-	-	-	-	-	-
65-69	720	1,788 ^R	480	1,129 ^R	700	1,454 ^R
70-74	980	1,808 *	640	1,093 *	680	1,107 *
75-79	1,020	1,569 *	1,060	1,513 *	1,140	1,587 *
80-84	660	1,189 *	840	1,371 *	660	1,059 *
85+	580	1,196 *	760	1,480 *	300	534 *
Race	-	-	-	-	-	-
White	3,520	1,528 ^R	3,260	1,328 ^R	3,000	1,168 ^R
Black	380	1,927 *	380	1,818 *	380	1,693 *
Hispanic	40	897 *	120	2,214 *	0	0 *
Asian	0	0 *	0	0 *	20	321 *
Other	0	0 *	20	312 *	60	870 *
Unknown	20	2,564 *	0	0 *	20	1,667 *
Region	-	-	-	-	-	-
Northeast	580	1,075 *	680	1,198 *	520	869 *
South	1,840	1,888 *	1,420	1,374 *	1,400	1,279 *
Midwest	980	1,461 ^R	1,260	1,776 ^R	1,040	1,424 ^R
West	560	1,245 *	420	797 *	520	904 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	3,280	1,040	2,940	915	3,180	1,055
Age	-	-	-	-	-	-
65-69	500	986 ^R	320	616 ^R	680	1,454 ^R
70-74	880	1,392 [*]	760	1,233 [*]	780	1,299 [*]
75-79	760	1,014 [*]	700	953 [*]	700	1,024 [*]
80-84	480	724 [*]	620	884 [*]	560	860 [*]
85+	660	1,099 [*]	540	840 [*]	460	754 [*]
Race	-	-	-	-	-	-
White	2,860	1,060 ^R	2,660	957 ^R	2,700	1,026 ^R
Black	340	1,368 [*]	200	785 [*]	340	1,534 [*]
Hispanic	0	0 [*]	20	328 [*]	0	0 [*]
Asian	0	0 [*]	20	407 [*]	0	0 [*]
Other	60	932 [*]	20	326 [*]	80	1,487 [*]
Unknown	20	2,326 [*]	20	2,326 [*]	60	9,677 [*]
Region	-	-	-	-	-	-
Northeast	680	1,099 [*]	440	693 [*]	540	896 [*]
South	1,040	899 [*]	1,020	818 [*]	1,240	1,058 [*]
Midwest	1,080	1,422 ^R	980	1,262 ^R	1,020	1,457 ^R
West	480	777 [*]	500	900 [*]	380	706 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	2,480	794	2,460	758	2,520	760
Age	-	-	-	-	-	-
65-69	480	951 ^R	360	671 ^R	500	904 ^R
70-74	560	862 [*]	680	1,053 [*]	580	883 [*]
75-79	660	976 [*]	620	879 [*]	560	791 [*]
80-84	460	688 [*]	380	550 [*]	480	685 [*]
85+	320	514 [*]	420	628 [*]	400	573 [*]
Race	-	-	-	-	-	-
White	2,120	776 ^R	2,220	786 ^R	2,160	752 ^R
Black	280	1,246 [*]	180	764 [*]	200	819 [*]
Hispanic	0	0 [*]	0	0 [*]	20	305 [*]
Asian	20	422 [*]	40	697 [*]	40	627 [*]
Other	60	1,010 [*]	20	346 [*]	100	1,672 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	480	748 [*]	520	803 [*]	560	853 [*]
South	860	714 [*]	880	704 [*]	720	563 [*]
Midwest	780	1,070 ^R	740	985 ^R	720	934 ^R
West	360	659 [*]	320	536 [*]	520	853 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

Table M11 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Stress Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	2,500	740
Age	-	-
65-69	400	714 ^R
70-74	600	855 *
75-79	600	844 *
80-84	360	532 *
85+	540	743 *
Race	-	-
White	2,060	709 ^R
Black	380	1,471 *
Hispanic	20	306 *
Asian	20	292 *
Other	20	313 *
Unknown	0	0 *
Region	-	-
Northeast	700	1,055
South	740	562 *
Midwest	740	988 ^R
West	320	495 *

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of stress urinary incontinence.