

Table M13 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2002 - 2004

	2002 Count of X	2002 Rate per 100,000 Patients	2003 Count of X	2003 Rate per 100,000 Patients	2004 Count of X	2004 Rate per 100,000 Patients
Total	200	76	80	28	80	27
Age	-	-	-	-	-	-
65-69	40	99 ^R	0	0 ^R	60	125 ^R
70-74	20	37 [*]	0	0 [*]	0	0 [*]
75-79	60	92 [*]	0	0 [*]	0	0 [*]
80-84	20	36 [*]	40	65 [*]	20	32 [*]
85+	60	124 [*]	40	78 [*]	0	0 [*]
Race	-	-	-	-	-	-
White	200	87 ^R	80	33 ^R	80	31 ^R
Black	0	0 [*]	0	0 [*]	0	0 [*]
Hispanic	0	0 [*]	0	0 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	20	37 [*]	0	0 [*]	20	33
South	100	103 [*]	40	39	20	18
Midwest	40	60 ^R	20	28 ^R	20	27 ^R
West	40	89 [*]	20	38 [*]	20	35 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2005 - 2007

	2005 Count of X	2005 Rate per 100,000 Patients	2006 Count of X	2006 Rate per 100,000 Patients	2007 Count of X	2007 Rate per 100,000 Patients
Total	60	19	80	25	100	33
Age	-	-	-	-	-	-
65-69	0	0 ^R	0	0 ^R	0	0 ^R
70-74	40	63 [*]	0	0 [*]	20	33 [*]
75-79	0	0 [*]	60	82 [*]	40	59 [*]
80-84	0	0 [*]	20	29 [*]	40	61 [*]
85+	20	33 [*]	0	0 [*]	0	0 [*]
Race	-	-	-	-	-	-
White	40	15 ^R	60	22 ^R	80	30 ^R
Black	20	81 [*]	20	79 [*]	20	90 [*]
Hispanic	0	0 [*]	0	0 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	0	0 [*]	0	0 [*]	0	0 [*]
South	20	17	20	16 [*]	100	85 [*]
Midwest	20	26 ^R	40	52 ^R	0	0 ^R
West	20	32 [*]	20	36 [*]	0	0 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2008 - 2010

	2008 Count of X	2008 Rate per 100,000 Patients	2009 Count of X	2009 Rate per 100,000 Patients	2010 Count of X	2010 Rate per 100,000 Patients
Total	40	13	100	31	60	18
Age	-	-	-	-	-	-
65-69	20	40 ^R	20	37 ^R	0	0 ^R
70-74	0	0 [*]	40	62 [*]	0	0 [*]
75-79	20	30 [*]	0	0 [*]	0	0 [*]
80-84	0	0 [*]	20	29 [*]	20	29 [*]
85+	0	0 [*]	20	30 [*]	40	57 [*]
Race	-	-	-	-	-	-
White	40	15 ^R	100	35 ^R	60	21 ^R
Black	0	0 [*]	0	0 [*]	0	0 [*]
Hispanic	0	0 [*]	0	0 [*]	0	0 [*]
Asian	0	0 [*]	0	0 [*]	0	0 [*]
Other	0	0 [*]	0	0 [*]	0	0 [*]
Unknown	0	0 [*]	0	0 [*]	0	0 [*]
Region	-	-	-	-	-	-
Northeast	0	0 [*]	20	31 [*]	20	31 [*]
South	20	17	40	32 [*]	20	16 [*]
Midwest	20	27 ^R	40	53 ^R	0	0 ^R
West	0	0 [*]	0	0 [*]	20	33 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

* Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

Table M13 - Hospitalizations by Male Medicare Beneficiaries with a Primary Diagnosis of Other Urinary Incontinence, 2011

	2011 Count of X	2011 Rate per 100,000 Patients
Total	40	12
Age	-	-
65-69	20	36 ^R
70-74	0	0 [*]
75-79	20	28 [*]
80-84	0	0 [*]
85+	0	0 [*]
Race	-	-
White	20	7 ^R
Black	20	77 [*]
Hispanic	0	0 [*]
Asian	0	0 [*]
Other	0	0 [*]
Unknown	0	0 [*]
Region	-	-
Northeast	0	0 [*]
South	40	30 [*]
Midwest	0	0 ^R
West	0	0 [*]

Source: Centers for Medicare and Medicaid Services, 5% Claims Files, 2002-2011

^R Indicates the reference group

^{*} Indicates that the rate is significantly different from the reference group

NOTE: X represents the number of male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.

NOTE: Unweighted counts have been multiplied by 20 to arrive at the number of counts for this table.

NOTE: Counts less than 600 should be interpreted with caution.

NOTE: Rate is per 100,000 male adult Medicare beneficiaries who were hospitalized for a primary diagnosis of other urinary incontinence.